

# Particle–Hole Ladders

Joel Feldman\*

Department of Mathematics  
University of British Columbia

Vancouver, B.C.

CANADA V6T 1Z2

feldman@math.ubc.ca

<http://www.math.ubc.ca/~feldman/>

Horst Knörrer, Eugene Trubowitz

Mathematik

ETH-Zentrum

CH-8092 Zürich

SWITZERLAND

knoerrer@math.ethz.ch, trub@math.ethz.ch

<http://www.math.ethz.ch/~knoerrer/>

**Abstract.** A self contained analysis demonstrates that the sum of all particle-hole ladder contributions for a two dimensional, weakly coupled fermion gas with a strictly convex Fermi curve at temperature zero is bounded. This is used in our construction of two dimensional Fermi liquids.

---


\* Research supported in part by the Natural Sciences and Engineering Research Council of Canada and the Forschungsinstitut für Mathematik, ETH Zürich

# Table of Contents

|                                                      |   |     |
|------------------------------------------------------|---|-----|
| §I Introduction                                      | p | 1   |
| Ladders in Momentum Space                            | p | 1   |
| Scales and Sectors                                   | p | 4   |
| Particle–Hole Ladders                                | p | 8   |
| Norms                                                | p | 9   |
| The Propagators                                      | p | 12  |
| Resectorization                                      | p | 13  |
| Compound Particle–Hole Ladders                       | p | 14  |
| §II Reduction to Bubble Estimates                    | p | 17  |
| Combinatorial Structure of Compound Ladders          | p | 17  |
| Spin Independence                                    | p | 21  |
| Scaled Norms                                         | p | 24  |
| Bubble and Double Bubble Bounds                      | p | 27  |
| The Infrared Limit                                   | p | 42  |
| §III Bubbles                                         | p | 46  |
| The Infrared Limit – Nonzero Transfer Momentum       | p | 77  |
| The Infrared Limit – Reduction to Factorized Cutoffs | p | 79  |
| §IV Double Bubbles                                   | p | 83  |
| <b>Appendices</b>                                    |   |     |
| §A Bounds on Propagators                             | p | 97  |
| §B Bound on the Generalized Model Bubble             | p | 102 |
| §C Sector Counting with Specified Transfer Momentum  | p | 117 |
| References                                           | p | 125 |
| Notation                                             | p | 127 |


# I. Introduction

In this paper we study the contributions of generalized particle-hole ladders to the four-point Green's function of a many fermion system. Formally, the amputated four-point Green's function,  $G_4((p_1, \sigma_1), (p_2, \sigma_2), (p_3, \sigma_3), (p_4, \sigma_4))$  with incoming particles of momenta  $p_1, p_4 \in \mathbb{R} \times \mathbb{R}^d$  and spins  $\sigma_1, \sigma_4 \in \{\uparrow, \downarrow\}$  and outgoing particles of momenta  $p_2, p_3$  and spins  $\sigma_2, \sigma_3$ , can be written as a sum of values of Feynman diagrams with four external legs. The propagator of these diagrams is  $C(k) = \frac{1}{ik_0 - e(\mathbf{k})}$ , where  $k = (k_0, \mathbf{k}) \in \mathbb{R} \times \mathbb{R}^d$  and the dispersion relation  $e(\mathbf{k})$  (into which the chemical potential has been absorbed) characterizes the independent fermion approximation. The interaction of the model determines the diagram vertices,  $V((k_1, \sigma_1), (k_2, \sigma_2), (k_3, \sigma_3), (k_4, \sigma_4))$ ,  $k_1 + k_4 = k_2 + k_3$ . Here, the incoming momenta are  $k_1, k_4$  and the outgoing momenta are  $k_2, k_3$ .



## Ladders in Momentum Space

The most important contributions to this four-point function are ladders. The contribution of the particle-hole ladder with  $\ell + 1$  rungs



is

$$\sum_{\substack{\tau_{i,1}, \tau_{i,2} \in \{\uparrow, \downarrow\} \\ i=1, \dots, \ell}} \int \frac{d^{d+1}k_1}{(2\pi)^{d+1}} \cdots \frac{d^{d+1}k_\ell}{(2\pi)^{d+1}} V((p_1, \sigma_1), (p_2, \sigma_2), (p_1 + k_1, \tau_{1,1}), (p_2 + k_1, \tau_{1,2})) C(p_1 + k_1) C(p_2 + k_1) \\ V((p_1 + k_1, \tau_{1,1}), (p_2 + k_1, \tau_{1,2}), \dots) \cdots V(\dots, (p_1 + k_\ell, \tau_{\ell,1}), (p_2 + k_\ell, \tau_{\ell,2})) \\ C(p_1 + k_\ell) C(p_2 + k_\ell) V((p_1 + k_\ell, \tau_{\ell,1}), (p_2 + k_\ell, \tau_{\ell,2}), (p_3, \sigma_3), (p_4, \sigma_4))$$

The contribution of the particle-particle ladder with  $\ell + 1$  rungs



is

$$\sum_{\substack{\tau_{i,1}, \tau_{i,2} \in \{\uparrow, \downarrow\} \\ i=1, \dots, \ell}} \int \frac{d^{d+1}k_i}{(2\pi)^{d+1}} \cdots \frac{d^{d+1}k_\ell}{(2\pi)^{d+1}} V((p_1, \sigma_1), (p_1+k_1, \tau_{1,1}), (p_4-k_1, \tau_{1,2}), (p_4, \sigma_4)) C(p_1+k_1) C(p_4-k_1) \\ V((p_1+k_1, \tau_{1,1}), \dots, (p_4-k_1, \tau_{1,2})) \cdots V(\dots, (p_1+k_\ell, \tau_{\ell,1}), (p_4-k_\ell, \tau_{\ell,2}), \dots) \\ C(p_1+k_\ell) C(p_4-k_\ell) V((p_1+k_\ell, \tau_{\ell,1}), (p_2, \sigma_2), (p_3, \sigma_3), (p_4-k_\ell, \tau_{\ell,2}))$$

Ladders with two rungs are called bubbles. The values of the bubbles with dispersion relation  $e(\mathbf{k}) = \frac{|\mathbf{k}|^2}{2m} - \mu$  and interaction  $V((p_1, \sigma_1), (p_2, \sigma_2), (p_3, \sigma_3), (p_4, \sigma_4)) = \lambda(\delta_{\sigma_1, \sigma_2} \delta_{\sigma_3, \sigma_4} - \delta_{\sigma_1, \sigma_3} \delta_{\sigma_2, \sigma_4})$  are well-known for  $d = 2, 3$  [FHN]. The particle-particle bubble has a logarithmic singularity [FKST, Proposition II.1b] at transfer momentum  $p_1 + p_4 = 0$  which is responsible for the formation of Cooper pairs and the onset of superconductivity. This singularity persists in models having dispersion relations that are symmetric about the origin, i.e.  $e(\mathbf{k}) = e(-\mathbf{k})$ . On the other hand, if  $e(\mathbf{k})$  is strongly asymmetric in the sense of Definition I.10 of [FKTf1] then the particle-particle bubble remains continuous and, in particular, bounded [FKLT1, page 297].

For the particle-hole bubble with  $d = 2$  and  $e(\mathbf{k}) = \frac{|\mathbf{k}|^2}{2m} - \mu$

$$\int_{\mathbb{R}^3} \frac{d^3k}{(2\pi)^3} C(k+p_1) C(k+p_2) = \int_{\mathbb{R}^3} \frac{d^3k}{(2\pi)^3} \frac{1}{i(k_0+t_0/2)-e(\mathbf{k}+\mathbf{t}/2)} \frac{1}{i(k_0-t_0/2)-e(\mathbf{k}-\mathbf{t}/2)} \\ = \begin{cases} -\frac{m}{2\pi} + \frac{m}{2\pi|\mathbf{t}|^2} \operatorname{Re} \sqrt{|\mathbf{t}|^2(|\mathbf{t}|^2 - 4k_F^2) - 4m^2t_0^2 - 4im t_0 |\mathbf{t}|^2} & \text{if } t_0, |\mathbf{t}| \neq 0 \text{ or } |\mathbf{t}| \geq 2k_F \\ -\frac{m}{2\pi} & \text{if } t_0 = 0 \text{ and } 0 < |\mathbf{t}| \leq 2k_F \\ 0 & \text{if } t_0 \neq 0 \text{ and } \mathbf{t} = 0 \end{cases}$$

where  $t = p_1 - p_2$  is the transfer momentum,  $k_F = \sqrt{2m\mu}$  is the radius of the Fermi surface and  $\sqrt{\phantom{x}}$  is the square root with nonnegative real part and cut along the negative real axis. See, for example, [FHN (2.22) or FKST, Proposition II.1a]. This is  $C^\infty$  on  $\{t \in \mathbb{R} \times \mathbb{R}^2 \mid t_0 \neq 0 \text{ or } |\mathbf{t}| > 2k_F\}$ , is Hölder continuous of degree 1 in a neighbourhood of any  $t$  with  $t_0 = 0$ ,  $0 < |\mathbf{t}| < 2k_F$  and is Hölder continuous of degree  $\frac{1}{2}$  in a neighbourhood of any  $t$  with  $t_0 = 0$ ,  $|\mathbf{t}| = 2k_F$ , but cannot be continuously extended to  $t = 0$ . However its restriction to  $t_0 = 0$  does have a  $C^\infty$  extension at the point  $\mathbf{t} = 0$ . The discontinuity at  $t = 0$  persists for general, even strongly asymmetric,  $e(\mathbf{k})$ . For this reason, bounds on particle-hole ladders in position space are not straight forward.

That the restriction of the particle-hole bubble to  $t_0 = 0$  does have a  $C^\infty$  extension for a large class of smooth dispersion relations may be seen by the following argument, which was shown to us by Manfred Salmhofer [S]. A generalization of this argument is used in Proposition III.27.

**Lemma I.1** *Choose a “scale parameter”  $M > 1$  and a function  $\nu \in C_0^\infty([\frac{1}{M}, 2M])$  that takes values in  $[0, 1]$ , is identically 1 on  $[\frac{2}{M}, M]$  is monotone on  $[\frac{1}{M}, \frac{2}{M}]$  and  $[M, 2M]$  and*

obeys

$$\sum_{j=0}^{\infty} \nu(M^{2j}x) = 1 \quad (\text{I.1})$$

for  $0 < x < 1$ . Set  $\nu_0^{[0,j]}(k_0) = \sum_{\ell=0}^j \nu(M^{2\ell}k_0^2)$  and let  $u(k, \mathbf{t})$  be a bounded  $C^\infty$  function with compact support in  $\mathbf{k}$  and bounded derivatives. Let  $e(\mathbf{k})$  be a  $C^\infty$  function that obeys  $\lim_{|\mathbf{k}| \rightarrow \infty} e(\mathbf{k}) = +\infty$ . Assume that the gradient of  $e(\mathbf{k})$  does not vanish on the Fermi surface  $F = \{ \mathbf{k} \in \mathbb{R}^d \mid e(\mathbf{k}) = 0 \}$ . Then

$$B(\mathbf{t}) = \lim_{j \rightarrow \infty} \int dk \frac{\nu_0^{[0,j]}(k_0)u(k, \mathbf{t})}{[ik_0 - e(\mathbf{k})][ik_0 - e(\mathbf{k} + \mathbf{t})]}$$

is  $C^\infty$  for  $\mathbf{t}$  in a neighbourhood of 0.

**Proof:** Write

$$\begin{aligned} B_j(\mathbf{t}) &= \int dk \frac{\nu_0^{[0,j]}(k_0)u(k, \mathbf{t})}{[ik_0 - e(\mathbf{k})][ik_0 - e(\mathbf{k} + \mathbf{t})]} = \int dk \frac{\nu_0^{[0,j]}(k_0)u(k, \mathbf{t})}{e(\mathbf{k}) - e(\mathbf{k} + \mathbf{t})} \left[ \frac{1}{ik_0 - e(\mathbf{k})} - \frac{1}{ik_0 - e(\mathbf{k} + \mathbf{t})} \right] \\ &= \int dk \frac{\nu_0^{[0,j]}(k_0)u(k, \mathbf{t})}{e(\mathbf{k}) - e(\mathbf{k} + \mathbf{t})} \int_0^1 ds \frac{d}{ds} \frac{1}{ik_0 - E(\mathbf{k}, \mathbf{t}, s)} = \int dk \int_0^1 ds \frac{\nu_0^{[0,j]}(k_0)u(k, \mathbf{t})}{[ik_0 - E(\mathbf{k}, \mathbf{t}, s)]^2} \end{aligned}$$

where

$$E(\mathbf{k}, \mathbf{t}, s) = se(\mathbf{k}) + (1-s)e(\mathbf{k} + \mathbf{t})$$

Make, for each fixed  $s$  and  $k_0$ , the change of variables from  $\mathbf{k}$  to  $E$  and  $d-1$  variables  $\theta$  on  $F$ . Denote by  $J(E, \mathbf{t}, \theta, s)$  the Jacobian of this change of variables and set

$$f(k_0, E, \theta, \mathbf{t}, s) = u((k_0, \mathbf{k}(E, \theta, \mathbf{t}, s)), \mathbf{t}) J(E, \theta, \mathbf{t}, s)$$

Because  $u$  has compact support in  $\mathbf{k}$ ,  $f$  vanishes unless  $|E| \leq \mathcal{E}$ , for some finite  $\mathcal{E}$ . Thus

$$B_j(\mathbf{t}) = \int_0^1 ds \int d\theta \int dk_0 \int_{-\mathcal{E}}^{\mathcal{E}} dE \frac{\nu_0^{[0,j]}(k_0)f(k_0, E, \theta, \mathbf{t}, s)}{[ik_0 - E]^2}$$

Set

$$B'_j(\mathbf{t}) = \int_0^1 ds \int d\theta \int dk_0 \int_{-\mathcal{E}}^{\mathcal{E}} dE \frac{\nu_0^{[0,j]}(k_0)f(k_0, 0, \theta, \mathbf{t}, s)}{[ik_0 - E]^2}$$

Since

$$\left| \partial_{\mathbf{t}}^\alpha \left[ \frac{\nu_0^{[0,j]}(k_0)f(k_0, E, \theta, \mathbf{t}, s)}{[ik_0 - E]^2} - \frac{\nu_0^{[0,j]}(k_0)f(k_0, 0, \theta, \mathbf{t}, s)}{[ik_0 - E]^2} \right] \right| \leq \text{const}_\alpha \frac{|E|}{k_0^2 + \mathcal{E}^2}$$

is integrable on  $\mathbb{R} \times [-\mathcal{E}, \mathcal{E}]$ ,  $\lim_{j \rightarrow \infty} B_j(\mathbf{t}) - B'_j(\mathbf{t})$  exists and is  $C^\infty$  by the Lebesgue dominated convergence theorem. So it suffices to consider

$$B'_j(\mathbf{t}) = -2\mathcal{E} \int_0^1 ds \int d\theta \int dk_0 \frac{\nu_0^{[0,j]}(k_0)f(k_0, 0, \theta, \mathbf{t}, s)}{k_0^2 + \mathcal{E}^2}$$

Since

$$\left| \partial_{\mathbf{t}}^\alpha \frac{\nu_0^{[0,j]}(k_0) f(k_0, 0, \theta, \mathbf{t}, s)}{k_0^2 + \mathcal{E}^2} \right| \leq \text{const}_\alpha \frac{1}{k_0^2 + \mathcal{E}^2}$$

is integrable on  $\mathbb{R}$ ,  $\lim_{j \rightarrow \infty} B'_j(\mathbf{t})$  exists and is  $C^\infty$  by the Lebesgue dominated convergence theorem. ■

## Scales and Sectors

In this paper, we derive position space bounds for generalized particle–hole ladders in two space dimensions as they arise in a multiscale analysis. The main result is Theorem I.20, which is used in [FKTf2], under the name Theorem D.2, to help construct a Fermi liquid. We assume that the dispersion relation  $e(\mathbf{k})$  is  $C^{r_e+3}$  for some  $r_e \geq 6$ , that its gradient does not vanish on the Fermi curve  $F = \{ \mathbf{k} \in \mathbb{R}^2 \mid e(\mathbf{k}) = 0 \}$  and that the Fermi curve is nonempty, connected, compact and strictly convex (meaning that its curvature does not vanish anywhere). We also fix the number  $r_0 \geq 6$  of derivatives in  $k_0$  that we wish to control.

We introduce scales as in [FKTf1, Definition I.2] and [FKTo2, §VIII]:

### Definition I.2

i) For  $j \geq 1$ , the  $j^{\text{th}}$  scale function on  $\mathbb{R} \times \mathbb{R}^2$  is defined as

$$\nu^{(j)}(k) = \nu(M^{2j}(k_0^2 + e(\mathbf{k})^2))$$

where  $\nu$  is the function of (I.1). It may be constructed by choosing a function  $\varphi \in C_0^\infty((-2, 2))$  that is identically one on  $[-1, 1]$  and setting  $\nu(x) = \varphi(x/M) - \varphi(Mx)$  for  $x > 0$  and zero otherwise. By construction,  $\nu^{(j)}$  is identically one on

$$\{ k = (k_0, \mathbf{k}) \in \mathbb{R} \times \mathbb{R}^2 \mid \sqrt{\frac{2}{M}} \frac{1}{M^j} \leq |ik_0 - e(\mathbf{k})| \leq \sqrt{M} \frac{1}{M^j} \}$$

The support of  $\nu^{(j)}$  is called the  $j^{\text{th}}$  shell. By construction, it is contained in

$$\{ k \in \mathbb{R} \times \mathbb{R}^2 \mid \frac{1}{\sqrt{M}} \frac{1}{M^j} \leq |ik_0 - e(\mathbf{k})| \leq \sqrt{2M} \frac{1}{M^j} \}$$

The momentum  $k$  is said to be of scale  $j$  if  $k$  lies in the  $j^{\text{th}}$  shell.

ii) For  $j \geq 1$ , set

$$\nu^{(\geq j)}(k) = \sum_{i \geq j} \nu^{(i)}(k)$$

for  $|ik_0 - e(\mathbf{k})| > 0$  and  $\nu^{(\geq j)}(k) = 1$  for  $|ik_0 - e(\mathbf{k})| = 0$ . Equivalently,  $\nu^{(\geq j)}(k) = \varphi(M^{2j-1}(k_0^2 + e(\mathbf{k})^2))$ . By construction,  $\nu^{(\geq j)}$  is identically 1 on

$$\{ k \in \mathbb{R} \times \mathbb{R}^2 \mid |ik_0 - e(\mathbf{k})| \leq \sqrt{M} \frac{1}{M^j} \}$$

The support of  $\nu^{(\geq j)}$  is called the  $j^{\text{th}}$  neighbourhood of the Fermi surface. By construction, it is contained in

$$\{ k \in \mathbb{R} \times \mathbb{R}^2 \mid |ik_0 - e(\mathbf{k})| \leq \sqrt{2M} \frac{1}{M^j} \}$$

The support of  $\varphi(M^{2j-2}(k_0^2 + e(\mathbf{k})^2))$  is called the  $j^{\text{th}}$  extended neighbourhood. It is contained in

$$\{ k \in \mathbb{R} \times \mathbb{R}^d \mid |ik_0 - e(\mathbf{k})| \leq \sqrt{2M} \frac{1}{M^j} \}$$

To estimate functions in position space and still make use of conservation of momentum, we use sectorization. See [FKTf1, Example A.1]. The following definition is also made in [FKTf2, §VI] and [FKTo3, §XII].

**Definition I.3 (Sectors and sectorizations)**

i) Let  $I$  be an interval on the Fermi surface  $F$  and  $j \geq 1$ . Then

$$s = \{ k \text{ in the } j^{\text{th}} \text{ neighbourhood} \mid \pi_F(k) \in I \}$$

is called a sector of length  $|I|$  at scale  $j$ . Here  $k \mapsto \pi_F(k)$  is a projection on the Fermi surface. Two different sectors  $s$  and  $s'$  are called neighbours if  $s' \cap s \neq \emptyset$ .

- ii) A sectorization of length  $\mathfrak{l}$  at scale  $j$  is a set  $\Sigma$  of sectors of length  $\mathfrak{l}$  at scale  $j$  that obeys
- the set  $\Sigma$  of sectors covers the Fermi surface
  - each sector in  $\Sigma$  has precisely two neighbours in  $\Sigma$ , one to its left and one to its right
  - if  $s, s' \in \Sigma$  are neighbours then  $\frac{1}{16}\mathfrak{l} \leq |s \cap s' \cap F| \leq \frac{1}{8}\mathfrak{l}$

Observe that there are at most  $2 \text{length}(F)/\mathfrak{l}$  sectors in  $\Sigma$ .

In the renormalization group map of [FKTf1] and [FKTo3], we integrate over fields whose arguments  $(x, \sigma, s)$  lie in  $\mathcal{B}^\uparrow \times \Sigma$ , where  $\mathcal{B}^\uparrow = (\mathbb{R} \times \mathbb{R}^2) \times \{\uparrow, \downarrow\}$  is the set of all “(positions, spins)”. On the other hand, we are interested in the dependence of the two and four-point functions on external momenta. To distinguish between the set of all positions and the set of all momenta, we denote by  $\mathbb{M} = \mathbb{R} \times \mathbb{R}^2$ , the set of all possible momenta. The set of all possible positions shall still be denoted  $\mathbb{R} \times \mathbb{R}^2$ . Thus the external variables  $(k, \sigma)$  lie in  $\check{\mathcal{B}}^\uparrow = \mathbb{M} \times \{\uparrow, \downarrow\}$ . In total, legs of four-legged kernels may lie in the disjoint union  $\mathfrak{Y}_\Sigma^\uparrow = \check{\mathcal{B}}^\uparrow \cup (\mathcal{B}^\uparrow \times \Sigma)$  for some sectorization  $\Sigma$ . The four-legged kernels over  $\mathfrak{Y}_\Sigma^\uparrow$  that we consider here arise in [FKTf2, §VII] as particle-hole reductions (as in Definition VII.4 of [FKTf2]) of four-legged kernels on  $\mathfrak{X}_\Sigma = \check{\mathcal{B}} \cup (\mathcal{B} \times \Sigma)$  where  $\check{\mathcal{B}} = \check{\mathcal{B}}^\uparrow \times \{0, 1\}$  and

$\mathcal{B} = \mathcal{B}^\dagger \times \{0, 1\}$  and  $\{0, 1\}$  is the set of creation/annihilation indices. Particle–hole reduction sets the creation/annihilation index to zero for legs number one and four and to one for legs number two and three. To simplify the notation in this paper, we shall eliminate the spin variables so that the legs lie in

$$\mathfrak{Y}_\Sigma = \mathbb{M} \cup ((\mathbb{R} \times \mathbb{R}^2) \times \Sigma)$$

Sometimes a four–legged kernel will have different sectorizations  $\Sigma, \Sigma'$  on its two left hand legs and on its two right hand legs. Therefore, we introduce the space

$$\mathfrak{Y}_{\Sigma, \Sigma'}^{(4)} = \mathfrak{Y}_\Sigma^2 \times \mathfrak{Y}_{\Sigma'}^2$$

Since  $\mathfrak{Y}_\Sigma$  is the disjoint union of  $\mathbb{M}$  and  $(\mathbb{R} \times \mathbb{R}^2) \times \Sigma$ , the space  $\mathfrak{Y}_{\Sigma, \Sigma'}^{(4)}$  is the disjoint union

$$\mathfrak{Y}_{\Sigma, \Sigma'}^{(4)} = \bigcup_{i_1, i_2, i_3, i_4 \in \{0, 1\}} \mathfrak{Y}_{i_1, \Sigma} \times \mathfrak{Y}_{i_2, \Sigma} \times \mathfrak{Y}_{i_3, \Sigma'} \times \mathfrak{Y}_{i_4, \Sigma'} \quad (\text{I.2})$$

where  $\mathfrak{Y}_{0, \Sigma} = \mathbb{M}$  and  $\mathfrak{Y}_{1, \Sigma} = (\mathbb{R} \times \mathbb{R}^2) \times \Sigma$ . If  $f$  is a function on  $\mathfrak{Y}_{\Sigma, \Sigma'}^{(4)}$ , we denote by  $f|_{(i_1, \dots, i_4)}$  its restriction to  $\mathfrak{Y}_{i_1, \Sigma} \times \mathfrak{Y}_{i_2, \Sigma} \times \mathfrak{Y}_{i_3, \Sigma'} \times \mathfrak{Y}_{i_4, \Sigma'}$  under the identification (I.2).

**Definition I.4 (Translation invariance)** Let  $\Sigma$  and  $\Sigma'$  be sectorizations.

i) Let  $y \in \mathfrak{Y}_\Sigma$  and  $t \in \mathbb{R} \times \mathbb{R}^2$ . We set

$$T_t y = \begin{cases} k & \text{if } y = k \in \mathbb{M} \\ (x + t, s) & \text{if } y = (x, s) \in (\mathbb{R} \times \mathbb{R}^2) \times \Sigma \end{cases}$$

ii) Let  $i_1, \dots, i_4 \in \{0, 1\}$ . A function  $f$  on  $\mathfrak{Y}_{i_1, \Sigma} \times \mathfrak{Y}_{i_2, \Sigma} \times \mathfrak{Y}_{i_3, \Sigma'} \times \mathfrak{Y}_{i_4, \Sigma'}$  is called translation invariant, if for all  $t \in \mathbb{R} \times \mathbb{R}^2$

$$f(T_t y_1, \dots, T_t y_4) = \left( \prod_{\substack{1 \leq \mu \leq 4 \\ i_\mu = 0}} e^{i(-1)^{b_\mu} \langle y_\mu, t \rangle_-} \right) f(y_1, \dots, y_4)$$

where

$$b_\mu = \begin{cases} 0 & \text{if } \mu = 1, 4 \\ 1 & \text{if } \mu = 2, 3 \end{cases} \quad (\text{I.3})$$

and  $\langle k, x \rangle_- = -k_0 x_0 + \mathbf{k}_1 \mathbf{x}_1 + \mathbf{k}_2 \mathbf{x}_2$ . This choice of  $b_\mu$  reflects our image of  $f$  as a particle–hole kernel, with first and fourth, resp. second and third, arguments being creation, resp. annihilation, arguments.

iii) A function  $f$  on  $\mathfrak{Y}_{\Sigma, \Sigma'}^{(4)}$  is translation invariant if  $f|_{(i_1, \dots, i_4)}$  is translation invariant for all  $i_1, \dots, i_4 \in \{0, 1\}$ .

A function  $f$  on  $(\mathfrak{Y}_\Sigma^\dagger)^4$  is translation invariant if  $f((\cdot, \sigma_1), (\cdot, \sigma_2), (\cdot, \sigma_3), (\cdot, \sigma_4))$  is translation invariant for all  $\sigma_1, \dots, \sigma_4 \in \{\uparrow, \downarrow\}$ .



**Definition I.5 (Fourier transform)** Let  $\Sigma, \Sigma'$  be sectorizations. Set  $\mathfrak{Y}_{2,\Sigma} = \mathbb{M} \times \Sigma$ .

i) Let  $i_1, \dots, i_4 \in \{0, 1, 2\}$  and  $1 \leq \mu \leq 4$  such that  $i_\mu = 1$ . The Fourier transform of a function  $f$  on  $\mathfrak{Y}_{i_1,\Sigma} \times \mathfrak{Y}_{i_2,\Sigma} \times \mathfrak{Y}_{i_3,\Sigma'} \times \mathfrak{Y}_{i_4,\Sigma'}$  with respect to the  $\mu^{\text{th}}$  variable is the function on  $\mathfrak{Y}_{i'_1,\Sigma} \times \mathfrak{Y}_{i'_2,\Sigma} \times \mathfrak{Y}_{i'_3,\Sigma'} \times \mathfrak{Y}_{i'_4,\Sigma'}$  with

$$i'_\nu = \begin{cases} i_\nu & \text{if } \nu \neq \mu \\ 2 & \text{if } \nu = \mu \end{cases}$$

defined by

$$(\Phi_\mu f)(y_1, \dots, y_{\mu-1}, (k, s), y_{\mu+1}, \dots, y_4) = \int e^{i(-1)^{b_\mu} \langle k, x \rangle} f(y_1, \dots, y_{\mu-1}, (x, s), y_{\mu+1}, \dots, y_4) d^3x$$

ii) Let  $i_1, \dots, i_4 \in \{0, 1\}$  with  $i_\mu = 1$  for at least one  $1 \leq \mu \leq 4$ . The total Fourier transform  $\check{f}$  of a translation invariant function  $f$  on  $\mathfrak{Y}_{i_1,\Sigma} \times \mathfrak{Y}_{i_2,\Sigma} \times \mathfrak{Y}_{i_3,\Sigma'} \times \mathfrak{Y}_{i_4,\Sigma'}$  is defined by

$$\check{f}(y_1, y_2, y_3, y_4) (2\pi)^3 \delta(k_1 - k_2 - k_3 + k_4) = \left( \prod_{\substack{1 \leq \mu \leq 4 \\ i_\mu = 1}} \Phi_\mu f \right)(y_1, y_2, y_3, y_4)$$

where  $y_\mu = k_\mu$  when  $i_\mu = 0$  and  $y_\mu = (k_\mu, s_\mu)$  when  $i_\mu = 1$ .  $\check{f}$  is defined on the set of all  $(y_1, y_2, y_3, y_4) \in \mathfrak{Y}_{2i_1,\Sigma} \times \mathfrak{Y}_{2i_2,\Sigma} \times \mathfrak{Y}_{2i_3,\Sigma'} \times \mathfrak{Y}_{2i_4,\Sigma'}$  for which  $k_1 - k_2 = k_3 - k_4$ .

**Definition I.6 (Sectorized Functions)** Let  $\Sigma$  and  $\Sigma'$  be sectorizations.

i) Let  $i_1, \dots, i_4 \in \{0, 1\}$ . A translation invariant function  $f$  on  $\mathfrak{Y}_{i_1,\Sigma} \times \mathfrak{Y}_{i_2,\Sigma} \times \mathfrak{Y}_{i_3,\Sigma'} \times \mathfrak{Y}_{i_4,\Sigma'}$  is sectorized if, for each  $1 \leq \mu \leq 4$  with  $i_\mu = 1$ , the total Fourier transform  $\check{f}(y_1, \dots, y_{\mu-1}, (k, s), y_{\mu+1}, \dots, y_4)$  vanishes unless  $k$  is in the  $j^{\text{th}}$  extended neighbourhood and  $\pi_F(k) \in s$ .

ii) A translation invariant function  $f$  on  $\mathfrak{Y}_{\Sigma,\Sigma'}^{(4)}$  is sectorized if  $f|_{(i_1, \dots, i_4)}$  is sectorized for all  $i_1, \dots, i_4 \in \{0, 1\}$ .

A translation invariant function  $f$  on  $(\mathfrak{Y}_\Sigma^\uparrow)^4$  is sectorized if  $f((\cdot, \sigma_1), (\cdot, \sigma_2), (\cdot, \sigma_3), (\cdot, \sigma_4))$  is sectorized for all  $\sigma_1, \dots, \sigma_4 \in \{\uparrow, \downarrow\}$ .

**Remark I.7** If  $f$  is a function in the space  $\check{\mathcal{F}}_{4,\Sigma}$  of Definition XIV.6 of [FKTf2] (or Definition XVI.7.iii of [FKTo3]), then its particle-hole reduction is a sectorized function on  $(\mathfrak{Y}_\Sigma^\uparrow)^4$ .

## Particle–Hole Ladders

### Definition I.8

- i) A (spin independent) propagator is a translation invariant function on  $(\mathbb{R} \times \mathbb{R}^2)^2$ . If  $A(x, x')$  is a propagator, then its transpose is  $A^t(x, x') = A(x', x)$ .
- ii) A (spin independent) bubble propagator is a translation invariant function on  $(\mathbb{R} \times \mathbb{R}^2)^4$ . If  $A$  and  $B$  are propagators, we define the bubble propagator

$$A \otimes B(x_1, x_2, x_3, x_4) = A(x_1, x_3)B(x_2, x_4)$$

We set

$$\begin{aligned} \mathcal{C}(A, B) &= (A + B) \otimes (A + B)^t - B \otimes B^t \\ &= A \otimes A^t + A \otimes B^t + B \otimes A^t \\ &= \begin{array}{c} \text{Diagram 1: } \overbrace{A} \quad \underbrace{A} \\ \text{Diagram 2: } \overbrace{A} \quad \underbrace{B} \\ \text{Diagram 3: } \overbrace{B} \quad \underbrace{A} \end{array} \end{aligned}$$

- iii) Let  $\Sigma, \Sigma', \Sigma''$  be sectorizations,  $P$  be a bubble propagator and  $F$  be a function on  $\mathfrak{Y}_{i_1, \Sigma''} \times \mathfrak{Y}_{i_2, \Sigma''} \times (\mathbb{R} \times \mathbb{R}^2)^2$ . If  $K$  is a function on  $\mathfrak{Y}_{\Sigma} \times \mathfrak{Y}_{\Sigma} \times \mathfrak{Y}_{1, \Sigma'} \times \mathfrak{Y}_{1, \Sigma'}$ , we set

$$(K \bullet P)(y_1, y_2, x_3, x_4) = \sum_{s'_1, s'_2 \in \Sigma'} \int dx'_1 dx'_2 K(y_1, y_2, (x'_1, s'_1), (x'_2, s'_2)) P(x'_1, x'_2, x_3, x_4)$$

If  $K$  is a function on  $\mathfrak{Y}_{1, \Sigma} \times \mathfrak{Y}_{1, \Sigma} \times \mathfrak{Y}_{i_3, \Sigma'} \times \mathfrak{Y}_{i_4, \Sigma'}$ , we set, when  $i_1, i_2, i_3, i_4$  are not all 0,

$$(F \bullet K)(y_1, y_2, y_3, y_4) = \sum_{s_1, s_2 \in \Sigma} \int dx_1 dx_2 F(y_1, y_2, x_1, x_2) K((x_1, s_1), (x_2, s_2), y_3, y_4)$$

and when  $i_1, i_2, i_3, i_4 = 0$ ,

$$(F \bullet K)(k_1, k_2, k_3, k_4) (2\pi)^3 \delta(k_1 - k_2 - k_3 + k_4) = \sum_{s_1, s_2 \in \Sigma} \int dx_1 dx_2 F(k_1, k_2, x_1, x_2) K((x_1, s_1), (x_2, s_2), k_3, k_4)$$

Observe that  $K \bullet P$  is a function on  $\mathfrak{Y}_{\Sigma}^2 \times (\mathbb{R} \times \mathbb{R}^2)^2$  and  $F \bullet K$  is a function on  $\mathfrak{Y}_{\Sigma, \Sigma'}^{(4)}$ . If  $K'$  is a function on  $(\mathfrak{Y}_{\Sigma}^{\uparrow})^4$  and  $F'$  is a function on  $(\mathfrak{Y}_{\Sigma}^{\uparrow})^2 \times (\mathcal{B}^{\uparrow})^2$  we set

$$(K' \bullet P)((\cdot, \sigma_1), (\cdot, \sigma_2), (\cdot, \sigma_3), (\cdot, \sigma_4)) = K'((\cdot, \sigma_1), (\cdot, \sigma_2), (\cdot, \sigma_3), (\cdot, \sigma_4)) \bullet P$$

and

$$\begin{aligned} (F' \bullet K')((\cdot, \sigma_1), (\cdot, \sigma_2), (\cdot, \sigma_3), (\cdot, \sigma_4)) \\ = \sum_{\tau_1, \tau_2 \in \{\uparrow, \downarrow\}} F'((\cdot, \sigma_1), (\cdot, \sigma_2), (\cdot, \tau_1), (\cdot, \tau_2)) \bullet K'((\cdot, \tau_1), (\cdot, \tau_2), (\cdot, \sigma_3), (\cdot, \sigma_4)) \end{aligned}$$

iv) Let  $\ell \geq 1$ . Let, for  $1 \leq i \leq \ell + 1$ ,  $\Sigma^{(i)}$ ,  $\Sigma'^{(i)}$  be sectorizations and  $K_i$  a function on  $\mathfrak{Y}_{\Sigma^{(i)}, \Sigma'^{(i)}}^{(4)}$ . Furthermore, let  $P_1, \dots, P_\ell$  be bubble propagators. The ladder with rungs  $K_1, \dots, K_{\ell+1}$  and bubble propagators  $P_1, \dots, P_\ell$  is defined to be

$$K_1 \bullet P_1 \bullet K_2 \bullet P_2 \bullet \dots \bullet K_\ell \bullet P_\ell \bullet K_{\ell+1}$$

If  $\Sigma$  is a sectorization and  $K'_1, \dots, K'_{\ell+1}$  are functions on  $(\mathfrak{Y}_\Sigma^\dagger)^4$ , the ladder with rungs  $K'_1, \dots, K'_{\ell+1}$  and bubble propagators  $P_1, \dots, P_\ell$  is defined to be

$$K'_1 \bullet P_1 \bullet K'_2 \bullet P_2 \bullet \dots \bullet K'_\ell \bullet P_\ell \bullet K'_{\ell+1}$$

**Remark I.9** We typically use  $\mathcal{C}(A, B)$  with  $A$  being the part,  $\nu^{(j)}(k)C(k)$ , of the propagator,  $C(k)$ , having momentum in the  $j^{\text{th}}$  shell and  $B$  being the part,  $\nu^{(\geq j+1)}(k)C(k)$ , of the propagator having momentum in the  $(j+1)^{\text{st}}$  neighbourhood. The bubble propagator  $\mathcal{C}(A, B)$  always contains at least one “hard line”  $A$  and may or may not contain one “soft line”  $B$ . The latter are created by Wick ordering. See [FKTf1, §II, subsection 9].

**Remark I.10** If  $F_1, F_2$  are functions on  $(\mathfrak{X}_\Sigma)^4$  and  $A, B$  are propagators over  $\mathcal{B}$  in the sense of Definition VII.1.i of [FKTf2], then the particle-hole reduction of  $F_1 \bullet \mathcal{C}(A, B) \bullet F_2$  (with the  $\mathcal{C}(A, B)$  of Definition VII.1.i of [FKTf2]) is equal to

$$-F_1^{\text{ph}} \bullet \mathcal{C}(A((\cdot 1), (\cdot 0)), B((\cdot 1), (\cdot 0))) \bullet F_2^{\text{ph}}$$

(with the  $\mathcal{C}$  of Definition I.8) since  $B((x, \sigma, 0), (x', \sigma', 1)) = -B((\cdot 1), (\cdot 0))^t((x, \sigma), (x', \sigma'))$ .

## Norms

In the momentum space variables, we take suprema of the function and its derivatives. In the position space variables, we will apply the  $L^1$ – $L^\infty$  norm of Definition I.11, below, to the function and to the function multiplied by various coordinate differences.

**Definition I.11** Let  $f$  be a function on  $(\mathbb{R} \times \mathbb{R}^2)^n$ . Its  $L^1$ – $L^\infty$  norm is

$$\|f\|_{1,\infty} = \max_{1 \leq j_0 \leq n} \sup_{x_{j_0} \in \mathbb{R} \times \mathbb{R}^2} \int \prod_{\substack{j=1, \dots, n \\ j \neq j_0}} dx_j |f(x_1, \dots, x_n)|$$

Multiple derivatives are labeled by a multiindex  $\delta = (\delta_0, \delta_1, \delta_2) \in \mathbb{N}_0 \times \mathbb{N}_0^2$ . For such a multiindex, we set  $|\delta| = \delta_0 + \delta_1 + \delta_2$ ,  $\delta! = \delta_0! \delta_1! \delta_2!$  and  $x^\delta = x_0^{\delta_0} x_1^{\delta_1} x_2^{\delta_2}$  for  $x \in \mathbb{R} \times \mathbb{R}^2$ .

**Definition I.12** Let  $\Sigma$  be a sectorization and  $A$  a function on  $((\mathbb{R} \times \mathbb{R}^2) \times \Sigma)^2$ . For a multiindex  $\delta \in \mathbb{N}_0 \times \mathbb{N}_0^2$ , we define

$$|A|_{1,\Sigma}^\delta = \max_{i=1,2} \max_{s_i \in \Sigma} \sum_{s_{3-i} \in \Sigma} \left\| (x-y)^\delta A((x, s_1), (y, s_2)) \right\|_{1,\infty}$$

Variables for four-point functions may be momenta or position/sector pairs. Therefore we introduce differential-decay operators that differentiate momentum space variables and multiply position space variables by coordinate differences. We again use the identification

$$\mathfrak{Y}_{\Sigma, \Sigma'}^{(4)} = \bigcup_{i_1, i_2, i_3, i_4 \in \{0,1\}} \mathfrak{Y}_{i_1, \Sigma} \times \mathfrak{Y}_{i_2, \Sigma} \times \mathfrak{Y}_{i_3, \Sigma'} \times \mathfrak{Y}_{i_4, \Sigma'}$$

of (I.2).

**Definition I.13 (Differential-decay operators)** Let  $\Sigma$  and  $\Sigma'$  be sectorizations,  $\delta = (\delta_0, \delta_1, \delta_2) \in \mathbb{N}_0 \times \mathbb{N}_0^2$  a multiindex and  $\mu, \mu' \in \{1, 2, 3, 4\}$  with  $\mu \neq \mu'$ .

i) Let  $i_1, \dots, i_4 \in \{0, 1\}$  and  $f$  be a function on  $\mathfrak{Y}_{i_1, \Sigma} \times \mathfrak{Y}_{i_2, \Sigma} \times \mathfrak{Y}_{i_3, \Sigma'} \times \mathfrak{Y}_{i_4, \Sigma'}$ . If  $i_\mu = 0$ , multiplication by the  $\delta^{\text{th}}$  power of the position variable dual to  $k_\mu$  (see Definition I.5) is implemented by

$$D_\mu^\delta f(\dots, k_\mu, \dots) = (-1)^{\delta_1 + \delta_2} (-1)^{b_\mu |\delta|} i_\mu^{|\delta|} \frac{\partial^{\delta_0}}{\partial k_{\mu,0}^{\delta_0}} \frac{\partial^{\delta_1}}{\partial \mathbf{k}_{\mu,1}^{\delta_1}} \frac{\partial^{\delta_2}}{\partial \mathbf{k}_{\mu,2}^{\delta_2}} f(\dots, k_\mu, \dots)$$

In general, set

$$D_{\mu; \mu'}^\delta f = \begin{cases} (D_\mu^\delta - D_{\mu'}^\delta) f & \text{if } i_\mu = i_{\mu'} = 0 \\ (D_\mu^\delta - x_{\mu'}^\delta) f & \text{if } i_\mu = 0, i_{\mu'} = 1 \\ (x_\mu^\delta - D_{\mu'}^\delta) f & \text{if } i_\mu = 1, i_{\mu'} = 0 \\ (x_\mu^\delta - x_{\mu'}^\delta) f & \text{if } i_\mu = i_{\mu'} = 1 \end{cases}$$

Here, when  $i_\mu = 1$ , the  $\mu^{\text{th}}$  argument of  $f$  is  $(x_\mu, s_\mu)$ .

ii) If  $f$  is a function on  $\mathfrak{Y}_{\Sigma, \Sigma'}^{(4)}$ , then  $(D_{\mu; \mu'}^\delta f)|_{(i_1, \dots, i_4)} = D_{\mu; \mu'}^\delta (f|_{(i_1, \dots, i_4)})$  for all  $i_1, \dots, i_4 \in \{0, 1\}$ .

**Definition I.14** Let  $\Sigma, \Sigma'$  be sectorizations.

i) Let  $i_1, \dots, i_4 \in \{0, 1\}$  and  $f$  be a function on  $\mathfrak{Y}_{i_1, \Sigma} \times \mathfrak{Y}_{i_2, \Sigma} \times \mathfrak{Y}_{i_3, \Sigma'} \times \mathfrak{Y}_{i_4, \Sigma'}$ . For multiindices  $\delta_l, \delta_c, \delta_r \in \mathbb{N}_0 \times \mathbb{N}_0^2$ , we define

$$|f|_{\Sigma, \Sigma'}^{(\delta_l, \delta_c, \delta_r)} = \max_{\substack{s_\nu \in \Sigma \\ \nu=1,2 \\ \text{with } i_\nu=1}} \max_{\substack{s_\nu \in \Sigma' \\ \nu=3,4 \\ \text{with } i_\nu=1}} \sup_{\substack{k_\nu \in \mathbb{M} \\ \nu=1,2,3,4 \\ \text{with } i_\nu=0}} \max_{\substack{\mu=1,2 \\ \mu'=3,4}} \left\| D_{1;2}^{\delta_l} D_{\mu; \mu'}^{\delta_c} D_{3;4}^{\delta_r} f \right\|_{1,\infty}$$

Here, the  $\nu^{\text{th}}$  argument of  $f$  is  $k_\nu$  when  $i_\nu = 0$  and  $(x_\nu, s_\nu)$  when  $i_\nu = 1$ . The  $\|\cdot\|_{1,\infty}$  of Definition I.11 is applied to all spatial arguments of  $D_{1;2}^{\delta_l} D_{\mu;\mu'}^{\delta_c} D_{3;4}^{\delta_r} f$ .

ii) If  $f$  is a function on  $\mathfrak{Y}_{\Sigma,\Sigma'}^{(4)}$ , we define

$$|f|_{\Sigma,\Sigma'}^{(\delta_l,\delta_c,\delta_r)} = \sum_{i_1,i_2,i_3,i_4 \in \{0,1\}} |f|_{(i_1,\dots,i_4)}^{(\delta_l,\delta_c,\delta_r)}|_{\Sigma,\Sigma'}$$

In this Definition, the system  $(\delta_l, \delta_c, \delta_r)$  of multiindices indicates, roughly speaking, that one takes  $\delta_l$  derivatives with respect to the momentum flowing between the two left legs,  $\delta_r$  derivatives with respect to the momentum flowing between the two right legs and  $\delta_c$  derivatives with respect to momenta flowing from the left hand side to the right hand side.

In [FKTf1–f3] and [FKTo1–o4], we combine the norms of all derivatives of a function in a formal power series. We denote by  $\mathfrak{N}_3$  the set of all formal power series  $X = \sum_{\delta \in \mathbb{N}_0 \times \mathbb{N}_0^2} X_\delta t^\delta$  in the variables  $t = (t_0, t_1, t_2)$  with coefficients  $X_\delta \in \mathbb{R}_+ \cup \{\infty\}$ . See Definition V.2 of [FKTf2] or Definition II.4 of [FKTo1].

A quantity in  $\mathfrak{N}_3$  characteristic of the power counting for derivatives in scale  $j$  is

$$\mathfrak{c}_j = \sum_{\substack{\delta_1 + \delta_2 \leq r_e \\ |\delta_0| \leq r_0}} M^{j|\delta|} t^\delta + \sum_{\substack{\delta_1 + \delta_2 > r_e \\ \text{or } |\delta_0| > r_0}} \infty t^\delta \quad (\text{I.4})$$

**Definition I.15** Let  $\Sigma$  be a sectorization.

i) For a function  $A$  on  $((\mathbb{R} \times \mathbb{R}^2) \times \Sigma)^2$ , we define

$$|A|_{1,\Sigma} = \sum_{\delta \in \mathbb{N}_0 \times \mathbb{N}_0^2} \frac{1}{\delta!} |A|_{1,\Sigma}^\delta t^\delta$$

ii) For a function  $f$  on  $\mathfrak{Y}_\Sigma^4 = \mathfrak{Y}_{\Sigma,\Sigma}^{(4)}$ , we define

$$|f|_\Sigma = \sum_{\delta \in \mathbb{N}_0 \times \mathbb{N}_0^2} \frac{1}{\delta!} \left( \max_{\delta_1 + \delta_c + \delta_r = \delta} |f|_{\Sigma,\Sigma}^{(\delta_l,\delta_c,\delta_r)} \right) t^\delta$$

iii) For a function  $f$  on  $(\mathfrak{Y}_\Sigma^\uparrow)^4$ , we define

$$|f|_\Sigma = \sum_{\sigma_1, \dots, \sigma_4 \in \{\uparrow, \downarrow\}} |f((\cdot, \sigma_1), (\cdot, \sigma_2), (\cdot, \sigma_3), (\cdot, \sigma_4))|_\Sigma$$

The following Lemma, whose proof follows immediately from the various definitions and Lemma D.2.ii of [FKTo3], compares these norms with the norms of Definition VI.6 of [FKTf2].

**Lemma I.16** *Let  $\Sigma$  be a sectorization.*

i) *Let  $f$  be a sectorized, translation invariant function on  $(\mathfrak{Y}_\Sigma^\dagger)^4$  and  $V_{\text{ph}}(f)$  its particle-hole value as in Definition VII.4 of [FKTf2]. Let  $|\cdot|_{3,\Sigma}$  be the norm of Definition XIII.12 of [FKTf3] (or Definition XVI.4 of [FKTo3]). Then there is a constant  $\text{const}$ , that depends only on  $r_0$  and  $r$ , such that*

$$|V_{\text{ph}}(f)|_{3,\Sigma}^{\sim} \leq \text{const} |f|_\Sigma + \sum_{\substack{\delta_1 + \delta_2 > r \\ \text{or } \delta_0 > r_0}} \infty t^\delta$$

ii) *Let  $g$  be a function in the space  $\tilde{\mathcal{F}}_{4,\Sigma}$  of Definition XIV.6 of [FKTf2] (or Definition XVI.7.iii of [FKTo3]) and  $g^{\text{ph}}$  its particle-hole reduction as in Definition VII.4 of [FKTf2]. Then there is a universal  $\text{const}$  such that*

$$|g^{\text{ph}}|_\Sigma \leq \text{const} |g|_{3,\Sigma}^{\sim}$$

## The Propagators

The propagators we use in the multiscale analysis of [FKTf1–f3] are of the form

$$C_v^{(j)}(k) = \frac{\nu^{(j)}(k)}{ik_0 - e(\mathbf{k}) - v(k)} \quad C_v^{(\geq j)}(k) = \frac{\nu^{(\geq j)}(k)}{ik_0 - e(\mathbf{k}) - v(k)}$$

with functions  $v(k)$  satisfying  $|v(k)| \leq \frac{1}{2}|ik_0 - e(\mathbf{k})|$ . Their Fourier transforms are

$$\begin{aligned} C_v^{(j)}(x, y) &= \int \frac{d^3 k}{(2\pi)^3} e^{i\langle k, x-y \rangle} C_v^{(j)}(k) & C_v^{(\geq j)}(x, y) &= \int \frac{d^3 k}{(2\pi)^3} e^{i\langle k, x-y \rangle} C_v^{(\geq j)}(k) \\ C_v^{(j)}(y) &= \int \frac{d^3 k}{(2\pi)^3} e^{-i\langle k, y \rangle} C_v^{(j)}(k) & C_v^{(\geq j)}(y) &= \int \frac{d^3 k}{(2\pi)^3} e^{-i\langle k, y \rangle} C_v^{(\geq j)}(k) \end{aligned}$$

The function  $v(k)$  will be the sum of Fourier transforms of sectorized, translation invariant functions  $p((x, s), (x, s'))$  on  $\left((\mathbb{R} \times \mathbb{R}^2) \times \Sigma\right)^2$  for various sectorizations  $\Sigma$ . The Fourier transform of such a function is defined as

$$\check{p}(k) = \sum_{s, s' \in \Sigma} \int d^3 x e^{i\langle k, x \rangle} p((0, s), (x, s'))$$

## Resectorization

We now fix  $\frac{1}{2} < \aleph < \frac{2}{3}$  and set  $\mathfrak{l}_j = \frac{1}{M^{\aleph j}}$ . Furthermore, we select, for each  $j \geq 1$ , a sectorization  $\Sigma_j$  of length  $\mathfrak{l}_j$  at scale  $j$  and a partition of unity  $\{ \chi_s \mid s \in \Sigma_j \}$  of the  $j^{\text{th}}$  neighbourhood which fulfills Lemma XII.3 of [FKTo3] with  $\Sigma = \Sigma_j$ . The Fourier transform of  $\chi_s$  is

$$\hat{\chi}_s(x) = \int e^{-i\langle k, x \rangle} \chi_s(k) \frac{d^3 k}{(2\pi)^3}$$

**Definition I.17 (Resectorization)** Let  $j, j', j_l, j'_l, j_r, j'_r \geq 1$ .

i) Let  $p$  be a sectorized, translation invariant function on  $\left( (\mathbb{R} \times \mathbb{R}^2) \times \Sigma_j \right)^2$ . Then, for  $j' \neq j$ , the  $j'$ -resectorization of  $p$  is

$$p_{\Sigma_{j'}}((x_1, s_1), (x_2, s_2)) = \sum_{s'_1, s'_2 \in \Sigma_j} \int dx'_1 dx'_2 \hat{\chi}_{s_1}(x_1 - x'_1) p((x'_1, s'_1), (x'_2, s'_2)) \hat{\chi}_{s_2}(x'_2 - x_2)$$

It is a sectorized, translation invariant function on  $\left( (\mathbb{R} \times \mathbb{R}^2) \times \Sigma_{j'} \right)^2$ . If  $j = j'$ , we set  $p_{\Sigma_{j'}} = p$ .

ii) Let  $i_1, \dots, i_4 \in \{0, 1\}$  and  $f$  be a function on  $\mathfrak{Y}_{i_1, \Sigma_{j_l}} \times \mathfrak{Y}_{i_2, \Sigma_{j_l}} \times \mathfrak{Y}_{i_3, \Sigma_{j_r}} \times \mathfrak{Y}_{i_4, \Sigma_{j_r}}$  that is sectorized and translation invariant. Then the  $(j'_l, j'_r)$ -resectorization of  $f$  is the sectorized, translation invariant function on  $\mathfrak{Y}_{i_1, \Sigma_{j'_l}} \times \mathfrak{Y}_{i_2, \Sigma_{j'_l}} \times \mathfrak{Y}_{i_3, \Sigma_{j'_r}} \times \mathfrak{Y}_{i_4, \Sigma_{j'_r}}$  defined by

$$f_{\Sigma_{j'_l}, \Sigma_{j'_r}}(y_1, y_2, y_3, y_4) = \sum_{\substack{s'_\mu \in \Sigma_{j_l} \\ \mu \in \{1, 2\} \cap S}} \sum_{\substack{s'_\mu \in \Sigma_{j_r} \\ \mu \in \{3, 4\} \cap S}} \int \prod_{\mu \in S} \left( dx'_\mu \hat{\chi}_{s_\mu}((-1)^{b_\mu}(x_\mu - x'_\mu)) \right) f(y'_1, y'_2, y'_3, y'_4)$$

where

$$S = \{ \mu \mid i_\mu = 1 \} \cap \begin{cases} \{1, 2, 3, 4\} & \text{if } j'_l \neq j_l, j'_r \neq j_r \\ \{1, 2\} & \text{if } j'_l \neq j_l, j'_r = j_r \\ \{3, 4\} & \text{if } j'_l = j_l, j'_r \neq j_r \\ \emptyset & \text{if } j'_l = j_l, j'_r = j_r \end{cases}$$

and  $y'_\mu = y_\mu$  for  $\mu \notin S$  and, for  $\mu \in S$ ,

$$y_\mu = (x_\mu, s_\mu) \quad y'_\mu = (x'_\mu, s'_\mu)$$

iii) If  $f$  is a sectorized, translation invariant function on  $\mathfrak{Y}_{\Sigma_{j_l}, \Sigma_{j_r}}^{(4)}$ , then  $(f_{\Sigma_{j'_l}, \Sigma_{j'_r}})|_{(i_1, \dots, i_4)} = (f|_{(i_1, \dots, i_4)})_{\Sigma_{j'_l}, \Sigma_{j'_r}}$  for all  $i_1, \dots, i_4 \in \{0, 1\}$ . If  $j'_l = j'_r = j'$ , we set  $f_{\Sigma_{j'}, \Sigma_{j'}} = f_{\Sigma_{j'}, \Sigma_{j'}}$ .

iv) If  $f$  is a sectorized, translation invariant function on  $(\mathfrak{Y}_{\Sigma_j}^\uparrow)^4$ , then

$$f_{\Sigma_{j'}}((\cdot, \sigma_1), (\cdot, \sigma_2), (\cdot, \sigma_3), (\cdot, \sigma_4)) = \left( f((\cdot, \sigma_1), (\cdot, \sigma_2), (\cdot, \sigma_3), (\cdot, \sigma_4)) \right)_{\Sigma_{j'}}$$

for all  $\sigma_1, \dots, \sigma_4 \in \{\uparrow, \downarrow\}$ .

**Remark I.18** Let  $K$  and  $H$  be sectorized translation invariant functions on  $\mathfrak{Y}_{\Sigma_{i_1}, \Sigma_{j_1}}^{(4)}$  and  $\mathfrak{Y}_{\Sigma_{i_r}, \Sigma_{j_r}}^{(4)}$  respectively. Let  $P$  be a bubble propagator. If the Fourier transform

$$\int \prod_{\mu=1}^4 dx_\mu \prod_{\mu=1}^4 e^{-i(-1)^{b_\mu} \langle k_\mu, x_\mu \rangle} P(x_1, x_2, x_3, x_4)$$

of  $P$  is supported on the  $\max\{j'_1, i'_r\}^{\text{th}}$  neighbourhood, then

$$\left[ K \bullet P \bullet H \right]_{\Sigma_{i'_1}, \Sigma_{j'_r}} = K_{\Sigma_{i'_1}, \Sigma_{j'_1}} \bullet P \bullet H_{\Sigma_{i'_r}, \Sigma_{j'_r}}$$

## Compound Particle–Hole Ladders

Define, for any set  $\mathcal{Z}$  and any function  $K$  on  $\mathcal{Z}^4$ , the flipped function

$$K^f(z_1, z_2, z_3, z_4) = -K(z_1, z_3, z_2, z_4) \quad (\text{I.5})$$

**Definition I.19** Let  $\vec{F} = (F^{(2)}, F^{(3)}, \dots)$  be a sequence of sectorized, translation invariant functions  $F^{(i)}$  on  $(\mathfrak{Y}_{\Sigma_i}^\uparrow)^4$  and  $v(k)$  a function on  $\mathbb{M}$  such that  $|v(k)| \leq \frac{1}{2}|\imath k_0 - e(\mathbf{k})|$ . We define, recursively on  $0 \leq j < \infty$ , the compound particle–hole (or wrong way) ladders up to scale  $j$ , denoted by  $\mathcal{L}^{(j)} = \mathcal{L}_v^{(j)}(\vec{F})$ , as

$$\begin{aligned} \mathcal{L}^{(0)} &= 0 \\ \mathcal{L}^{(j+1)} &= \mathcal{L}_{\Sigma_j}^{(j)} + \sum_{\ell=1}^{\infty} (F + \mathcal{L}_{\Sigma_j}^{(j)} + \mathcal{L}_{\Sigma_j}^{(j)f}) \bullet \mathcal{C}^{(j)} \bullet \dots \bullet \mathcal{C}^{(j)} \bullet (F + \mathcal{L}_{\Sigma_j}^{(j)} + \mathcal{L}_{\Sigma_j}^{(j)f}) \end{aligned}$$

where  $F = \sum_{i=2}^j F_{\Sigma_j}^{(i)}$  and the  $\ell^{\text{th}}$  term has  $\ell$  bubble propagators  $\mathcal{C}^{(j)} = \mathcal{C}(C_v^{(j)}, C_v^{(\geq j+1)})$ . Observe that  $\mathcal{L}^{(1)} = \mathcal{L}^{(2)} = 0$ .



**Theorem I.20** For every  $\varepsilon > 0$  there are constants  $\rho_0, \text{const}^{(1)}$  such that the following holds. Let  $\vec{F} = (F^{(2)}, F^{(3)}, \dots)$  be a sequence of sectorized, translation invariant spin independent<sup>(2)</sup> functions  $F^{(i)}$  on  $(\mathfrak{Y}_{\Sigma_i}^\uparrow)^4$  and  $\vec{p} = (p^{(2)}, p^{(3)}, \dots)$  be a sequence of sectorized, translation invariant functions  $p^{(i)}$  on  $((\mathbb{R} \times \mathbb{R}^2) \times \Sigma_i)^2$ . Assume that there is  $\rho \leq \rho_0$  such that for  $i \geq 2$


$$|F^{(i)}|_{\Sigma_i} \leq \frac{\rho}{M^{\varepsilon i}} \mathfrak{c}_i \quad |p^{(i)}|_{1, \Sigma_i} \leq \frac{\rho \mathfrak{l}_i}{M^i} \mathfrak{c}_i \quad \check{p}^{(i)}(0, \mathbf{k}) = 0$$

Set  $v(k) = \sum_{i=2}^{\infty} \check{p}^{(i)}(k)$ . Then for all  $j \geq 1$

$$|\mathcal{L}_v^{(j+1)}(\vec{F})|_{\Sigma_j} \leq \text{const } \rho^2 \mathfrak{c}_j$$

**Remark I.21** Theorem I.20 and Theorem D.2 of [FKTf3] are equivalent. If one replaces the functions  $F^{(i)}$  of Theorem D.2 of [FKTf3] by 24 times their particle–hole reductions, then, by Corollary D.7 of [FKTf3] and Remark I.10, the concepts of compound ladders of Definition I.19 and Definition D.1 of [FKTf3] coincide. Hence Theorem I.20 and Theorem D.2 of [FKTf3] are equivalent by Lemma I.16.

Theorem I.20 will be proven following Corollary II.24. The core of the proof consists of bounds on two types of ladder fragments, that look like



and are called particle–hole bubbles and double bubbles, and a combinatorial result, Corollary II.12, that enables one to express general ladders in terms of these fragments. The most subtle part of the bound, Theorem II.19, on particle–hole bubbles is a generalization of Lemma I.1. The bound, Theorem II.20, on double bubbles also exploits “volume improvement due to overlapping loops”. A simple introduction to this phenomenon is provided at the beginning of §IV.

Ladders with external momenta have an infrared limit that behaves much like the model bubble of Lemma I.1.

---

<sup>(1)</sup> Throughout this paper we use “const” to denote unimportant constants that depend only on the dispersion relation  $e(\mathbf{k})$  and the scale parameter  $M$ . In particular, they do not depend on the scale  $j$ .

<sup>(2)</sup> “Spin independence” is formally defined in Definition II.6.

**Theorem I.22** *Under the hypotheses of Theorem I.20, the limit*

$$\mathfrak{L}(q, q', t, \sigma_1, \dots \sigma_4) = \lim_{j \rightarrow \infty} \mathcal{L}_v^{(j)}(\vec{F})|_{i_1, i_2, i_3, i_4=0}((q+\frac{t}{2}, \sigma_1), (q-\frac{t}{2}, \sigma_2), (q'+\frac{t}{2}, \sigma_3), (q'-\frac{t}{2}, \sigma_4))$$

*exists for transfer momentum  $t \neq 0$  and is continuous in  $(q, q', t)$  for  $t \neq 0$ . The restrictions to  $\mathbf{t} = 0$  and to  $t_0 = 0$ , namely,  $\mathfrak{L}(q, q', (t_0, \mathbf{0}), \sigma_1, \dots \sigma_4)$  and  $\mathfrak{L}(q, q', (0, \mathbf{t}), \sigma_1, \dots \sigma_4)$ , have continuous extensions to  $t = 0$ .*

This Theorem is proven following Lemma II.29. Notation tables are provided at the end of the paper.

## II. Reduction to Bubble Estimates

For the rest of the paper, we fix a sequence  $\vec{F} = (F^{(2)}, F^{(3)}, \dots)$  of sectorized, translation invariant, spin independent functions  $F^{(i)}$  on  $(\mathfrak{Y}_{\Sigma_i}^\uparrow)^4$  and a sequence  $\vec{p} = (p^{(2)}, p^{(3)}, \dots)$  of sectorized, translation invariant functions  $p^{(i)}$  on  $((\mathbb{R} \times \mathbb{R}^2) \times \Sigma_i)^2$  as in Theorem I.20 and we set  $v(k) = \sum_{i=2}^\infty \vec{p}^{(i)}(k)$ . Denote  $\mathcal{L}^{(j+1)} = \mathcal{L}_v^{(j+1)}(\vec{F})$  and define the particle-hole bubble propagator of scale  $j$  by

$$\mathcal{C}^{(j)} = \mathcal{C}(C_v^{(j)}, C_v^{(\geq j+1)}) = \sum_{\substack{i_1, i_2 \geq 1 \\ \min(i_1, i_2) = j}} C_v^{(i_1)} \otimes C_v^{(i_2)} t$$

and set

$$\mathcal{C}^{[j_1, j_2]} = \sum_{j=j_1}^{j_2} \mathcal{C}^{(j)} = C_v^{(\geq j_1)} \otimes C_v^{(\geq j_1)} t - C_v^{(\geq j_2+1)} \otimes C_v^{(\geq j_2+1)} t$$

### Combinatorial Structure of Compound Ladders

In this section, we use the following

**Convention II.1** Let  $K$  and  $K'$  be functions on  $(\mathfrak{Y}_{\Sigma_j})^4$  and  $(\mathfrak{Y}_{\Sigma_{j'}})^4$ , respectively. Then the notation  $K + K'$  denotes the function  $K_{\Sigma_{\max\{j, j'\}}} + K'_{\Sigma_{\max\{j, j'\}}}$  on  $(\mathfrak{Y}_{\Sigma_{\max\{j, j'\}}})^4$ . The same convention is used when  $K$  and  $K'$  are functions on  $(\mathfrak{Y}_{\Sigma_j}^\uparrow)^4$  and  $(\mathfrak{Y}_{\Sigma_{j'}}^\uparrow)^4$ .

**Definition II.2** We define, recursively on  $0 \leq j < \infty$ , sectorized, translation invariant, spin independent functions  $L^{(j)}$ , on  $(\mathfrak{Y}_{\Sigma_{j-1}}^\uparrow)^4$  by


$$L^{(0)} = L^{(1)} = L^{(2)} = 0$$

$$L^{(j+1)} = \sum_{\ell=1}^{\infty} \sum'_{\substack{i_1, \dots, i_{\ell+1} \geq 2 \\ j_1, \dots, j_\ell \geq 0}} \left[ (F^{(i_1)} + L^{(i_1)} f) \bullet \mathcal{C}^{(j_1)} \bullet \dots \bullet \mathcal{C}^{(j_\ell)} \bullet (F^{(i_{\ell+1})} + L^{(i_{\ell+1})} f) \right]_{\Sigma_j}$$

where the sum  $\sum'$  imposes the constraints

$$\begin{aligned} \max\{j_1, \dots, j_\ell\} &= j \\ i_m &\leq \min\{j_{m-1}, j_m\} \quad \text{for all } 1 \leq m \leq \ell + 1 \end{aligned}$$

When  $m = 1$ ,  $\min\{j_{m-1}, j_m\} = j_1$  and when  $m = \ell + 1$ ,  $\min\{j_{m-1}, j_m\} = j_\ell$ .



Observe that  $L^{(j)}$  depends only on the components  $F^{(2)}, \dots, F^{(j-1)}$  of  $\vec{F}$ .

### Proposition II.3

$$\begin{aligned}
i) \quad \mathcal{L}^{(j+1)} &= \sum_{i=0}^{j+1} L_{\Sigma_j}^{(i)} \\
ii) \quad \mathcal{L}^{(j+1)} &= \sum_{\ell=1}^{\infty} \sum_{i_1, \dots, i_{\ell+1}=2}^j \left[ (F^{(i_1)} + L^{(i_1)} f) \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet (F^{(i_2)} + L^{(i_2)} f) \bullet \dots \right. \\
&\quad \left. \dots \bullet \mathcal{C}^{[\max\{i_{\ell}, i_{\ell+1}\}, j]} \bullet (F^{(i_{\ell+1})} + L^{(i_{\ell+1})} f) \right]_{\Sigma_j} \\
iii) \quad L^{(j+1)} &= \left( \sum_{i=2}^j F_{\Sigma_j}^{(i)} + \mathcal{L}_{\Sigma_j}^{(j)} f + \mathcal{L}_{\Sigma_j}^{(j)} \right) \bullet \mathcal{C}^{(j)} \bullet \left( \sum_{i=2}^j F_{\Sigma_j}^{(i)} + \mathcal{L}_{\Sigma_j}^{(j)} f + \mathcal{L}^{(j+1)} \right)
\end{aligned}$$

To prove Proposition II.3, we define

$$\tilde{\mathcal{L}}^{(j+1)} = \sum_{i=0}^{j+1} L_{\Sigma_j}^{(i)}$$

and verify, in Lemmas II.4 and II.5, parts (ii) and (iii) of the Proposition, but with  $\mathcal{L}^{(k)}$  replaced by  $\tilde{\mathcal{L}}^{(k)}$ . Then we prove that  $\tilde{\mathcal{L}}^{(k)} = \mathcal{L}^{(k)}$ .

### Lemma II.4

$$\begin{aligned}
\tilde{\mathcal{L}}^{(j+1)} &= \sum_{\ell=1}^{\infty} \sum_{i_1, \dots, i_{\ell+1}=2}^j \left[ (F^{(i_1)} + L^{(i_1)} f) \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet (F^{(i_2)} + L^{(i_2)} f) \bullet \dots \right. \\
&\quad \left. \dots \bullet \mathcal{C}^{[\max\{i_{\ell}, i_{\ell+1}\}, j]} \bullet (F^{(i_{\ell+1})} + L^{(i_{\ell+1})} f) \right]_{\Sigma_j}
\end{aligned}$$

**Proof:**

$$\begin{aligned}
\tilde{\mathcal{L}}^{(j+1)} &= \sum_{\ell=1}^{\infty} \sum_{j_1, \dots, j_{\ell}=0}^j \sum_{\substack{i_m=2 \\ 1 \leq m \leq \ell+1}}^{\min\{j_{m-1}, j_m\}} \left[ (F^{(i_1)} + L^{(i_1)} f) \bullet \mathcal{C}^{(j_1)} \bullet (F^{(i_2)} + L^{(i_2)} f) \bullet \dots \right. \\
&\quad \left. \dots \bullet \mathcal{C}^{(j_{\ell})} \bullet (F^{(i_{\ell+1})} + L^{(i_{\ell+1})} f) \right]_{\Sigma_j} \\
&= \sum_{\ell=1}^{\infty} \sum_{i_1, \dots, i_{\ell+1}=2}^j \sum_{\substack{j_m=\max\{i_m, i_{m+1}\} \\ 1 \leq m \leq \ell}}^j \left[ (F^{(i_1)} + L^{(i_1)} f) \bullet \mathcal{C}^{(j_1)} \bullet (F^{(i_2)} + L^{(i_2)} f) \bullet \dots \right. \\
&\quad \left. \dots \bullet \mathcal{C}^{(j_{\ell})} \bullet (F^{(i_{\ell+1})} + L^{(i_{\ell+1})} f) \right]_{\Sigma_j}
\end{aligned}$$

■

**Lemma II.5**

$$\begin{aligned}
i) \quad L^{(j+1)} &= \left( \sum_{i=2}^j F_{\Sigma_j}^{(i)} + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} f + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} \right) \bullet \mathcal{C}^{(j)} \bullet \left( \sum_{i=2}^j F_{\Sigma_j}^{(i)} + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} f + \tilde{\mathcal{L}}_{\Sigma_j}^{(j+1)} \right) \\
ii) \quad L^{(j+1)} &= \sum_{\ell=1}^{\infty} \left[ \left( \sum_{i=2}^j F_{\Sigma_j}^{(i)} + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} f + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} \right) \bullet \mathcal{C}^{(j)} \right]^{\ell} \bullet \left( \sum_{i=2}^j F_{\Sigma_j}^{(i)} + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} f + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} \right)
\end{aligned}$$

**Proof:** i)

$$\begin{aligned}
L^{(j+1)} &= \sum_{\ell=1}^{\infty} \sum_{\ell'=1}^{\ell} \sum_{\substack{j_1, \dots, j_{\ell}=0 \\ j_1, \dots, j_{\ell'-1} \leq j-1 \\ j_{\ell'}=j}}^j \sum_{\substack{i_m=2 \\ 1 \leq m \leq \ell+1}}^{\min\{j_{m-1}, j_m\}} \left[ (F^{(i_1)} + L^{(i_1)} f) \bullet \mathcal{C}^{(j_1)} \bullet (F^{(i_2)} + L^{(i_2)} f) \bullet \dots \right. \\
&\quad \left. \dots \bullet \mathcal{C}^{(j_{\ell})} \bullet (F^{(i_{\ell+1})} + L^{(i_{\ell+1})} f) \right]_{\Sigma_j} \\
&= \text{Diagram showing a sequence of nodes } i_1, i_2, \dots, i_{\ell'}, i_{\ell'+1}, \dots, i_{\ell+1} \text{ with directed edges } j_1, j_{\ell'}=j, j_{\ell'+1} \text{ and self-loops.}
\end{aligned}$$

Splitting up the sum according to whether  $\ell' = 1$ ,  $1 < \ell' < \ell$  or  $\ell' = \ell$ , we have

$$\begin{aligned}
L^{(j+1)} &= \left[ \sum_{i=2}^j (F_{\Sigma_j}^{(i)} + L_{\Sigma_j}^{(i)} f) \right] \bullet \mathcal{C}^{(j)} \bullet \left[ \sum_{i=2}^j (F_{\Sigma_j}^{(i)} + L_{\Sigma_j}^{(i)} f) \right] \\
&+ \sum_{\ell=2}^{\infty} \sum_{\substack{j_1, \dots, j_{\ell}=0 \\ j_1=j}}^j \sum_{i=2}^j \sum_{\substack{\min\{j_{m-1}, j_m\} \\ i_m=2 \\ 2 \leq m \leq \ell+1}} (F_{\Sigma_j}^{(i)} + L_{\Sigma_j}^{(i)} f) \bullet \mathcal{C}^{(j)} \bullet \\
&\quad \left[ (F^{(i_2)} + L^{(i_2)} f) \bullet \mathcal{C}^{(j_2)} \bullet \dots \bullet \mathcal{C}^{(j_{\ell})} \bullet (F^{(i_{\ell+1})} + L^{(i_{\ell+1})} f) \right]_{\Sigma_j} \\
&+ \sum_{\ell=3}^{\infty} \sum_{\ell'=2}^{\ell-1} \left[ \sum_{\substack{j_1, \dots, j_{\ell'-1}=0 \\ i_1, \dots, i_{\ell'} \geq 2 \\ i_m \leq \min\{j_{m-1}, j_m\} \\ \text{for } m=1, \dots, \ell'-1 \\ i_{\ell'} \leq j_{\ell'-1}}}^{j-1} (F^{(i_1)} + L^{(i_1)} f) \bullet \mathcal{C}^{(j_1)} \dots (F^{(i_{\ell'})} + L^{(i_{\ell'})} f) \right]_{\Sigma_j} \bullet \mathcal{C}^{(j)} \\
&\bullet \left[ \sum_{\substack{j_{\ell'+1}, \dots, j_{\ell}=0 \\ i_{\ell'+1}, \dots, i_{\ell+1} \geq 2 \\ i_m \leq \min\{j_{m-1}, j_m\} \\ \text{for } m=\ell'+2, \dots, \ell+1 \\ i_{\ell'+1} \leq j_{\ell'+1}}}^j (F^{(i_{\ell'+1})} + L^{(i_{\ell'+1})} f) \bullet \mathcal{C}^{(j_{\ell'+1})} \dots (F^{(i_{\ell+1})} + L^{(i_{\ell+1})} f) \right]_{\Sigma_j} \\
&+ \sum_{\ell=2}^{\infty} \sum_{j_1, \dots, j_{\ell-1}=0}^{j-1} \sum_{\substack{\min\{j_{m-1}, j_m\} \\ i_m=2 \\ 1 \leq m \leq \ell}} \sum_{i=2}^j \left[ (F^{(i_1)} + L^{(i_1)} f) \bullet \mathcal{C}^{(j_1)} \dots \mathcal{C}^{(j_{\ell-1})} \bullet (F^{(i_{\ell})} + L^{(i_{\ell})} f) \right]_{\Sigma_j} \\
&\bullet \mathcal{C}^{(j)} \bullet (F_{\Sigma_j}^{(i)} + L_{\Sigma_j}^{(i)} f) \\
&= \left[ \sum_{i=2}^j (F_{\Sigma_j}^{(i)} + L_{\Sigma_j}^{(i)} f) \right] \bullet \mathcal{C}^{(j)} \bullet \left[ \sum_{i=2}^j (F_{\Sigma_j}^{(i)} + L_{\Sigma_j}^{(i)} f) \right] + \left[ \sum_{i=2}^j (F_{\Sigma_j}^{(i)} + L_{\Sigma_j}^{(i)} f) \right] \bullet \mathcal{C}^{(j)} \bullet \tilde{\mathcal{L}}^{(j+1)} \\
&\quad + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} \bullet \mathcal{C}^{(j)} \bullet \tilde{\mathcal{L}}^{(j+1)} + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} \bullet \mathcal{C}^{(j)} \bullet \left[ \sum_{i=2}^j (F_{\Sigma_j}^{(i)} + L_{\Sigma_j}^{(i)} f) \right] \\
&= \left[ \sum_{i=2}^j F_{\Sigma_j}^{(i)} + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} f + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} \right] \bullet \mathcal{C}^{(j)} \bullet \left[ \sum_{i=2}^j F_{\Sigma_j}^{(i)} + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} f + \tilde{\mathcal{L}}^{(j+1)} \right]
\end{aligned}$$

ii) Substituting  $\tilde{\mathcal{L}}^{(j+1)} = \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} + L^{(j+1)}$  into part (i) gives

$$\begin{aligned}
L^{(j+1)} &= \left( \sum_{i=2}^j F_{\Sigma_j}^{(i)} + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} f + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} \right) \bullet \mathcal{C}^{(j)} \bullet \left( \sum_{i=2}^j F_{\Sigma_j}^{(i)} + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} f + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} \right) \\
&\quad + \left( \sum_{i=2}^j F_{\Sigma_j}^{(i)} + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} f + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} \right) \bullet \mathcal{C}^{(j)} \bullet L^{(j+1)}
\end{aligned}$$

Now just iterate. ■

**Proof of Proposition II.3:** By Lemma II.5.ii,

$$\begin{aligned}\tilde{\mathcal{L}}^{(j+1)} &= \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} + L^{(j+1)} \\ &= \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} + \sum_{\ell=1}^{\infty} (F + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)f}) \bullet \mathcal{C}^{(j)} \bullet \dots \bullet \mathcal{C}^{(j)} \bullet (F + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)} + \tilde{\mathcal{L}}_{\Sigma_j}^{(j)f})\end{aligned}$$

where  $F = \sum_{i=2}^j F_{\Sigma_j}^{(i)}$  and the  $\ell^{\text{th}}$  term has  $\ell$  bubble propagators  $\mathcal{C}^{(j)}$ . Thus  $\tilde{\mathcal{L}}^{(j)}$  obeys the same initial condition and recursion relation as that defining  $\mathcal{L}^{(j)}$  in Definition I.19. Therefore, they are equal. Hence the Proposition follows from Lemma II.4 and Lemma II.5.i.  $\blacksquare$

## Spin Independence

The following discussion shows how spin independent functions on  $(\mathfrak{Y}_{\Sigma}^{\uparrow})^4$  are related to functions on  $\mathfrak{Y}_{\Sigma}^4$ .

**Definition II.6 (Spin Independence)** Let  $\mathfrak{Z}_l$  and  $\mathfrak{Z}_r$  be sets and let  $f$  be a function on  $(\mathfrak{Z}_l \times \{\uparrow, \downarrow\})^2 \times (\mathfrak{Z}_r \times \{\uparrow, \downarrow\})^2$ . Set, for each  $A \in SU(2)$ ,

$$f^A((\cdot, \sigma_1), \dots, (\cdot, \sigma_4)) = \sum_{\tau_1, \dots, \tau_4} f((\cdot, \tau_1), \dots, (\cdot, \tau_4)) A_{\tau_1, \sigma_1} \bar{A}_{\tau_2, \sigma_2} \bar{A}_{\tau_3, \sigma_3} A_{\tau_4, \sigma_4}$$

$f$  is called (particle-hole) spin independent if  $f = f^A$  for all  $A \in SU(2)$ .

**Remark II.7** Let  $F$  be a four-legged kernel on  $\mathfrak{X}_{\Sigma}$ . If  $F$  is spin independent in the sense of Definition B.1.S of [FKTo2], then its particle-hole reduction is spin independent in the sense of Definition II.6.

**Lemma II.8 (Charge Spin Representation)** Let  $\mathfrak{Z}_l$  and  $\mathfrak{Z}_r$  be sets and let  $f$  be a spin independent function on  $(\mathfrak{Z}_l \times \{\uparrow, \downarrow\})^2 \times (\mathfrak{Z}_r \times \{\uparrow, \downarrow\})^2$ . Then, there are functions  $f_C$  and  $f_S$  on  $\mathfrak{Z}_l^2 \times \mathfrak{Z}_r^2$  such that

$$\begin{aligned}f((z_1, \sigma_1), (z_2, \sigma_2), (z_3, \sigma_3), (z_4, \sigma_4)) &= \frac{1}{2} f_C(z_1, z_2, z_3, z_4) \delta_{\sigma_1, \sigma_2} \delta_{\sigma_3, \sigma_4} \\ &\quad + f_S(z_1, z_2, z_3, z_4) [\delta_{\sigma_1, \sigma_3} \delta_{\sigma_2, \sigma_4} - \frac{1}{2} \delta_{\sigma_1, \sigma_2} \delta_{\sigma_3, \sigma_4}]\end{aligned}$$

**Proof:** The statement is essentially [N, (1–7)]. The proof is outlined in [N] between (3–40) and (3–41). For the readers convenience, we include a detailed proof.

The  $z$ 's play no role, so we suppress them. Then the function  $f(\sigma_1, \sigma_2, \sigma_3, \sigma_4)$  can be viewed as an element of  $\mathbb{C}^{16} = \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$  and  $M_A : f \mapsto f^A$  is a linear map on  $\mathbb{C}^{16}$ . The map  $A \mapsto M_A$  is a representation of  $SU(2)$  on  $\mathbb{C}^{16}$ . Denote by  $S_n$  the

standard  $(2n + 1)$  dimensional “spin  $n$ ” irreducible representation of  $SU(2)$ . In particular, the identity representation  $A \mapsto A$  is  $S_{1/2}$ . Since the representation  $A \mapsto \bar{A}$  is unitarily equivalent to  $S_{1/2}$ , the representation  $A \mapsto M_A$  is unitarily equivalent to  $S_{1/2} \otimes S_{1/2} \otimes S_{1/2} \otimes S_{1/2} \cong (S_0 \oplus S_1) \otimes (S_0 \oplus S_1) \cong 2S_0 \oplus 3S_1 \oplus S_2$ . Thus the dimension of the subspace  $\{ f \in \mathbb{C}^{16} \mid f = f^A \ \forall A \in SU(2) \}$  is exactly two. Since  $f(\sigma_1, \sigma_2, \sigma_3, \sigma_4) = \delta_{\sigma_1, \sigma_2} \delta_{\sigma_3, \sigma_4}$  and  $f(\sigma_1, \sigma_2, \sigma_3, \sigma_4) = \delta_{\sigma_1, \sigma_3} \delta_{\sigma_2, \sigma_4} - \frac{1}{2} \delta_{\sigma_1, \sigma_2} \delta_{\sigma_3, \sigma_4}$  are two independent elements of that subspace, every  $f \in \mathbb{C}^{16}$  obeying  $f = f^A$  for all  $A \in SU(2)$  is a linear combination of  $\delta_{\sigma_1, \sigma_2} \delta_{\sigma_3, \sigma_4}$  and  $\delta_{\sigma_1, \sigma_3} \delta_{\sigma_2, \sigma_4} - \frac{1}{2} \delta_{\sigma_1, \sigma_2} \delta_{\sigma_3, \sigma_4}$ . ■

**Remark II.9**

$$\begin{aligned} f_C &= f((\cdot, \uparrow), (\cdot, \uparrow), (\cdot, \uparrow), (\cdot, \uparrow)) + f((\cdot, \uparrow), (\cdot, \uparrow), (\cdot, \downarrow), (\cdot, \downarrow)) \\ &= f((\cdot, \downarrow), (\cdot, \downarrow), (\cdot, \downarrow), (\cdot, \downarrow)) + f((\cdot, \downarrow), (\cdot, \downarrow), (\cdot, \uparrow), (\cdot, \uparrow)) \\ f_S &= f((\cdot, \uparrow), (\cdot, \downarrow), (\cdot, \uparrow), (\cdot, \downarrow)) = f((\cdot, \downarrow), (\cdot, \uparrow), (\cdot, \downarrow), (\cdot, \uparrow)) \end{aligned}$$

**Lemma II.10** *If  $K$  is a spin independent function on  $(\mathfrak{Z} \times \{\uparrow, \downarrow\})^4$ , then*

$$(K^f)_C = \frac{1}{2}(K_C + 3K_S)^f \quad (K^f)_S = \frac{1}{2}(K_C - K_S)^f$$

where  $K^f$  is the flipped function of (I.5).

**Proof:**

$$\begin{aligned} K^f((z_1, \sigma_1), (z_2, \sigma_2), (z_3, \sigma_3), (z_4, \sigma_4)) &= -K((z_1, \sigma_1), (z_3, \sigma_3), (z_2, \sigma_2), (z_4, \sigma_4)) \\ &= -\frac{1}{2}K_C(z_1, z_3, z_2, z_4)\delta_{\sigma_1, \sigma_3}\delta_{\sigma_2, \sigma_4} - K_S(z_1, z_3, z_2, z_4)[\delta_{\sigma_1, \sigma_2}\delta_{\sigma_3, \sigma_4} - \frac{1}{2}\delta_{\sigma_1, \sigma_3}\delta_{\sigma_2, \sigma_4}] \\ &= K_S^f(z_1, z_2, z_3, z_4)\delta_{\sigma_1, \sigma_2}\delta_{\sigma_3, \sigma_4} + \frac{1}{2}(K_C^f - K_S^f)(z_1, z_2, z_3, z_4)\delta_{\sigma_1, \sigma_3}\delta_{\sigma_2, \sigma_4} \\ &= \frac{1}{4}(K_C^f + 3K_S^f)(z_1, z_2, z_3, z_4)\delta_{\sigma_1, \sigma_2}\delta_{\sigma_3, \sigma_4} \\ &\quad + \frac{1}{2}(K_C^f - K_S^f)(z_1, z_2, z_3, z_4)[\delta_{\sigma_1, \sigma_3}\delta_{\sigma_2, \sigma_4} - \frac{1}{2}\delta_{\sigma_1, \sigma_2}\delta_{\sigma_3, \sigma_4}] \end{aligned}$$

**Lemma II.11** *If  $H'$  and  $K'$  are spin independent functions on  $(\mathfrak{Y}_{\Sigma}^{\uparrow})^4$  and  $P$  is a bubble propagator, then*

$$\begin{aligned} (H' \bullet P \bullet K')_C &= H'_C \bullet P \bullet K'_C \\ (H' \bullet P \bullet K')_S &= H'_S \bullet P \bullet K'_S \end{aligned}$$

**Proof:** This Lemma follows directly from Remark II.9. ■



Parts (ii) and (iii) of Proposition II.3, Lemma II.10 and Lemma II.11 give a coupled system of recursion relations for  $\mathcal{L}_C^{(j)}$ ,  $\mathcal{L}_S^{(j)}$ ,  $L_C^{(j)}$  and  $L_S^{(j)}$ .

**Corollary II.12**

$$\begin{aligned}
i) \quad \mathcal{L}_C^{(j+1)} &= \sum_{\ell=1}^{\infty} \sum_{i_1, \dots, i_{\ell+1}=2}^j \left[ \left( F_C^{(i_1)} + \frac{1}{2} L_C^{(i_1)} f + \frac{3}{2} L_S^{(i_1)} f \right) \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet \right. \\
&\quad \left. \dots \bullet \mathcal{C}^{[\max\{i_{\ell}, i_{\ell+1}\}, j]} \bullet \left( F_C^{(i_{\ell+1})} + \frac{1}{2} L_C^{(i_{\ell+1})} f + \frac{3}{2} L_S^{(i_{\ell+1})} f \right) \right]_{\Sigma_j} \\
\mathcal{L}_S^{(j+1)} &= \sum_{\ell=1}^{\infty} \sum_{i_1, \dots, i_{\ell+1}=2}^j \left[ \left( F_S^{(i_1)} + \frac{1}{2} L_C^{(i_1)} f - \frac{1}{2} L_S^{(i_1)} f \right) \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet \right. \\
&\quad \left. \dots \bullet \mathcal{C}^{[\max\{i_{\ell}, i_{\ell+1}\}, j]} \bullet \left( F_S^{(i_{\ell+1})} + \frac{1}{2} L_C^{(i_{\ell+1})} f - \frac{1}{2} L_S^{(i_{\ell+1})} f \right) \right]_{\Sigma_j} \\
ii) \quad L_C^{(j+1)} &= \left( \sum_{i=2}^j F_{C\Sigma_j}^{(i)} + \frac{1}{2} \mathcal{L}_C^{(j)} f + \frac{3}{2} \mathcal{L}_S^{(j)} f + \mathcal{L}_C^{(j)} \right) \bullet \mathcal{C}^{(j)} \\
&\quad \bullet \left( \sum_{i=2}^j F_{C\Sigma_j}^{(i)} + \frac{1}{2} \mathcal{L}_C^{(j)} f + \frac{3}{2} \mathcal{L}_S^{(j)} f + \mathcal{L}_C^{(j+1)} \right) \\
L_S^{(j+1)} &= \left( \sum_{i=2}^j F_{S\Sigma_j}^{(i)} + \frac{1}{2} \mathcal{L}_C^{(j)} f - \frac{1}{2} \mathcal{L}_S^{(j)} f + \mathcal{L}_S^{(j)} \right) \bullet \mathcal{C}^{(j)} \\
&\quad \bullet \left( \sum_{i=2}^j F_{S\Sigma_j}^{(i)} + \frac{1}{2} \mathcal{L}_C^{(j)} f - \frac{1}{2} \mathcal{L}_S^{(j)} f + \mathcal{L}_S^{(j+1)} \right)
\end{aligned}$$

Theorem I.20 will be proven by bounding each term on the right hand side of Corollary II.12.i. Each such term is a particle-hole ladder of the form

$$(G^{(i_1)} + K^{(i_1)} f) \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet \dots \bullet \mathcal{C}^{[\max\{i_{\ell}, i_{\ell+1}\}, j]} \bullet (G^{(i_{\ell+1})} + K^{(i_{\ell+1})} f)$$


where  $G^{(i)}$  is either  $F_C^{(i)}$  or  $F_S^{(i)}$  and  $K^{(i)}$  is a linear combination of  $L_C^{(i)}$  and  $L_S^{(i)}$ . This ladder has rungs  $(G^{(i_{\nu})} + K^{(i_{\nu})} f)$  which are connected by particle-hole propagators  $\mathcal{C}^{[i, j]}$ . The induction step will consist in adding an additional rung to the left of the ladder. More precisely, we will prove a bound on

$$(G^{(i_1)} + K^{(i_1)} f) \bullet \mathcal{C}^{[i, j]} \bullet H$$

with  $H = (G^{(i_2)} + K^{(i_2)} f) \bullet \mathcal{C}^{[\max\{i_2, i_3\}, j]} \bullet \dots \bullet (G^{(i_{\ell+1})} + K^{(i_{\ell+1})} f)$ , assuming bounds on  $H$ . The expression

$$G^{(i_1)} \bullet \mathcal{C}^{[i, j]} \bullet H$$


is a particle-hole bubble



We will derive the necessary bounds on general particle-hole bubbles in Theorem II.19. By Corollary II.12.ii,

$$K^{(i_1)f} \bullet \mathcal{C}^{[i,j]} \bullet H = \left( G_1^{(i_1)} \bullet \mathcal{C}^{(i_1-1)} \bullet G_2^{(i_1)} \right)^f \bullet \mathcal{C}^{[i,j]} \bullet H$$

with  $G_1^{(i)}$  and  $G_2^{(i)}$  linear combinations of  $\sum_{k=2}^{i-1} F_C^{(k)}$ ,  $\sum_{k=2}^{i-1} F_S^{(k)}$ ,  $\mathcal{L}_C^{(i-1)}$ ,  $\mathcal{L}_S^{(i-1)}$ ,  $\mathcal{L}_C^{(i-1)f}$ ,  $\mathcal{L}_S^{(i-1)f}$ ,  $\mathcal{L}_C^{(i)}$ ,  $\mathcal{L}_S^{(i)}$ . It is a double bubble



Bounds on double bubbles will be obtained in Theorem II.20.

### Scaled Norms

In the induction procedure outlined above the various ladders naturally have different sectorization scales at their left and right hand ends. This was the motivation for Definition I.14.

**Convention II.13** *Introduce, for scales  $\ell, r$ , the short hand notation*

$$\mathfrak{Y}_{\ell,r} = \mathfrak{Y}_{\Sigma_\ell, \Sigma_r}^{(4)}$$

**Definition II.14** For a function  $f$  on  $\mathfrak{Y}_{\ell,r}$  and multiindices  $\delta_1, \delta_c, \delta_r \in \mathbb{N}_0 \times \mathbb{N}_0^2$ , set

$$\begin{aligned} \|f\|_{\ell,r}^{(\delta_1, \delta_c, \delta_r)} &= \frac{1}{M^{\ell|\delta_1| + |\delta_c| \max(\ell, r) + r|\delta_r|}} |f|_{\Sigma_\ell, \Sigma_r}^{(\delta_1, \delta_c, \delta_r)} \\ |f|_{\ell,r}^{[\delta_1, \delta_c, \delta_r]} &= \max_{\substack{\delta'_1 \leq \delta_1 \\ \delta'_c \leq \delta_c \\ \delta'_r \leq \delta_r}} \|f\|_{\ell,r}^{(\delta'_1, \delta'_c, \delta'_r)} \end{aligned}$$

The norm  $\|\cdot\|_{\Sigma_\ell, \Sigma_r}^{(\delta_1, \delta_c, \delta_r)}$  was defined in Definition I.14. If  $\ell = r = j$ , set

$$|f|_j^{[\delta]} = \max_{\substack{\delta_1, \delta_c, \delta_r \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \delta_1 + \delta_c + \delta_r \leq \delta}} \|f\|_{j,j}^{(\delta_1, \delta_c, \delta_r)}$$

Set

$$\begin{aligned}\Delta &= \{ \delta \in \mathbb{N}_0 \times \mathbb{N}_0^2 \mid \delta_0 \leq r_0, \delta_1 + \delta_2 \leq r_e \} \\ \vec{\Delta} &= \{ \vec{\delta} = (\delta_l, \delta_c, \delta_r) \in (\mathbb{N}_0 \times \mathbb{N}_0^2)^3 \mid \delta_l + \delta_c + \delta_r \in \Delta \}\end{aligned}\tag{II.1}$$

where  $r_e + 3$  is the degree of differentiability of the dispersion relation  $e(\mathbf{k})$  and  $r_0$  is the number of  $k_0$  derivatives that we wish to control. The numbers  $r_e$  and  $r_0$  also determine the number of finite coefficients in the formal power series  $\mathbf{c}_j$  of (I.4). The following remark relates the formal power series norms of Definition I.15.ii to the norms of Definition II.14.

**Remark II.15** There is a constant  $const$ , depending only on  $r_e$  and  $r_0$  such that the following holds. Let  $f$  be a sectorized, translation invariant function on  $\mathfrak{Y}_{\Sigma_j}^4$ .

i)

$$|f|_{\Sigma_j} \leq \left[ \max_{\delta \in \Delta} |f|_j^{[\delta]} \right] \mathbf{c}_j$$

ii) If there is a number  $\gamma$  such that  $|f|_{\Sigma_j} \leq \gamma \mathbf{c}_j$ , then

$$|f|_j^{[\delta]} \leq const \gamma \quad \text{for all } \delta \in \Delta$$

Thus to prove Theorem I.20, it suffices to prove that

$$\max_{\delta \in \Delta} |\mathcal{L}_C^{(j+1)}|_j^{[\delta]} \leq const \rho^2 \quad \max_{\delta \in \Delta} |\mathcal{L}_S^{(j+1)}|_j^{[\delta]} \leq const \rho^2$$

**Definition II.16 (Norms and Resectorization)** Let  $\ell, \ell', r, r' \geq 0$ . For a sectorized, translation invariant, function  $f$  on  $\mathfrak{Y}_{\ell', r'}$  and multiindices  $\vec{\delta} \in (\mathbb{N}_0 \times \mathbb{N}_0^2)^3$ , set

$$|f|_{\ell, r}^{[\vec{\delta}]} = |f_{\Sigma_\ell, \Sigma_r}|_{\ell, r}^{[\vec{\delta}]}$$

If  $\ell = r = j$  and  $\delta \in \mathbb{N}_0 \times \mathbb{N}_0^2$ , set

$$|f|_j^{[\delta]} = \max_{\substack{\delta_l, \delta_c, \delta_r \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \delta_l + \delta_c + \delta_r \leq \delta}} |f|_{j, j}^{[\delta_l, \delta_c, \delta_r]}$$

As in Proposition XIX.4 of [FKTo4], one proves

**Lemma II.17** Let  $\ell \geq \ell' \geq 1$  and  $r \geq r' \geq 1$ . Let  $f$  be a sectorized, translation invariant, function on  $\mathfrak{Y}_{\ell', r'}$  and let  $\vec{\delta} = (\delta_l, \delta_c, \delta_r) \in \vec{\Delta}$ . Then

$$\begin{aligned}|f|_{\ell, r}^{[\vec{\delta}]} &\leq const \left\{ \frac{1}{M^{\ell-\ell'}} \frac{1}{M^{r-r'}} |f|_{\ell', r'}^{[\vec{\delta}]} + \frac{1}{M^{\ell-\ell'}} |f|_{\ell', r'}^{[\delta_l, \delta_c, 0]} + \frac{1}{M^{r-r'}} |f|_{\ell', r'}^{[0, \delta_c, \delta_r]} + |f|_{\ell', r'}^{[0, \delta_c, 0]} \right\} \\ &\leq const |f|_{\ell', r'}^{[\vec{\delta}]}\end{aligned}$$

The constant  $const$  depends only on  $\Delta$ .

**Proof:** Let  $f$  be a function on  $\mathfrak{Y}_{i_1, \Sigma_{\ell'}} \times \mathfrak{Y}_{i_2, \Sigma_{\ell'}} \times \mathfrak{Y}_{i_3, \Sigma_{r'}} \times \mathfrak{Y}_{i_4, \Sigma_{r'}}$ . We consider the case  $i_1 = i_2 = i_3 = i_4 = 1$  and  $\ell' < \ell$ ,  $r' < r$ . The other cases are similar, but easier. Recall from Definition I.17 that,

$$\begin{aligned} f_{\Sigma_{\ell}, \Sigma_r}((x_1, s_1), (x_2, s_2), (x_3, s_3), (x_4, s_4)) \\ = \sum_{\substack{s'_\nu \in \Sigma_{\ell'} \\ \nu \in \{1, 2\}}} \sum_{\substack{s'_\nu \in \Sigma_{r'} \\ \nu \in \{3, 4\}}} \int \prod_{\nu=1}^4 \left( dx'_\nu \hat{\chi}_{s'_\nu}((-1)^{b_\nu}(x_\nu - x'_\nu)) \right) f((x'_1, s'_1), (x'_2, s'_2), (x'_3, s'_3), (x'_4, s'_4)) \end{aligned}$$

First observe that, for any fixed  $s_1, \dots, s_4$ , there are at most  $3^4$  choices of  $(s'_1, \dots, s'_4)$  for which the integral  $\int \prod_{\nu=1}^4 \left( dx'_\nu \hat{\chi}_{s'_\nu}(\dots) \right) f(\dots)$  fails to vanish identically, because  $f$  is sectorized and  $\ell' < \ell$ ,  $r' < r$ . So it suffices to consider any fixed  $s'_1, \dots, s'_4$ . Hence by Leibniz's Rule (Lemma II.21),  $\|f_{\Sigma_{\ell}, \Sigma_r}\|_{\ell, r}^{(\delta_1, \delta_c, \delta_r)}$  is bounded by a constant, which depends only on  $\Delta$ , times the maximum of

$$\begin{aligned} & \frac{1}{M^{\ell|\delta_1|+|\delta_c|\max(\ell, r)+r|\delta_r|}} \int \prod_{\nu=2}^4 dx_\nu \int \prod_{\nu=1}^4 \left( dx'_\nu |(x_\nu - x'_\nu)^{\beta_\nu} \hat{\chi}_{s'_\nu}((-1)^{b_\nu}(x_\nu - x'_\nu))| \right) \\ & \quad \left| D_{1;2}^{\alpha_1} D_{\mu; \mu'}^{\alpha_c} D_{3;4}^{\alpha_r} f((x'_1, s'_1), (x'_2, s'_2), (x'_3, s'_3), (x'_4, s'_4)) \right| \\ & \leq \frac{1}{M^{\ell|\delta_1|+|\delta_c|\max(\ell, r)+r|\delta_r|}} \left( \prod_{\nu=1}^4 \|x_\nu^{\beta_\nu} \hat{\chi}_{s'_\nu}(x_\nu)\|_{L^1} \right) \|f\|_{\Sigma_{\ell'}, \Sigma_{r'}}^{(\alpha_1, \alpha_c, \alpha_r)} \\ & = \frac{M^{\ell'|\alpha_1|+|\alpha_c|\max(\ell', r')+r'|\alpha_r|}}{M^{\ell|\delta_1|+|\delta_c|\max(\ell, r)+r|\delta_r|}} \left( \prod_{\nu=1}^4 \|x_\nu^{\beta_\nu} \hat{\chi}_{s'_\nu}(x_\nu)\|_{L^1} \right) \|f\|_{\Sigma_{\ell'}, \Sigma_{r'}}^{(\alpha_1, \alpha_c, \alpha_r)} \end{aligned}$$

over  $x_1, s_1, \dots, s_4, s'_1, \dots, s'_4$  and  $\mu \in \{1, 2\}$ ,  $\mu' \in \{3, 4\}$  and  $\alpha_1, \alpha_c, \alpha_r$  and

$$\beta_\nu = \beta_{\nu,1} + \beta_{\nu,c} + \beta_{\nu,r} \quad \nu = 1, \dots, 4$$

obeying

$$\begin{aligned} \beta_{1,1} + \alpha_1 + \beta_{2,1} &= \delta_1 & \beta_{\mu,c} + \alpha_c + \beta_{\mu',c} &= \delta_c & \beta_{3,r} + \alpha_r + \beta_{4,r} &= \delta_r \\ \beta_{1,r} &= \beta_{2,r} = \beta_{3,1} = \beta_{4,1} = \beta_{\nu,c} = 0 & \text{for } \nu &\neq \mu, \mu' \end{aligned}$$

In particular

$$\ell|\delta_1| + |\delta_c|\max(\ell, r) + r|\delta_r| \geq \ell|\alpha_1 + \beta_1 + \beta_2| + |\alpha_c|\max(\ell, r) + r|\alpha_r + \beta_3 + \beta_4|$$

By Lemma XII.3 of [FKTo3]

$$\|x_\nu^{\beta_\nu} \hat{\chi}_{s'_\nu}(x_\nu)\|_{L^1} \leq \text{const} \begin{cases} M^{|\beta_\nu|\ell} & \text{if } \nu \in \{1, 2\} \\ M^{|\beta_\nu|r} & \text{if } \nu \in \{3, 4\} \end{cases} \quad (\text{II.2})$$

so that

$$\begin{aligned} & \frac{M^{\ell'|\alpha_1|+|\alpha_c|\max(\ell', r')+r'|\alpha_r|}}{M^{\ell|\delta_1|+|\delta_c|\max(\ell, r)+r|\delta_r|}} \prod_{\nu=1}^4 \|x_\nu^{\beta_\nu} \hat{\chi}_{s'_\nu}(x_\nu)\|_{L^1} \\ & \leq \text{const} \frac{M^{\ell'|\alpha_1|+|\alpha_c|\max(\ell', r')+r'|\alpha_r|}}{M^{\ell|\delta_1|+|\delta_c|\max(\ell, r)+r|\delta_r|}} M^{\ell|\beta_1+\beta_2|+r|\beta_3+\beta_4|} \\ & \leq \text{const} \frac{1}{M^{(\ell-\ell')|\alpha_1|+(r-r')|\alpha_r|}} \end{aligned}$$

and

$$\|f_{\Sigma_\ell, \Sigma_r}\|_{\ell, r}^{(\delta_1, \delta_c, \delta_r)} \leq \text{const} \max_{\substack{\alpha_1 \leq \delta_1 \\ \alpha_c \leq \delta_c \\ \alpha_r \leq \delta_r}} \frac{1}{M^{(\ell-\ell')|\alpha_1|+(r-r')|\alpha_r|}} \|f\|_{\Sigma_{\ell'}, \Sigma_{r'}}^{(\alpha_1, \alpha_c, \alpha_r)}$$

and the Lemma follows. ■

## Bubble and Double Bubble Bounds

**Definition II.18** Let  $i \leq j$ . Then

$$\mathcal{C}^{[i,j]} = \mathcal{C}_{\text{top}}^{[i,j]} + \mathcal{C}_{\text{mid}}^{[i,j]} + \mathcal{C}_{\text{bot}}^{[i,j]}$$

where

$$\begin{aligned} \mathcal{C}_{\text{top}}^{[i,j]} &= \sum_{\substack{i \leq i_t \leq j \\ i_b > j}} C_v^{(i_t)} \otimes C_v^{(i_b) t} \\ \mathcal{C}_{\text{mid}}^{[i,j]} &= \sum_{\substack{i \leq i_t \leq j \\ i \leq i_b \leq j}} C_v^{(i_t)} \otimes C_v^{(i_b) t} \\ \mathcal{C}_{\text{bot}}^{[i,j]} &= \sum_{\substack{i_t > j \\ i \leq i_b \leq j}} C_v^{(i_t)} \otimes C_v^{(i_b) t} \end{aligned}$$

**Theorem II.19 (Bubble Bound)** Let  $1 \leq i, \ell \leq j$  and  $\delta_1, \delta_r \in \Delta$ . Let  $g$  and  $h$  be sectorized, translation invariant functions on  $\mathfrak{Y}_{\ell, i}$  and  $\mathfrak{Y}_{i, j}$  respectively. Then

a)

$$|g \bullet \mathcal{C}^{[i,j]} \bullet h|_{\ell, j}^{[\delta_1, 0, \delta_r]} \leq \text{const} \max_{\substack{\alpha_r, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_r| + |\alpha_1| \leq 3}} |g|_{\ell, i}^{[\delta_1, 0, \alpha_r]} |h|_{i, j}^{[\alpha_1, 0, \delta_r]}$$

b) For any  $\beta \in \Delta$

$$\begin{aligned} \frac{1}{M^{|\beta|j}} \|g \bullet D_{1;3}^\beta \mathcal{C}_{\text{top}}^{[i,j]} \bullet h\|_{\ell, j}^{(\delta_1, 0, \delta_r)} &\leq \text{const} \|g\|_{\ell, i}^{(\delta_1, 0, 0)} \|h\|_{i, j}^{(0, 0, \delta_r)} \\ \frac{1}{M^{|\beta|j}} \|g \bullet D_{2;4}^\beta \mathcal{C}_{\text{bot}}^{[i,j]} \bullet h\|_{\ell, j}^{(\delta_1, 0, \delta_r)} &\leq \text{const} \|g\|_{\ell, i}^{(\delta_1, 0, 0)} \|h\|_{i, j}^{(0, 0, \delta_r)} \end{aligned}$$

c)

$$\|g \bullet \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h\|_{\ell, j}^{(\delta_1, 0, \delta_r)} \leq \text{const} |j - i + 1| \|g\|_{\ell, i}^{(\delta_1, 0, 0)} \|h\|_{i, j}^{(0, 0, \delta_r)}$$

and for any  $\beta \in \Delta$  with  $|\beta| \geq 1$  and  $(\mu, \mu') = (1, 3), (2, 4)$

$$\frac{1}{M^{|\beta|j}} \|g \bullet D_{\mu; \mu'}^\beta \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h\|_{\ell, j}^{(\delta_1, 0, \delta_r)} \leq \text{const} \|g\|_{\ell, i}^{(\delta_1, 0, 0)} \|h\|_{i, j}^{(0, 0, \delta_r)}$$

This Theorem is proven in §III.

**Theorem II.20 (Double Bubble Bound)** *Let  $1 \leq \ell \leq i \leq j$ ,  $\nu \in \mathbb{N}_0 \times \mathbb{N}_0^2$  and  $\delta_l, \delta_r \in \Delta$ . Let  $g_1, g_2$  and  $h$  be sectorized, translation invariant functions on  $\mathfrak{Y}_{\ell,\ell}, \mathfrak{Y}_{\ell,\ell}$  and  $\mathfrak{Y}_{i,j}$  respectively. Let  $\mathcal{D}$  be either*

$$\mathcal{D}_{\nu,\text{up}}^{(\ell)}(x_1, x_2, x_3, x_4) = \frac{1}{M^{|\nu|_\ell}} \sum_{m=\ell}^{\infty} D_{1;3}^\nu C_v^{(\ell)}(x_1, x_3) C_v^{(m)}(x_4, x_2)$$

or

$$\mathcal{D}_{\nu,\text{dn}}^{(\ell)}(x_1, x_2, x_3, x_4) = \frac{1}{M^{|\nu|_\ell}} \sum_{m=\ell+1}^{\infty} C_v^{(m)}(x_1, x_3) D_{2;4}^\nu C_v^{(\ell)}(x_4, x_2)$$

a) *If  $\nu + \delta_l + \alpha \in \Delta$  for all  $|\alpha| \leq 3$ , then*

$$|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{C}^{[i,j]} \bullet h|_{\ell,j}^{[\delta_l,0,\delta_r]} \leq \text{const } i \sqrt{\mathfrak{l}_\ell} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} |g_1|_{\ell}^{\llbracket \delta_l + \alpha_{\text{up}} \rrbracket} |g_2|_{\ell}^{\llbracket \delta_l + \alpha_{\text{dn}} \rrbracket} |h|_{i,j}^{[\alpha_1,0,\delta_r]}$$

b) *If  $\nu + \delta_l \in \Delta$ , then for any  $\beta \in \Delta$*

$$\begin{aligned} \frac{1}{M^{|\beta|_j}} \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet D_{1;3}^\beta \mathcal{C}_{\text{top}}^{[i,j]} \bullet h\|_{\ell,j}^{(\delta_l,0,\delta_r)} &\leq \text{const } \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell,\ell}^{[0,\delta_l,0]} |g_2|_{\ell,\ell}^{[0,\delta_l,0]} \|h\|_{i,j}^{(0,0,\delta_r)} \\ \frac{1}{M^{|\beta|_j}} \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet D_{2;4}^\beta \mathcal{C}_{\text{bot}}^{[i,j]} \bullet h\|_{\ell,j}^{(\delta_l,0,\delta_r)} &\leq \text{const } \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell,\ell}^{[0,\delta_l,0]} |g_2|_{\ell,\ell}^{[0,\delta_l,0]} \|h\|_{i,j}^{(0,0,\delta_r)} \end{aligned}$$

c) *If  $\nu + \delta_l \in \Delta$ , then*

$$\|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h\|_{\ell,j}^{(\delta_l,0,\delta_r)} \leq \text{const } |j - i + 1| \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell,\ell}^{[0,\delta_l,0]} |g_2|_{\ell,\ell}^{[0,\delta_l,0]} \|h\|_{i,j}^{(0,0,\delta_r)}$$

and for any  $\beta \in \Delta$  with  $|\beta| \geq 1$  and  $(\mu, \mu') = (1, 3), (2, 4)$

$$\frac{1}{M^{|\beta|_j}} \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet D_{\mu;\mu'}^\beta \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h\|_{\ell,j}^{(\delta_l,0,\delta_r)} \leq \text{const } \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell,\ell}^{[0,\delta_l,0]} |g_2|_{\ell,\ell}^{[0,\delta_l,0]} \|h\|_{i,j}^{(0,0,\delta_r)}$$

This Theorem is proven in §IV.

**Remark.** Observe that

$$\mathcal{C}^{(\ell)} = \mathcal{D}_{0,\text{up}}^{(\ell)} + \mathcal{D}_{0,\text{dn}}^{(\ell)}$$

We use Leibniz's rule to convert Theorems II.19 and II.20 into bounds on derivatives of  $g \bullet \mathcal{C}^{[i,j]} \bullet h$  and  $(g_1 \bullet \mathcal{C}^{(\ell)} \bullet g_2)^f \bullet \mathcal{C}^{[i,j]} \bullet h$  with respect to transfer momenta. These bounds are stated in Corollaries II.22, II.23 and II.24, below.

**Lemma II.21 (Leibniz's Rule)** Let  $\ell_1, r_1, \ell_2, r_2 \geq 1$ ,  $P$  a bubble propagator and  $K_1, K_2$  sectorized, translation invariant functions on  $\mathfrak{Y}_{\ell_1, r_1}$  and  $\mathfrak{Y}_{\ell_2, r_2}$ , respectively. Let  $\mu, \nu \in \{1, 2\}$ ,  $\mu', \nu' \in \{3, 4\}$  and  $\delta \in \mathbb{N}_0 \times \mathbb{N}_0^2$ . Then,

$$D_{\nu; \nu'}^\delta(K_1 \bullet P \bullet K_2) = \sum_{\substack{\beta_1, \beta_2, \beta_3 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \beta_1 + \beta_2 + \beta_3 = \delta}} \binom{\delta}{\beta_1, \beta_2, \beta_3} (D_{\nu; \mu+2}^{\beta_1} K_1) \bullet (D_{\mu; \mu'}^{\beta_2} P) \bullet (D_{\mu'-2; \nu'}^{\beta_3} K_2)$$

Here  $\binom{\delta}{\beta_1, \beta_2, \beta_3} = \frac{\delta!}{\beta_1! \beta_2! \beta_3!}$ .

**Proof:** The proof is trivial. ■

**Corollary II.22** Let  $1 \leq \ell \leq i \leq j$  and  $\delta_l, \delta_c, \delta_r \in \Delta$ . Let  $g$  and  $h$  be sectorized, translation invariant functions on  $\mathfrak{Y}_{\ell, \ell}$  and  $\mathfrak{Y}_{i, j}$  respectively.

a)

$$\begin{aligned} \|g \bullet \mathcal{C}_{\text{top}}^{[i, j]} \bullet h\|_{\ell, j}^{(\delta_l, \delta_c, \delta_r)} &\leq \text{const } |g|_{\ell, \ell}^{[\delta_l, \delta_c, 0]} |h|_{i, j}^{[0, \delta_c, \delta_r]} \\ \|g \bullet \mathcal{C}_{\text{bot}}^{[i, j]} \bullet h\|_{\ell, j}^{(\delta_l, \delta_c, \delta_r)} &\leq \text{const } |g|_{\ell, \ell}^{[\delta_l, \delta_c, 0]} |h|_{i, j}^{[0, \delta_c, \delta_r]} \end{aligned}$$

b) For  $\mu \in \{1, 2\}$  and  $\mu' \in \{3, 4\}$

$$D_{\mu; \mu'}^{\delta_c}(g \bullet \mathcal{C}_{\text{mid}}^{[i, j]} \bullet h) = \sum_{\substack{\beta_1, \beta_2, \beta_3 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \beta_1 + \beta_2 + \beta_3 = \delta_c}} \binom{\delta_c}{\beta_1, \beta_2, \beta_3} D_{\mu; 3}^{\beta_1} g \bullet D_{1; 3}^{\beta_2} \mathcal{C}_{\text{mid}}^{[i, j]} \bullet D_{1; \mu'}^{\beta_3} h$$

and, for all  $\beta_1 + \beta_2 + \beta_3 = \delta_c$ ,

$$\begin{aligned} \frac{1}{M^{|\delta_c|j}} \|D_{\mu; 3}^{\beta_1} g \bullet D_{1; 3}^{\beta_2} \mathcal{C}_{\text{mid}}^{[i, j]} \bullet D_{1; \mu'}^{\beta_3} h\|_{\ell, j}^{(\delta_l, 0, \delta_r)} \\ \leq \frac{\text{const}}{M^{|\beta_1|(j-\ell)}} \begin{cases} (j-i+1) \|g\|_{\ell, \ell}^{(\delta_l, \beta_1, 0)} \|h\|_{i, j}^{(0, \beta_3, \delta_r)} & \text{if } \beta_2 = 0 \\ i \max_{|\alpha_r + \alpha_l| \leq 3} |g|_{\ell, \ell}^{[\delta_l, \beta_1, \alpha_r]} |h|_{i, j}^{[\alpha_l, \beta_3, \delta_r]} & \text{if } \beta_2 = 0 \\ \|g\|_{\ell, \ell}^{(\delta_l, \beta_1, 0)} \|h\|_{i, j}^{(0, \beta_3, \delta_r)} & \text{if } \beta_2 \neq 0 \end{cases} \end{aligned}$$

**Proof:** a) We consider the case of top. By Leibniz,

$$D_{\mu; \mu'}^{\delta_c}(g \bullet \mathcal{C}_{\text{top}}^{[i, j]} \bullet h) = \sum_{\substack{\beta_1, \beta_2, \beta_3 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \beta_1 + \beta_2 + \beta_3 = \delta_c}} \binom{\delta_c}{\beta_1, \beta_2, \beta_3} D_{\mu; 3}^{\beta_1} g \bullet D_{1; 3}^{\beta_2} \mathcal{C}_{\text{top}}^{[i, j]} \bullet D_{1; \mu'}^{\beta_3} h$$

The desired inequality follows by the triangle inequality, Theorem II.19b and Lemma II.17, with  $D_{\mu; 3}^{\beta_1} g$  in place of  $g$  and  $D_{1; \mu'}^{\beta_3} h$  in place of  $h$ .

b) The first statement is again Leibniz's rule. By the first statement of Theorem II.19.c, with  $D_{\mu;3}^{\beta_1}g$  in place of  $g$  and  $D_{1;\mu'}^{\beta_3}h$  in place of  $h$ ,

$$\begin{aligned} \frac{1}{M^{|\delta_c|j}} \|D_{\mu;3}^{\beta_1}g \bullet \mathcal{C}_{\text{mid}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3}h\|_{\ell,j}^{(\delta_1,0,\delta_r)} &\leq \text{const} \frac{j-i+1}{M^{|\delta_c|j}} \|D_{\mu;3}^{\beta_1}g\|_{\ell,i}^{(\delta_1,0,0)} \|D_{1;\mu'}^{\beta_3}h\|_{i,j}^{(0,0,\delta_r)} \\ &\leq \text{const} \frac{j-i+1}{M^{|\delta_c|j}} \|D_{\mu;3}^{\beta_1}g\|_{\ell,\ell}^{(\delta_1,0,0)} \|D_{1;\mu'}^{\beta_3}h\|_{i,j}^{(0,0,\delta_r)} \\ &\leq \text{const} \frac{j-i+1}{M^{|\delta_c|j}} M^{|\beta_1|\ell} \|g\|_{\ell,\ell}^{(\delta_1,\beta_1,0)} M^{|\beta_3|j} \|h\|_{i,j}^{(0,\beta_3,\delta_r)} \\ &\leq \text{const} \frac{j-i+1}{M^{|\beta_1|(|j-\ell|)}} \|g\|_{\ell,\ell}^{(\delta_1,\beta_1,0)} \|h\|_{i,j}^{(0,\beta_3,\delta_r)} \end{aligned}$$

For the second inequality, we used the variant

$$\|D_{\mu;3}^{\beta_1}g\|_{\ell,i}^{(\delta_1,0,0)} = \frac{1}{M^{|\delta_1|\ell}} |D_{1;2}^{\delta_1} D_{\mu;3}^{\beta_1}g|_{\ell,i}^{[0,0,0]} \leq \text{const} \frac{1}{M^{|\delta_1|\ell}} |D_{1;2}^{\delta_1} D_{\mu;3}^{\beta_1}g|_{\ell,\ell}^{[0,0,0]} = \text{const} \|D_{\mu;3}^{\beta_1}g\|_{\ell,\ell}^{(\delta_1,0,0)}$$

of Lemma II.17. The proof of the second case is similar, but with

$$\begin{aligned} \|g \bullet \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h\|_{\ell,j}^{(\delta_1,0,\delta_r)} &\leq |g \bullet \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h|_{\ell,j}^{[\delta_1,0,\delta_r]} + |g \bullet \mathcal{C}_{\text{top}}^{[i,j]} \bullet h|_{\ell,j}^{[\delta_1,0,\delta_r]} + |g \bullet \mathcal{C}_{\text{bot}}^{[i,j]} \bullet h|_{\ell,j}^{[\delta_1,0,\delta_r]} \\ &\leq \text{const} \max_{\substack{\alpha_r, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_r| + |\alpha_1| \leq 3}} |g|_{\ell,i}^{[\delta_1,0,\alpha_r]} |h|_{i,j}^{[\alpha_1,0,\delta_r]} \end{aligned}$$

(by Theorem II.19.a,b) used in place of the first statement of Theorem II.19.c. The proof of the third case is again similar, but with the second statement of Theorem II.19.c used in place of the first statement of Theorem II.19.c.  $\blacksquare$

**Corollary II.23** *Let  $1 \leq \ell \leq i \leq j$  and  $\delta_1, \delta_c, \delta_r \in \Delta$ . Let  $g_1, g_2$  and  $h$  be sectorized, translation invariant functions on  $\mathfrak{Y}_{\ell,\ell}, \mathfrak{Y}_{\ell,\ell}$  and  $\mathfrak{Y}_{i,j}$  respectively. Let  $\mu \in \{1, 2\}$ ,  $\mu' \in \{3, 4\}$  and*

$$g = (g_1 \bullet \mathcal{C}^{(\ell)} \bullet g_2)^f$$

a) *If  $\delta_1 + \delta_c \in \Delta$ , then*

$$\begin{aligned} \|g \bullet \mathcal{C}_{\text{top}}^{[i,j]} \bullet h\|_{\ell,j}^{(\delta_1,\delta_c,\delta_r)} &\leq \text{const} \sqrt{\ell} |g_1|_{\ell}^{[\delta_1+\delta_c]} |g_2|_{\ell}^{[\delta_1+\delta_c]} |h|_{i,j}^{[0,\delta_c,\delta_r]} \\ \|g \bullet \mathcal{C}_{\text{bot}}^{[i,j]} \bullet h\|_{\ell,j}^{(\delta_1,\delta_c,\delta_r)} &\leq \text{const} \sqrt{\ell} |g_1|_{\ell,\ell}^{[\delta_1+\delta_c]} |g_2|_{\ell}^{[\delta_1+\delta_c]} |h|_{i,j}^{[0,\delta_c,\delta_r]} \end{aligned}$$

b) *Let  $\beta_1 + \beta_2 + \beta_3 = \delta_c$ . If  $\delta_1 + \beta_1 \in \Delta$ , then*

$$\begin{aligned} \frac{1}{M^{|\delta_c|j}} \|D_{\mu;3}^{\beta_1}g \bullet D_{1;3}^{\beta_2}\mathcal{C}_{\text{mid}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3}h\|_{\ell,j}^{(\delta_1,0,\delta_r)} \\ \leq \frac{\text{const}}{M^{|\beta_1|(|j-\ell|)}} \sqrt{\ell} \begin{cases} (j-i+1) |g_1|_{\ell}^{[\delta_1+\beta_1]} |g_2|_{\ell}^{[\delta_1+\beta_1]} \|h\|_{i,j}^{(0,\beta_3,\delta_r)} & \text{if } \beta_2 = 0 \\ |g_1|_{\ell}^{[\delta_1+\beta_1]} |g_2|_{\ell}^{[\delta_1+\beta_1]} \|h\|_{i,j}^{(0,\beta_3,\delta_r)} & \text{if } \beta_2 \neq 0 \end{cases} \end{aligned}$$



If  $\beta_2 = 0$  and  $\delta_1 + \beta_1 + \alpha \in \Delta$  for all  $|\alpha| \leq 3$ , then

$$\begin{aligned} & \frac{1}{M^{|\delta_c|j}} \left\| D_{\mu;3}^{\beta_1} g \bullet D_{1;3}^{\beta_2} \mathcal{C}_{\text{mid}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} h \right\|_{\ell,j}^{(\delta_1,0,\delta_r)} \\ & \leq \frac{\text{const}}{M^{|\beta_1|(j-\ell)}} i \sqrt{\ell} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} |g_1|_{\ell}^{\llbracket \delta_1 + \beta_1 + \alpha_{\text{up}} \rrbracket} |g_2|_{\ell}^{\llbracket \delta_1 + \beta_1 + \alpha_{\text{dn}} \rrbracket} |h|_{i,j}^{[\alpha_1, \beta_3, \delta_r]} \end{aligned}$$

**Proof:** a) We consider the case of top. By Leibniz,

$$\begin{aligned} D_{1;\mu'}^{\delta_c} (g \bullet \mathcal{C}_{\text{top}}^{[i,j]} \bullet h) &= \sum_{\substack{\beta_1, \beta_2, \beta_3 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \beta_1 + \beta_2 + \beta_3 = \delta_c}} \binom{\delta_c}{\beta_1, \beta_2, \beta_3} D_{1;3}^{\beta_1} (g_1 \bullet \mathcal{C}^{(\ell)} \bullet g_2)^f \bullet D_{1;3}^{\beta_2} \mathcal{C}_{\text{top}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} h \\ D_{2;\mu'}^{\delta_c} (g \bullet \mathcal{C}_{\text{top}}^{[i,j]} \bullet h) &= \sum_{\substack{\beta_1, \beta_2, \beta_3 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \beta_1 + \beta_2 + \beta_3 = \delta_c}} \binom{\delta_c}{\beta_1, \beta_2, \beta_3} D_{2;3}^{\beta_1} (g_1 \bullet \mathcal{C}^{(\ell)} \bullet g_2)^f \bullet D_{1;3}^{\beta_2} \mathcal{C}_{\text{top}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} h \end{aligned}$$

Substitute  $\mathcal{C}^{(\ell)} = \mathcal{D}_{0,\text{up}}^{(\ell)} + \mathcal{D}_{0,\text{dn}}^{(\ell)}$ . We consider the case of up. Then

$$\begin{aligned} D_{1;3}^{\beta_1} (g_1 \bullet \mathcal{D}_{0,\text{up}}^{(\ell)} \bullet g_2)^f \bullet D_{1;3}^{\beta_2} \mathcal{C}_{\text{top}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} h &= (D_{1;2}^{\beta_1} g_1 \bullet \mathcal{D}_{0,\text{up}}^{(\ell)} \bullet g_2)^f \bullet D_{1;3}^{\beta_2} \mathcal{C}_{\text{top}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} h \\ D_{2;3}^{\beta_1} (g_1 \bullet \mathcal{D}_{0,\text{up}}^{(\ell)} \bullet g_2)^f \bullet D_{1;3}^{\beta_2} \mathcal{C}_{\text{top}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} h &= (D_{3;2}^{\beta_1} (g_1 \bullet \mathcal{D}_{0,\text{up}}^{(\ell)} \bullet g_2))^f \bullet D_{1;3}^{\beta_2} \mathcal{C}_{\text{top}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} h \\ &= (-1)^{|\beta_1|} \sum_{\substack{\gamma_1, \gamma_2, \gamma_3 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \gamma_1 + \gamma_2 + \gamma_3 = \beta_1}} M^{|\gamma_2|\ell} \binom{\beta_1}{\gamma_1, \gamma_2, \gamma_3} (D_{2;3}^{\gamma_1} g_1 \bullet \mathcal{D}_{\gamma_2, \text{up}}^{(\ell)} \bullet D_{1;3}^{\gamma_3} g_2)^f \bullet D_{1;3}^{\beta_2} \mathcal{C}_{\text{top}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} h \end{aligned}$$

The  $\frac{1}{M^{|\delta_c|j}} \left\| \cdot \right\|_{\ell,j}^{(\delta_1,0,\delta_r)}$  norm of each term is bounded by Theorem II.20.b.

b) As above, we must estimate the  $\frac{1}{M^{|\delta_c|j}} \left\| \cdot \right\|_{\ell,j}^{(\delta_1,0,\delta_r)}$  norm of terms like

$$(D_{1;2}^{\beta_1} g_1 \bullet \mathcal{D}_{0,\text{up}}^{(\ell)} \bullet g_2)^f \bullet D_{1;3}^{\beta_2} \mathcal{C}_{\text{mid}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} h$$

and

$$M^{|\gamma_2|\ell} (D_{2;3}^{\gamma_1} g_1 \bullet \mathcal{D}_{\gamma_2, \text{up}}^{(\ell)} \bullet D_{1;3}^{\gamma_3} g_2)^f \bullet D_{1;3}^{\beta_2} \mathcal{C}_{\text{mid}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} h$$

with  $\gamma_1 + \gamma_2 + \gamma_3 = \beta_1$ . This is done using Theorem II.20. (In the last case, we write  $\mathcal{C}_{\text{mid}}^{[i,j]} = \mathcal{C}^{[i,j]} - \mathcal{C}_{\text{top}}^{[i,j]} - \mathcal{C}_{\text{bot}}^{[i,j]}$ .) ■

**Corollary II.24** Let  $1 \leq \ell \leq i \leq j$ ,  $1 \leq r \leq j$  and  $\delta_1, \delta_c, \delta_r \in \Delta$ . Let  $\mu \in \{1, 2\}$  and  $\mu' \in \{3, 4\}$ . Let  $h$  be a sectorized, translation invariant function on  $\mathfrak{Y}_{i,r}$  and let  $h' = h_{\Sigma_i, \Sigma_j}$  be its resectorization as in Definition I.17.i.

a) Let  $g$  be a sectorized, translation invariant function on  $\mathfrak{Y}_{\ell,i}$ . Then

$$\frac{1}{M^{j|\delta_c|}} \|g \bullet \mathcal{C}_{\text{mid}}^{[i,j]} \bullet D_{\mu;\mu'}^{\delta_c} h'\|_{\ell,j}^{(\delta_1,0,\delta_r)} \leq \text{const} \max_{\substack{\alpha_r, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_r| + |\alpha_1| \leq 3}} |g|_{\ell,i}^{[\delta_1,0,\alpha_r]} \left( \frac{j-i+1}{M^{j-i}} |h|_{i,r}^{[0,\delta_c,\delta_r]} + i |h|_{i,r}^{[\alpha_1,0,0]} \right)$$

b) Let  $g_1$  and  $g_2$  be sectorized, translation invariant functions on  $\mathfrak{Y}_{\ell,\ell}$ . If  $\delta_1 + \alpha \in \Delta$  for all  $|\alpha| \leq 3$ , then

$$\begin{aligned} \frac{1}{M^{j|\delta_c|}} \|(g_1 \bullet \mathcal{C}^{(\ell)} \bullet g_2)^f \bullet \mathcal{C}_{\text{mid}}^{[i,j]} \bullet D_{\mu;\mu'}^{\delta_c} h'\|_{\ell,j}^{(\delta_1,0,\delta_r)} \\ \leq \text{const} \sqrt{\ell} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} |g_1|_{\ell}^{[\delta_1+\alpha_{\text{up}}]} |g_2|_{\ell}^{[\delta_1+\alpha_{\text{dn}}]} \left( \frac{j-i+1}{M^{j-i}} |h|_{i,r}^{[0,\delta_c,\delta_r]} + i |h|_{i,r}^{[\alpha_1,0,0]} \right) \end{aligned}$$

**Proof:** We prove part a. First suppose that  $h$  is a function on  $\mathfrak{Y}_{\Sigma_i}^2 \times ((\mathbb{R} \times \mathbb{R}^2) \times \Sigma_r)^2$ . Then, for  $s_3, s_4 \in \Sigma_j$ ,

$$h'(\cdot, \cdot, (\cdot, s_3), (\cdot, s_4)) = h \bullet (\hat{\chi}_{s_3} \otimes \hat{\chi}_{s_4}^t)$$

We have

$$\frac{1}{M^{j|\delta_c|}} \|g \bullet \mathcal{C}_{\text{mid}}^{[i,j]} \bullet D_{\mu;\mu'}^{\delta_c} h'\|_{\ell,j}^{(\delta_1,0,\delta_r)} = \frac{1}{M^{j(|\delta_c|+|\delta_r|)}} \|g \bullet \mathcal{C}_{\text{mid}}^{[i,j]} \bullet D_{\mu;\mu'}^{\delta_c} D_{3;4}^{\delta_r} h'\|_{\ell,j}^{(\delta_1,0,0)}$$

Apply Leibniz to  $D_{\mu;\mu'}^{\delta_c} D_{3;4}^{\delta_r} (h \bullet (\hat{\chi}_{s_3} \otimes \hat{\chi}_{s_4}^t))$ , yielding a sum of terms of the form

$$D_{\mu;\mu'}^{\beta_1} D_{3;4}^{\gamma_1} h \bullet D_{1;3}^{\beta_2+\gamma_2} D_{2;4}^{\beta_3+\gamma_3} (\hat{\chi}_{s_3} \otimes \hat{\chi}_{s_4}^t)$$

with  $\beta_1 + \beta_2 + \beta_3 = \delta_c$  and  $\gamma_1 + \gamma_2 + \gamma_3 = \delta_r$ . If  $\beta_1 + \gamma_1 = 0$  we apply Theorem II.19a and otherwise we apply Theorem II.19c. The Lemma follows from

$$\begin{aligned} |h \bullet D_{1;3}^{\beta_2+\gamma_2} D_{2;4}^{\beta_3+\gamma_3} (\hat{\chi}_{s_3} \otimes \hat{\chi}_{s_4}^t)|_{i,j}^{[\alpha_1,0,0]} &\leq M^{j(|\delta_c|+|\delta_r|)} |h|_{i,r}^{[\alpha_1,0,0]} \\ \|D_{\mu;\mu'}^{\beta_1} D_{3;4}^{\gamma_1} h \bullet D_{1;3}^{\beta_2+\gamma_2} D_{2;4}^{\beta_3+\gamma_3} (\hat{\chi}_{s_3} \otimes \hat{\chi}_{s_4}^t)\|_{i,j}^{(0,0,0)} &\leq M^{j(|\beta_2+\gamma_2+\beta_3+\gamma_3|)} M^{|\beta_1|i+|\gamma_1|r} |h|_{i,r}^{[0,\delta_c,\delta_r]} \\ &\leq M^{j(|\delta_c|+|\delta_r|-1)} M^i |h|_{i,r}^{[0,\delta_c,\delta_r]} \quad \text{if } |\beta_1 + \gamma_1| \geq 1 \end{aligned}$$

which is proven in the same way as Lemma II.17 and, in particular, uses (II.2) with  $r = j$ . If one of the third or fourth arguments of  $h$  lie in momentum space,  $\mathbb{M}$ , the argument is similar, except that the corresponding  $\hat{\chi}_{s_3}$  or  $\hat{\chi}_{s_4}$  is omitted. The proof of part b is similar with Theorem II.20 used in place of Theorem II.19.  $\blacksquare$

**Proof of Theorem I.20** (assuming Theorems II.19 and II.20):

Let  $\delta \in \Delta$ . By the hypothesis of the Theorem and Remark II.15.ii, there is a constant  $c_F$  such that

$$\max_{\delta \in \Delta} |F_C^{(i)}|_i^{[\delta]} \leq \frac{c_F}{M^{\varepsilon i}} \rho \quad \max_{\delta \in \Delta} |F_S^{(i)}|_i^{[\delta]} \leq \frac{c_F}{M^{\varepsilon i}} \rho \quad (\text{II.3})$$

We prove by induction on  $j$  that

$$\max_{\delta \in \Delta} |\mathcal{L}_C^{(i)}|_{i-1}^{[\delta]} \leq c_{\mathcal{L}} \rho^2 \quad \max_{\delta \in \Delta} |\mathcal{L}_S^{(i)}|_{i-1}^{[\delta]} \leq c_{\mathcal{L}} \rho^2 \quad \text{for all } i \leq j \quad (\text{II.4})$$

with a constant  $c_{\mathcal{L}}$ , independent of  $j$ . See Remark II.15. By construction  $\mathcal{L}^{(0)} = \mathcal{L}^{(1)} = \mathcal{L}^{(2)} = 0$ . Now assume that (II.4) holds for some  $j \geq 2$ . We prove that

$$\max_{\delta \in \Delta} |\mathcal{L}_S^{(j+1)}|_j^{[\delta]} \leq c_{\mathcal{L}} \rho^2 \quad (\text{II.5})$$

The bound on  $\mathcal{L}_C^{(j+1)}$  is similar.

For  $i \leq j$  we have, by Corollary II.12.ii,

$$\begin{aligned} L_C^{(i)} &= G_{C,1}^{(i-1)} \bullet \mathcal{C}^{(i-1)} \bullet G_{C,2}^{(i-1)} \\ L_S^{(i)} &= G_{S,1}^{(i-1)} \bullet \mathcal{C}^{(i-1)} \bullet G_{S,2}^{(i-1)} \end{aligned}$$

with

$$\begin{aligned} G_{C,1}^{(i-1)} &= \sum_{i'=2}^{i-1} F_{C\Sigma_{i-1}}^{(i')} + \frac{1}{2} \mathcal{L}_{C\Sigma_{i-1}}^{(i-1)f} + \frac{3}{2} \mathcal{L}_{S\Sigma_{i-1}}^{(i-1)f} + \mathcal{L}_{C\Sigma_{i-1}}^{(i-1)} \\ G_{C,2}^{(i-1)} &= \sum_{i'=2}^{i-1} F_{C\Sigma_{i-1}}^{(i')} + \frac{1}{2} \mathcal{L}_{C\Sigma_{i-1}}^{(i-1)f} + \frac{3}{2} \mathcal{L}_{S\Sigma_{i-1}}^{(i-1)f} + \mathcal{L}_C^{(i)} \\ G_{S,1}^{(i-1)} &= \sum_{i'=2}^{i-1} F_{S\Sigma_{i-1}}^{(i')} + \frac{1}{2} \mathcal{L}_{C\Sigma_{i-1}}^{(i-1)f} - \frac{1}{2} \mathcal{L}_{S\Sigma_{i-1}}^{(i-1)f} + \mathcal{L}_{S\Sigma_{i-1}}^{(i-1)} \\ G_{S,2}^{(i-1)} &= \sum_{i'=2}^{i-1} F_{S\Sigma_{i-1}}^{(i')} + \frac{1}{2} \mathcal{L}_{C\Sigma_{i-1}}^{(i-1)f} - \frac{1}{2} \mathcal{L}_{S\Sigma_{i-1}}^{(i-1)f} + \mathcal{L}_S^{(i)} \end{aligned}$$

The hypotheses (II.3) on  $\vec{F}$  and the induction hypotheses (II.4) imply, via Lemma II.17, that, when  $\rho$  is small enough and  $M^\varepsilon$  is large enough,

$$\max_{\delta \in \Delta} |G_{C,\nu}^{(i-1)}|_{i-1}^{[\delta]} \leq c_F \rho \quad \max_{\delta \in \Delta} |G_{S,\nu}^{(i-1)}|_{i-1}^{[\delta]} \leq c_F \rho \quad (\text{II.6})$$

for  $i \leq j$ ,  $\nu = 1, 2$ .

**Remark II.25** For  $i \leq j$

$$\max_{\delta \in \Delta} |L_C^{(i)}|_{i-1}^{[\delta]} \leq \text{const } c_F^2 \rho^2 \quad \max_{\delta \in \Delta} |L_S^{(i)}|_{i-1}^{[\delta]} \leq \text{const } c_F^2 \rho^2$$

where const is  $(2 + 3^{r_0+2r_e})$  times the constant of Corollary II.22.

**Proof:** We prove the Remark for  $L_C^{(i)}$ . Fix  $(\delta_l, \delta_c, \delta_r) \in \vec{\Delta}$ . Decomposing

$$\mathcal{C}^{[i-1, i-1]} = (\mathcal{C}_{\text{top}}^{[i-1, i-1]} + \mathcal{C}_{\text{bot}}^{[i-1, i-1]}) + \mathcal{C}_{\text{mid}}^{[i-1, i-1]}$$

and applying Corollary II.22, parts a and b respectively, with  $\ell, i, j$  all replaced by  $i-1$ , we have

$$\begin{aligned} |G_{C,1}^{(i-1)} \bullet \mathcal{C}^{[i-1, i-1]} \bullet G_{C,2}^{(i-1)}|_{i-1, i-1}^{[\delta_l, \delta_c, \delta_r]} &\leq \text{const} \left( (1+1) + 3^{|\delta_c|} \right) |G_{C,1}^{(i-1)}|_{i-1, i-1}^{[\delta_l, \delta_c, 0]} |G_{C,2}^{(i-1)}|_{i-1, i-1}^{[0, \delta_c, \delta_r]} \\ &\leq \text{const} (2 + 3^{r_0+2r_e}) c_F^2 \rho^2 \end{aligned}$$

■

Set, for each  $i \geq 1$ ,

$$\mathbf{v}_i = i^2 \max \left\{ \sqrt{\mathfrak{l}_i}, \frac{1}{M^{\varepsilon i}} \right\}$$

and

$$K^{(i)} = F_S^{(i)} + \frac{1}{2} L_{C \Sigma_i}^{(i)f} - \frac{1}{2} L_{S \Sigma_i}^{(i)f}$$

Then, by Corollary II.12.i,

$$\mathcal{L}_S^{(j+1)} = \sum_{\ell=1}^{\infty} \sum_{i_1, \dots, i_{\ell+1}=2}^j \left[ K^{(i_1)} \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet K^{(i_2)} \bullet \dots \bullet \mathcal{C}^{[\max\{i_{\ell}, i_{\ell+1}\}, j]} \bullet K^{(i_{\ell+1})} \right]_{\Sigma_j}$$

We put the main estimates required to complete the proof of Theorem I.20 in

**Lemma II.26** *Let  $\ell \geq 1$  and  $i_1, \dots, i_{\ell+1} \leq j$ .*

a) *For  $|\alpha| \leq 3$  and  $\delta \in \Delta$ ,*

$$\left| K^{(i_1)} \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet K^{(i_2)} \bullet \dots \bullet K^{(i_{\ell+1})} \right|_{i_1, j}^{[\alpha, 0, \delta]} \leq \text{const}^{\ell} (c_F \rho)^{\ell+1} \frac{i_{\ell+1}}{i_1} \mathbf{v}_{i_1} \dots \mathbf{v}_{i_{\ell}}$$

b) *For  $(\delta_l, 0, \delta_r) \in \vec{\Delta}$*

$$\left| K^{(i_1)} \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet K^{(i_2)} \bullet \dots \bullet K^{(i_{\ell+1})} \right|_{j, j}^{[\delta_l, 0, \delta_r]} \leq \text{const}^{\ell} (c_F \rho)^{\ell+1} \mathbf{v}_{i_2} \dots \mathbf{v}_{i_{\ell}} \min \{ \mathbf{v}_{i_1}, \mathbf{v}_{i_{\ell+1}} \}$$

c) *For  $0 \neq \delta \in \Delta$ ,  $\mu \in \{1, 2\}$  and  $\mu' \in \{3, 4\}$ , there are sectorized, translation invariant functions  $k', k''$  on  $\mathfrak{Y}_{i_1, j}$  such that*

$$\frac{1}{M^{j|\delta|}} D_{\mu; \mu'}^{\delta} \left[ (K^{(i_1)} \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet \dots \bullet K^{(i_{\ell+1})})_{\Sigma_{i_1}, \Sigma_j} \right] = k' + k''$$

and, for all  $|\alpha| \leq 3$  and all  $\gamma$  with  $\gamma + \delta \in \Delta$ ,

$$\begin{aligned} |k'|_{i_1,j}^{[\alpha,0,\gamma]} &\leq \text{const}^\ell (c_F \rho)^{\ell+1} \frac{i_{\ell+1}}{i_1} \mathbf{v}_{i_1} \cdots \mathbf{v}_{i_\ell} \\ |k''|_{i_1,j}^{[0,0,\gamma]} &\leq \frac{j-i_1+1}{M^{j-i_1}} \text{const}^\ell (c_F \rho)^{\ell+1} \frac{i_{\ell+1}}{i_1} \mathbf{v}_{i_1} \cdots \mathbf{v}_{i_\ell} \end{aligned}$$

d) For  $(\delta_l, \delta_c, \delta_r) \in \vec{\Delta}$  with  $|\delta_c| \geq 1$ ,

$$\left| K^{(i_1)} \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet \dots \bullet K^{(i_{\ell+1})} \right|_{j,j}^{[\delta_l, \delta_c, \delta_r]} \leq \text{const}^\ell (c_F \rho)^{\ell+1} \mathbf{v}_{i_2} \cdots \mathbf{v}_{i_\ell} \min \{ \mathbf{v}_{i_1}, \mathbf{v}_{i_{\ell+1}} \}$$

Set  $i = \max\{i_1, i_2\}$  and write, for  $(\alpha_l, \alpha_c, \alpha_r) \in \vec{\Delta}$  and  $\alpha_{\text{up}}, \alpha_{\text{dn}} \in \Delta$ ,

$$\begin{aligned} q(\alpha_l, \alpha_c, \alpha_r; \alpha_{\text{up}}, \alpha_{\text{dn}}) &= i \left| F_S^{(i_1)} \right|_{i_1, i_1}^{[\alpha_l, \alpha_c, \alpha_r]} \\ &\quad + i \sqrt{l_{i_1-1}} \left[ \left| G_{C,1}^{(i_1-1)} \right|_{i_1-1}^{[\alpha_{\text{up}}]} \left| G_{C,2}^{(i_1-1)} \right|_{i_1-1}^{[\alpha_{\text{dn}}]} + \left| G_{S,1}^{(i_1-1)} \right|_{i_1-1}^{[\alpha_{\text{up}}]} \left| G_{S,2}^{(i_1-1)} \right|_{i_1-1}^{[\alpha_{\text{dn}}]} \right] \end{aligned}$$

By (II.3) and (II.6)

$$q(\alpha_l, \alpha_c, \alpha_r; \alpha_{\text{up}}, \alpha_{\text{dn}}) \leq i \frac{c_F \rho}{M^{\varepsilon i_1}} + i 2 c_F^2 \rho^2 \sqrt{l_{i_1-1}} \leq 2 c_F \rho \mathbf{v}_{i_1} \frac{i_2}{i_1} \quad (\text{II.7})$$

for  $\rho$  sufficiently small. The proof of Lemma II.26 follows

**Lemma II.27** *Let  $1 \leq i_1, i_2 \leq i \leq j$ ,  $1 \leq r \leq j$  and  $\delta_l, \delta_c, \delta_r \in \Delta$ . Let  $\mu \in \{1, 2\}$  and  $\mu' \in \{3, 4\}$  and  $\text{loc} \in \{\text{top}, \text{bot}, \text{mid}\}$ . If  $\text{loc} \in \{\text{top}, \text{mid}\}$ , set  $(\nu, \nu') = (1, 3)$ . If  $\text{loc} = \text{bot}$ , set  $(\nu, \nu') = (2, 4)$ . Let  $H$  be a sectorized, translation invariant function on  $\mathfrak{Y}_{i_2, r}$ .*

a) Let  $\beta_2 + \beta_3 = \delta$  with  $\beta_2 \neq 0$  and either  $|\delta_l| \leq 3$ ,  $\delta + \delta_r \in \Delta$  or  $\delta_l + \delta_r + \delta \in \Delta$ . Then

$$\frac{1}{M^{j|\delta|}} \left| K^{(i_1)} \bullet D_{\nu; \nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i, j]} \bullet D_{1; \mu'}^{\beta_3} H_{\Sigma_{i_2}, \Sigma_j} \right|_{i_1, j}^{[\delta_l, 0, \delta_r]} \leq \text{const } q(\delta_l, 0, 0; \delta_l, \delta_l) \left| H \right|_{i_2, j}^{[0, \beta_3, \delta_r]}$$

b) Let  $|\delta_l| \leq 3$  and  $\delta + \delta_r \in \Delta$ . Then

$$\begin{aligned} &\frac{1}{M^{j|\delta|}} \left| K^{(i_1)} \bullet \mathcal{C}_{\text{loc}}^{[i, j]} \bullet D_{1; \mu'}^\delta H_{\Sigma_{i_2}, \Sigma_j} \right|_{i_1, j}^{[\delta_l, 0, \delta_r]} \\ &\leq \text{const} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} q(\delta_l, 0, \alpha_{\text{up}} + \alpha_{\text{dn}}; \delta_l + \alpha_{\text{up}}, \delta_l + \alpha_{\text{dn}}) \left( \left| H \right|_{i_2, j}^{[0, \delta, \delta_r]} + \left| H \right|_{i_2, j}^{[\alpha_1, 0, 0]} \right) \end{aligned}$$

c) Let  $\beta_1 + \beta_2 + \beta_3 = \delta$  and  $\beta_1 + \delta_l, \delta, \delta_r \in \Delta$ . Then

$$\begin{aligned} &\frac{1}{M^{j|\delta|}} \left| D_{\mu; 3}^{\beta_1} K^{(i_1)} \bullet D_{\nu; \nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i, j]} \bullet D_{1; \mu'}^{\beta_3} H_{\Sigma_{i_2}, \Sigma_j} \right|_{i_1, j}^{[\delta_l, 0, \delta_r]} \\ &\leq \text{const} \frac{j-i+1}{M^{|\beta_1|(j-i_1)}} q(\delta_l, \beta_1, 0; \delta_l + \beta_1, \delta_l + \beta_1) \left| H \right|_{i_2, j}^{[0, \beta_3, \delta_r]} \end{aligned}$$

**Proof:** Subbing in the definition of  $K^{(i_1)}$  and applying Lemma II.17,

$$\begin{aligned}
& |D_{\mu;3}^{\beta_1} K^{(i_1)} \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} H_{\Sigma_{i_2}, \Sigma_j} \Big|_{i_1, j}^{[\delta_1, 0, \delta_r]} \\
& \leq |D_{\mu;3}^{\beta_1} F_S^{(i_1)} \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} H_{\Sigma_i, \Sigma_j} \Big|_{i_1, j}^{[\delta_1, 0, \delta_r]} \\
& \quad + \text{const} |D_{\mu;3}^{\beta_1} (G_{C,1}^{(i_1-1)} \bullet \mathcal{C}^{(i_1-1)} \bullet G_{C,2}^{(i_1-1)})^f \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} H_{\Sigma_i, \Sigma_j} \Big|_{i_1-1, j}^{[\delta_1, 0, \delta_r]} \\
& \quad + \text{const} |D_{\mu;3}^{\beta_1} (G_{S,1}^{(i_1-1)} \bullet \mathcal{C}^{(i_1-1)} \bullet G_{S,2}^{(i_1-1)})^f \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} H_{\Sigma_i, \Sigma_j} \Big|_{i_1-1, j}^{[\delta_1, 0, \delta_r]}
\end{aligned}$$

We were able to replace  $H_{\Sigma_{i_2}, \Sigma_j}$  by  $H_{\Sigma_i, \Sigma_j}$  without changing the  $\bullet$  products because  $i \geq i_2$  and  $\mathcal{C}_{\text{loc}}^{[i,j]}$  is supported in the  $i^{\text{th}}$  neighbourhood.

a) By Corollary II.22.b with  $\beta_1 = 0$  and  $\delta_c = \delta$  (and, when  $\text{loc} = \text{top}, \text{bot}$ , Theorem II.19.b with  $h = \frac{1}{M^j |\beta_3|} D_{1;\mu'}^{\beta_3} H_{\Sigma_i, \Sigma_j}$ )

$$\frac{1}{M^j |\delta|} |F_S^{(i_1)} \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} H_{\Sigma_i, \Sigma_j} \Big|_{i_1, j}^{[\delta_1, 0, \delta_r]} \leq \text{const} |F_S^{(i_1)} \Big|_{i_1}^{[\delta_1, 0, 0]} |H \Big|_{i_2, j}^{[0, \beta_3, \delta_r]}$$

provided  $\delta_l, \delta_r, \delta \in \Delta$ . For  $T = C, S$ ,

$$\begin{aligned}
& \frac{1}{M^j |\delta|} |(G_{T,1}^{(i_1-1)} \bullet \mathcal{C}^{(i_1-1)} \bullet G_{T,2}^{(i_1-1)})^f \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} H_{\Sigma_i, \Sigma_j} \Big|_{i_1-1, j}^{[\delta_1, 0, \delta_r]} \\
& \leq \text{const} \sqrt{l_{i_1-1}} |G_{T,1}^{(i_1-1)} \Big|_{i_1-1}^{[\llbracket \delta_l \rrbracket]} |G_{T,2}^{(i_1-1)} \Big|_{i_1-1}^{[\llbracket \delta_l \rrbracket]} |H \Big|_{i_2, j}^{[0, \beta_3, \delta_r]}
\end{aligned}$$

by Corollary II.23.b with  $\ell = i_1 - 1$ ,  $\beta_1 = 0$  and  $\delta_c = \delta$  (and Theorem II.20.b when  $\text{loc} = \text{top}, \text{bot}$ ), provided  $\delta_l, \delta_r, \delta \in \Delta$ .

b) By Corollary II.24.a with  $\delta_c = \delta$ ,  $\ell = i_1$  and  $r = j$  (and Corollary II.22.a when  $\text{loc} = \text{top}, \text{bot}$ )

$$\begin{aligned}
& \frac{1}{M^j |\delta|} |F_S^{(i_1)} \bullet \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\delta} H_{\Sigma_i, \Sigma_j} \Big|_{i_1, j}^{[\delta_1, 0, \delta_r]} \\
& \leq \text{const} \max_{\substack{\alpha_l, \alpha_r \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_l| + |\alpha_r| \leq 3}} |F_S^{(i_1)} \Big|_{i_1}^{[\delta_1, 0, \alpha_r]} \left( |H \Big|_{i_2, j}^{[0, \delta, \delta_r]} + i |H \Big|_{i_2, j}^{[\alpha_l, 0, 0]} \right)
\end{aligned}$$

provided  $\delta_l, \delta_r, \delta \in \Delta$ . For  $T = C, S$ ,

$$\begin{aligned}
& \frac{1}{M^j |\delta|} |(G_{T,1}^{(i_1-1)} \bullet \mathcal{C}^{(i_1-1)} \bullet G_{T,2}^{(i_1-1)})^f \bullet \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\delta} H_{\Sigma_i, \Sigma_j} \Big|_{i_1-1, j}^{[\delta_1, 0, \delta_r]} \\
& \leq \text{const} \sqrt{l_{i_1-1}} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_l \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_l| \leq 3}} |G_{T,1}^{(i_1-1)} \Big|_{i_1-1}^{[\delta_1 + \alpha_{\text{up}}]} |G_{T,2}^{(i_1-1)} \Big|_{i_1-1}^{[\delta_1 + \alpha_{\text{dn}}]} \left( |H \Big|_{i_2, j}^{[0, \delta, \delta_r]} + i |H \Big|_{i_2, j}^{[\alpha_l, 0, 0]} \right)
\end{aligned}$$

by Corollary II.24.b with  $\delta_c = \delta$  and  $\ell = i_1 - 1$  (and Corollary II.23.a when  $\text{loc} = \text{top}, \text{bot}$ ) provided  $\delta_l + \alpha \in \Delta$  for all  $|\alpha| \leq 3$  (which is certainly the case when  $|\delta_l| \leq 3$ ) and  $\delta_r, \delta \in \Delta$ .

c) By Corollary II.22.b, with  $\delta_c = \delta$  (and Theorem II.19.b with  $\beta = \beta_2$ ,  $g = \frac{1}{M^{i_1|\beta_1|}} D_{\mu;3}^{\beta_1} F_S^{(i_1)}$  and  $h = \frac{1}{M^{j|\beta_3|}} D_{1;\mu'}^{\beta_3} H_{\Sigma_i, \Sigma_j}$ , when loc = top, bot)

$$\begin{aligned} & \frac{1}{M^{j|\delta|}} \left| D_{\mu;3}^{\beta_1} F_S^{(i_1)} \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} H_{\Sigma_i, \Sigma_j} \right|_{i_1, j}^{[\delta_1, 0, \delta_r]} \\ & \leq \text{const} \frac{j-i+1}{M^{|\beta_1|(\beta_1-j-i_1)}} \left| F_S^{(i_1)} \right|_{i_1}^{[\delta_1, \beta_1, 0]} \left| H \right|_{i_2, j}^{[0, \beta_3, \delta_r]} \end{aligned}$$

provided  $\delta_1, \delta_r, \delta \in \Delta$ . For  $T = C, S$ ,

$$\begin{aligned} & \frac{1}{M^{j|\delta|}} \left| D_{\mu;3}^{\beta_1} (G_{T,1}^{(i_1-1)} \bullet \mathcal{C}^{(i_1-1)} \bullet G_{T,2}^{(i_1-1)})^f \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} H_{\Sigma_i, \Sigma_j} \right|_{i_1-1, j}^{[\delta_1, 0, \delta_r]} \\ & \leq \text{const} \frac{j-i+1}{M^{|\beta_1|(\beta_1-j-i_1)}} \sqrt{i_1-1} \left| G_{T,1}^{(i_1-1)} \right|_{i_1-1}^{[\delta_1+\beta_1]} \left| G_{T,2}^{(i_1-1)} \right|_{i_1-1}^{[\delta_1+\beta_1]} \left| H \right|_{i_2, j}^{[0, \beta_3, \delta_r]} \end{aligned}$$

by Corollary II.23.b with  $\delta_c = \delta$  and  $\ell = i_1 - 1$  (and Theorem II.20.b when loc = top, bot – the  $D_{\mu;3}^{\beta_1}$  is treated as in the proof of Corollary II.23.a) provided  $\delta_1 + \beta_1, \delta_r, \delta \in \Delta$ . ■

**Proof of Lemma II.26:** The proof is by induction on  $\ell$ . We begin the induction at  $\ell = 1$ . Observe that, by (II.3), Remark II.25 and Lemma II.17,

$$\max_{\delta \in \Delta} |K^{(i_2)}|_{i_2}^{[\delta]} \leq c_F \rho \quad (\text{II.8})$$

for all  $\delta \in \Delta$ , if  $\rho$  is sufficiently small.

a) By Lemma II.27.b, with  $\delta = 0$ ,  $\delta_1 = \alpha$  and  $\delta_r = \delta$ , (II.7), (II.8) and Lemma II.17,

$$\begin{aligned} & \left| K^{(i_1)} \bullet \mathcal{C}_{\text{loc}}^{[i,j]} \bullet K^{(i_2)} \right|_{i_1, j}^{[\alpha, 0, \delta]} \\ & \leq \text{const} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} q(\alpha, 0, \alpha_{\text{up}} + \alpha_{\text{dn}}; \alpha + \alpha_{\text{up}}, \alpha + \alpha_{\text{dn}}) \left[ \left| K^{(i_2)} \right|_{i_2}^{[0, 0, \delta]} + \left| K^{(i_2)} \right|_{i_2}^{[\alpha_1, 0, 0]} \right] \\ & \leq \text{const} c_F^2 \rho^2 \mathbf{v}_{i_1} \frac{i_2}{i_1} \end{aligned}$$

for all of loc = mid, top, bot. Observe that  $\alpha + \alpha_{\text{up}} + \alpha_{\text{dn}} + \alpha_1 \in \Delta$ , since, by hypothesis,  $r_e, r_0 \geq 6$ .

b) By symmetry, we may assume, without loss of generality that  $i_1 \geq i_2$ . Then  $\mathbf{v}_{i_2} \cdots \mathbf{v}_{i_\ell} \min \{ \mathbf{v}_{i_1}, \mathbf{v}_{i_{\ell+1}} \}$  reduces to  $\mathbf{v}_{i_1}$ . By Lemma II.17, Lemma II.27.c, with  $\beta_1 = \beta_2 = \beta_3 = 0$ , (II.7), (II.8) and part a of this Lemma with  $\ell = 1$ ,

$$\begin{aligned} & \left| K^{(i_1)} \bullet \mathcal{C}^{[i,j]} \bullet K^{(i_2)} \right|_{j, j}^{[\delta_1, 0, \delta_r]} \\ & \leq \frac{\text{const}}{M^{j-i}} \left| K^{(i_1)} \bullet \mathcal{C}^{[i,j]} \bullet K^{(i_2)} \right|_{i, j}^{[\delta_1, 0, \delta_r]} + \text{const} \left| K^{(i_1)} \bullet \mathcal{C}^{[i,j]} \bullet K^{(i_2)} \right|_{i, j}^{[0, 0, \delta_r]} \\ & \leq \frac{\text{const} (j-i+1)}{M^{j-i}} q(\delta_1, 0, 0; \delta_1, \delta_1) \left| K^{(i_2)} \right|_{i_2}^{[0, 0, \delta_r]} + \text{const} \left| K^{(i_1)} \bullet \mathcal{C}^{[i,j]} \bullet K^{(i_2)} \right|_{i, j}^{[0, 0, \delta_r]} \\ & \leq \text{const} c_F^2 \rho^2 \mathbf{v}_{i_1} \frac{i_2}{i_1} \leq \text{const} c_F^2 \rho^2 \mathbf{v}_{i_1} \end{aligned}$$

c) Substitute  $\mathcal{C}^{[i,j]} = \mathcal{C}_{\text{top}}^{[i,j]} + \mathcal{C}_{\text{mid}}^{[i,j]} + \mathcal{C}_{\text{bot}}^{[i,j]}$  into

$$D_{\mu;\mu'}^\delta \left[ (K^{(i_1)} \bullet \mathcal{C}^{[i,j]} \bullet K^{(i_2)})_{\Sigma_{i_1}, \Sigma_j} \right] = D_{\mu;\mu'}^\delta \left[ K^{(i_1)} \bullet \mathcal{C}^{[i,j]} \bullet K_{\Sigma_{i_2}, \Sigma_j}^{(i_2)} \right]$$

and apply Leibniz's rule (Lemma II.21) using the routing which gives  $D_{1;3}^{\beta_2} \mathcal{C}_{\text{top}}^{[i,j]}$ ,  $D_{1;3}^{\beta_2} \mathcal{C}_{\text{mid}}^{[i,j]}$  and  $D_{2;4}^{\beta_2} \mathcal{C}_{\text{bot}}^{[i,j]}$ . We define  $k'$  to be  $\frac{1}{M^j|\delta|}$  times the sum of all resulting terms having no derivatives acting on  $K^{(i_1)}$  and  $k''$  to be  $\frac{1}{M^j|\delta|}$  times the sum of all terms having at least one derivative acting on  $K^{(i_1)}$ . Fix any  $\alpha, \alpha'$  and  $\gamma, \gamma'$  obeying,  $|\alpha| \leq 3$ ,  $\gamma + \delta \in \Delta$  and  $\alpha' + \gamma' + \delta \in \Delta$ . We show

$$\begin{aligned} |k'|_{i_1,j}^{[\alpha,0,\gamma]} &\leq \text{const } c_F^2 \rho^2 \mathbf{v}_{i_1} \frac{i_2}{i_1} \\ |k''|_{i_1,j}^{[0,0,\gamma]} &\leq \text{const } c_F^2 \frac{j-i_1+1}{M^{j-i_1}} \rho^2 \mathbf{v}_{i_1} \frac{i_2}{i_1} \end{aligned} \quad (\text{II.9})$$

and

$$|k'|_{j,j}^{[\alpha',0,\gamma']} + |k''|_{j,j}^{[\alpha',0,\gamma']} \leq \text{const } c_F^2 \rho^2 \mathbf{v}_{i_1} \frac{i_2}{i_1} \quad (\text{II.10})$$

Let  $\text{loc} \in \{\text{mid}, \text{top}, \text{bot}\}$ . If  $\text{loc} \in \{\text{top}, \text{mid}\}$ , set  $(\nu, \nu') = (1, 3)$ . If  $\text{loc} = \text{bot}$ , set  $(\nu, \nu') = (2, 4)$ . The contributions to  $k'$  and  $k''$  coming from  $\mathcal{C}_{\text{loc}}^{[i,j]}$  are

$$\begin{aligned} k'_{\text{loc}} &= \frac{1}{M^j|\delta|} \sum_{\substack{\beta_2, \beta_3 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \beta_2 + \beta_3 = \delta}} \binom{\delta}{\beta_2, \beta_3} K^{(i_1)} \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} K_{\Sigma_{i_2}, \Sigma_j}^{(i_2)} \\ k''_{\text{loc}} &= \frac{1}{M^j|\delta|} \sum_{\substack{\beta_1, \beta_2, \beta_3 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \beta_1 + \beta_2 + \beta_3 = \delta \\ |\beta_1| > 0}} \binom{\delta}{\beta_1, \beta_2, \beta_3} D_{\mu;3}^{\beta_1} K^{(i_1)} \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} K_{\Sigma_{i_2}, \Sigma_j}^{(i_2)} \end{aligned}$$

We first bound  $k'_{\text{loc}}$ . Fix  $\beta_2 + \beta_3 = \delta$ . First consider  $\beta_2 \neq 0$ . Let  $(\delta_l, \delta_r) = (\alpha, \gamma)$  or  $(\alpha', \gamma')$ . By Lemma II.27.a,

$$\begin{aligned} \frac{1}{M^j|\delta|} \left| K^{(i_1)} \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} K_{\Sigma_{i_2}, \Sigma_j}^{(i_2)} \right|_{i_1,j}^{[\delta_l, 0, \delta_r]} &\leq \text{const } q(\delta_l, 0, 0; \delta_l, \delta_l) \left| K^{(i_2)} \right|_{i_2}^{[0, \beta_3, \delta_r]} \\ &\leq \text{const } c_F^2 \rho^2 \mathbf{v}_{i_1} \frac{i_2}{i_1} \end{aligned}$$

Next consider  $\beta_2 = 0$ . By Lemma II.27.b, with  $\delta_l = \alpha$  and  $\delta_r = \gamma$ ,

$$\begin{aligned} &\frac{1}{M^j|\delta|} \left| K^{(i_1)} \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} K_{\Sigma_{i_2}, \Sigma_j}^{(i_2)} \right|_{i_1,j}^{[\alpha, 0, \gamma]} \\ &= \frac{1}{M^j|\delta|} \left| K^{(i_1)} \bullet \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^\delta K_{\Sigma_{i_2}, \Sigma_j}^{(i_2)} \right|_{i_1,j}^{[\alpha, 0, \gamma]} \\ &\leq \text{const} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} q(\alpha, 0, \alpha_{\text{up}} + \alpha_{\text{dn}}; \alpha + \alpha_{\text{up}}, \alpha + \alpha_{\text{dn}}) \left[ \left| K^{(i_2)} \right|_{i_2}^{[0, \delta, \gamma]} + \left| K^{(i_2)} \right|_{i_2}^{[\alpha_1, 0, 0]} \right] \\ &\leq \text{const } c_F^2 \rho^2 \mathbf{v}_{i_1} \frac{i_2}{i_1} \end{aligned}$$



and

$$\begin{aligned}
& \frac{1}{M^{j|\delta|}} \left| K^{(i_1)} \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} K_{\Sigma_{i_2}, \Sigma_j}^{(i_2)} \right|_{j,j}^{[\alpha', 0, \gamma']} \\
& \leq \text{const} \frac{1}{M^{j|\delta|}} \left| K^{(i_1)} \bullet \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\delta} K_{\Sigma_{i_2}, \Sigma_j}^{(i_2)} \right|_{i_1,j}^{[0, 0, \gamma']} \\
& \quad + \text{const} \frac{1}{M^{j-i_1}} \frac{1}{M^{j|\delta|}} \left| K^{(i_1)} \bullet \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\delta} K_{\Sigma_{i_2}, \Sigma_j}^{(i_2)} \right|_{i_1,j}^{[\alpha', 0, \gamma']} \\
& \leq \text{const} \frac{1}{M^{j|\delta|}} \left| K^{(i_1)} \bullet \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\delta} K_{\Sigma_{i_2}, \Sigma_j}^{(i_2)} \right|_{i_1,j}^{[0, 0, \gamma']} \\
& \quad + \text{const} \frac{j-i+1}{M^{j-i_1}} q(\alpha', 0, 0; \alpha', \alpha') \left| K^{(i_2)} \right|_{i_2,j}^{[0, \delta, \gamma']} \\
& \leq \text{const} c_F^2 \rho^2 \mathfrak{v}_{i_1} \frac{i_2}{i_1}
\end{aligned}$$

In the first step we applied Lemma II.17. In the second, we applied Lemma II.27.c with  $\beta_1 = \beta_2 = 0$ ,  $\beta_3 = \delta$ ,  $\delta_1 = \alpha'$  and  $\delta_r = \gamma'$ . In the third step we applied the conclusion of the last estimate and Lemma II.17.

To bound  $|k''_{\text{loc}}|_{i_1,j}^{[\delta_1, 0, \delta_r]}$  with  $\delta_1 + \delta_r + \delta \in \Delta$  observe that, by Lemma II.27.c, for all  $\beta_1 + \beta_2 + \beta_3 = \delta$  with  $\beta_1 \neq 0$ ,

$$\begin{aligned}
& \frac{1}{M^{j|\delta|}} \left| D_{\mu;3}^{\beta_1} K^{(i_1)} \bullet D_{\nu;\nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1;\mu'}^{\beta_3} K_{\Sigma_{i_2}, \Sigma_j}^{(i_2)} \right|_{i_1,j}^{[\delta_1, 0, \delta_r]} \\
& \leq \text{const} \frac{j-i+1}{M^{|\beta_1|(j-i_1)}} q(\delta_1, \beta_1, 0; \delta_1 + \beta_1, \delta_1 + \beta_1) \left| K^{(i_2)} \right|_{i_2,j}^{[0, \beta_3, \delta_r]} \\
& \leq \text{const} \frac{j-i_1+1}{M^{j-i_1}} c_F^2 \rho^2 \mathfrak{v}_{i_1} \frac{i_2}{i_1}
\end{aligned}$$

Setting  $(\delta_1, \delta_r) = (0, \gamma)$ , we get the  $k''$  estimate of (II.9). Setting  $(\delta_1, \delta_r) = (\alpha', \gamma')$  and using Lemma II.17, we get the  $k''$  estimate of (II.10).

When  $\ell = 1$ , part c follows from (II.9).

d) Again, we may assume, without loss of generality that  $i_1 \geq i_2$ . By part c and (II.10)

$$\begin{aligned}
& \frac{1}{M^{j|\delta_c|}} \left| D_{\mu;\mu'}^{\delta_c} \left[ K^{(i_1)} \bullet \mathcal{C}^{[i,j]} \bullet K_{\Sigma_{i_2}, \Sigma_j}^{(i_2)} \right] \right|_{j,j}^{[\delta_1, 0, \delta_r]} \leq |k'|_{j,j}^{[\delta_1, 0, \delta_r]} + |k''|_{j,j}^{[\delta_1, 0, \delta_r]} \\
& \leq \text{const} c_F^2 \rho^2 \mathfrak{v}_{i_1}
\end{aligned}$$

This finishes the case  $\ell = 1$ .

**Induction step:** We assume that the Lemma holds for  $\ell - 1$ . Write

$$K^{(i_1)} \bullet \mathcal{C}^{[i,j]} \bullet K^{(i_2)} \bullet \dots \bullet K^{(i_{\ell+1})} = K^{(i_1)} \bullet \mathcal{C}^{[i,j]} \bullet H$$

with

$$H = K^{(i_2)} \bullet \mathcal{C}^{[\max\{i_2, i_3\}, j]} \bullet \dots \bullet K^{(i_{\ell+1})}$$

Set

$$\mathfrak{V} = \text{const}^{\ell-1} (c_F \rho)^\ell \mathbf{v}_{i_2} \cdots \mathbf{v}_{i_\ell}$$

The induction hypothesis applies to  $H$ . So, for all  $|\alpha| \leq 3$  and  $\delta \in \Delta$

$$|H|_{i_2, j}^{[\alpha, 0, \delta]} \leq \mathfrak{V} \frac{i_{\ell+1}}{i_2}$$

Furthermore, for each  $0 \neq \delta \in \Delta$ ,  $\mu \in \{1, 2\}$  and  $\mu' \in \{3, 4\}$ , there is a decomposition

$$\frac{1}{M^{j|\delta|}} D_{\mu; \mu'}^\delta H_{\Sigma_{i_2}, \Sigma_j} = h'_{\delta, \mu, \mu'} + h''_{\delta, \mu, \mu'}$$

with, for all  $|\alpha| \leq 3$  and all  $\gamma$  with  $\gamma + \delta \in \Delta$ ,

$$\begin{aligned} |h'_{\delta, \mu, \mu'}|_{i_2, j}^{[\alpha, 0, \gamma]} &\leq \mathfrak{V} \frac{i_{\ell+1}}{i_2} \\ |h''_{\delta, \mu, \mu'}|_{i_2, j}^{[0, 0, \gamma]} &\leq \frac{j-i_2+1}{M^{j-i_2}} \mathfrak{V} \frac{i_{\ell+1}}{i_2} \end{aligned}$$

In particular,

$$|H|_{i_2, j}^{[0, \delta_c, \delta_r]} \leq 2\mathfrak{V} \frac{i_{\ell+1}}{i_2}$$

for  $\delta_c + \delta_r \in \Delta$ .

a) By Lemma II.27.b, with  $\delta = 0$ ,  $\delta_l = \alpha$  and  $\delta_r = \delta$ , (II.7) and Lemma II.17,

$$\begin{aligned} \left| K^{(i_1)} \bullet \mathcal{C}^{[i, j]} \bullet H \right|_{i_1, j}^{[\alpha, 0, \delta]} &\leq \text{const} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_l \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_l| \leq 3}} q(\alpha, 0, \alpha_{\text{up}} + \alpha_{\text{dn}}; \alpha + \alpha_{\text{up}}, \alpha + \alpha_{\text{dn}}) |H|_{i_2, j}^{[\alpha_l, 0, \delta]} \\ &\leq \text{const } c_F \rho \mathbf{v}_{i_1} \frac{i_2}{i_1} \mathfrak{V} \frac{i_{\ell+1}}{i_2} \leq \text{const } c_F \rho \mathbf{v}_{i_1} \mathfrak{V} \frac{i_{\ell+1}}{i_1} \end{aligned}$$

b) Again, we may assume, without loss of generality that  $i_1 \geq i_{\ell+1}$ . Then the factor  $\text{const}^\ell (c_F \rho)^{\ell+1} \mathbf{v}_{i_2} \cdots \mathbf{v}_{i_\ell} \min \{ \mathbf{v}_{i_1}, \mathbf{v}_{i_{\ell+1}} \}$  in the right hand side of the statement reduces to  $\text{const } c_F \rho \mathbf{v}_{i_1} \mathfrak{V}$ . The remainder of the proof is virtually identical to that for  $\ell = 1$ .

c) Substitute  $\mathcal{C}^{[i, j]} = \mathcal{C}_{\text{top}}^{[i, j]} + \mathcal{C}_{\text{mid}}^{[i, j]} + \mathcal{C}_{\text{bot}}^{[i, j]}$  into

$$D_{\mu; \mu'}^\delta \left[ (K^{(i_1)} \bullet \mathcal{C}^{[i, j]} \bullet H)_{\Sigma_{i_1}, \Sigma_j} \right] = D_{\mu; \mu'}^\delta \left[ K^{(i_1)} \bullet \mathcal{C}^{[i, j]} \bullet H_{\Sigma_{i_2}, \Sigma_j} \right]$$

and apply Leibniz's rule (Lemma II.21) using the routing which gives  $D_{1;3}^{\beta_2} \mathcal{C}_{\text{top}}^{[i, j]}$ ,  $D_{1;3}^{\beta_2} \mathcal{C}_{\text{mid}}^{[i, j]}$  and  $D_{2;4}^{\beta_2} \mathcal{C}_{\text{bot}}^{[i, j]}$ . We define  $k'$  to be  $\frac{1}{M^{j|\delta|}}$  times the sum of all resulting terms having no derivatives acting on  $K^{(i_1)}$  and  $k''$  to be  $\frac{1}{M^{j|\delta|}}$  times the sum of all terms having at least one derivative acting on  $K^{(i_1)}$ . Fix any  $\alpha, \alpha'$  and  $\gamma, \gamma'$  obeying,  $|\alpha| \leq 3$ ,  $\gamma + \delta \in \Delta$  and  $\alpha' + \gamma' + \delta \in \Delta$ . We show

$$\begin{aligned} |k'|_{i_1, j}^{[\alpha, 0, \gamma]} &\leq \text{const } c_F \rho \mathbf{v}_{i_1} \mathfrak{V} \frac{i_{\ell+1}}{i_1} \\ |k''|_{i_1, j}^{[0, 0, \gamma]} &\leq \text{const } c_F \frac{j-i_1+1}{M^{j-i_1}} \rho \mathbf{v}_{i_1} \mathfrak{V} \frac{i_{\ell+1}}{i_1} \end{aligned} \tag{II.11}$$

and

$$|k'|_{j,j}^{[\alpha',0,\gamma']} + |k''|_{j,j}^{[\alpha',0,\gamma']} \leq \text{const } c_F \rho \mathbf{v}_{i_1} \mathfrak{V} \frac{i_{\ell+1}}{i_1} \quad (\text{II.12})$$

Let  $\text{loc} \in \{\text{mid}, \text{top}, \text{bot}\}$ . If  $\text{loc} \in \{\text{top}, \text{mid}\}$ , set  $(\nu, \nu') = (1, 3)$ . If  $\text{loc} = \text{bot}$ , set  $(\nu, \nu') = (2, 4)$ . The contributions to  $k'$  and  $k''$  coming from  $\mathcal{C}_{\text{loc}}^{[i,j]}$  are

$$\begin{aligned} k'_{\text{loc}} &= \frac{1}{M^{j|\delta|}} \sum_{\substack{\beta_2, \beta_3 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \beta_2 + \beta_3 = \delta}} \binom{\delta}{\beta_2, \beta_3} K^{(i_1)} \bullet D_{\nu; \nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1; \mu'}^{\beta_3} H_{\Sigma_{i_2}, \Sigma_j} \\ k''_{\text{loc}} &= \frac{1}{M^{j|\delta|}} \sum_{\substack{\beta_1, \beta_2, \beta_3 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \beta_1 + \beta_2 + \beta_3 = \delta \\ |\beta_1| > 0}} \binom{\delta}{\beta_1, \beta_2, \beta_3} D_{\mu; 3}^{\beta_1} K^{(i_1)} \bullet D_{\nu; \nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1; \mu'}^{\beta_3} H_{\Sigma_{i_2}, \Sigma_j} \end{aligned}$$

We first bound  $k'_{\text{loc}}$ . Fix  $\beta_2 + \beta_3 = \delta$ . First consider  $\beta_2 \neq 0$ . Let  $(\delta_1, \delta_r) = (\alpha, \gamma)$  or  $(\alpha', \gamma')$ . By Lemma II.27.a,

$$\begin{aligned} \frac{1}{M^{j|\delta|}} \left| K^{(i_1)} \bullet D_{\nu; \nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1; \mu'}^{\beta_3} H_{\Sigma_{i_2}, \Sigma_j} \right|_{i_1, j}^{[\delta_1, 0, \delta_r]} &\leq \text{const } q(\delta_1, 0, 0; \delta_1, \delta_1) |H|_{i_2, j}^{[0, \beta_3, \delta_r]} \\ &\leq \text{const } c_F \rho \mathbf{v}_{i_1} \frac{i_2}{i_1} \mathfrak{V} \frac{i_{\ell+1}}{i_2} \\ &\leq \text{const } c_F \rho \mathbf{v}_{i_1} \mathfrak{V} \frac{i_{\ell+1}}{i_1} \end{aligned}$$

Next consider  $\beta_2 = 0$ . By (II.7),

$$\begin{aligned} \frac{1}{M^{j|\delta|}} \left| K^{(i_1)} \bullet D_{\nu; \nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1; \mu'}^{\beta_3} H_{\Sigma_{i_2}, \Sigma_j} \right|_{i_1, j}^{[\alpha, 0, \gamma]} &= \frac{1}{M^{j|\delta|}} \left| K^{(i_1)} \bullet \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1; \mu'}^{\delta} H_{\Sigma_{i_2}, \Sigma_j} \right|_{i_1, j}^{[\alpha, 0, \gamma]} \\ &\leq \left| K^{(i_1)} \bullet \mathcal{C}_{\text{loc}}^{[i,j]} \bullet h'_{\delta, 1, \mu'} \right|_{i_1, j}^{[\alpha, 0, \gamma]} + \left| K^{(i_1)} \bullet \mathcal{C}_{\text{loc}}^{[i,j]} \bullet h''_{\delta, 1, \mu'} \right|_{i_1, j}^{[\alpha, 0, \gamma]} \\ &\leq \text{const} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} q(\alpha, 0, \alpha_{\text{up}} + \alpha_{\text{dn}}; \alpha + \alpha_{\text{up}}, \alpha + \alpha_{\text{dn}}) |h'_{\delta, 1, \mu'}|_{i_2, j}^{[\alpha_1, 0, \gamma]} \\ &\quad + \text{const } q(\alpha, 0, 0; \alpha, \alpha) (j - i + 1) |h''_{\delta, 1, \mu'}|_{i_2, j}^{[0, 0, \gamma]} \\ &\leq \text{const } c_F \rho \mathbf{v}_{i_1} \mathfrak{V} \frac{i_{\ell+1}}{i_1} \end{aligned}$$

since  $\frac{(j-i+1)(j-i_2-1)}{M^{j-i_2}} \leq \text{const}$ . The term with  $h'_{\delta, 1, \mu'}$  was bounded using Lemma II.27.b with  $\delta = 0$ . The term with  $h''_{\delta, 1, \mu'}$  was bounded using Lemma II.27.c with  $\beta_1 = \beta_2 = \beta_3 = 0$ . Again, with  $\beta_2 = 0$ ,

$$\frac{1}{M^{j|\delta|}} \left| K^{(i_1)} \bullet D_{\nu; \nu'}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet D_{1; \mu'}^{\beta_3} H_{\Sigma_{i_2}, \Sigma_j} \right|_{j, j}^{[\alpha', 0, \gamma']} \leq \text{const } c_F \rho \mathbf{v}_{i_1} \mathfrak{V} \frac{i_{\ell+1}}{i_1}$$

as in the proof of part (c) when  $\ell = 1$ .

We bound  $|k''_{\text{loc}}|_{i_1, j}^{[\delta_1, 0, \delta_r]}$  with  $\delta_1 + \delta_r + \delta \in \Delta$  as for  $\ell = 1$ . Observe that, by Lemma II.27.c, for all  $\beta_1 + \beta_2 + \beta_3 = \delta$  with  $\beta_1 \neq 0$ ,

$$\begin{aligned} & \frac{1}{M^{j|\delta|}} \left| D_{\mu; 3}^{\beta_1} K^{(i_1)} \bullet D_{1, 3}^{\beta_2} \mathcal{C}_{\text{loc}}^{[i, j]} \bullet D_{1; \mu'}^{\beta_3} H_{\Sigma_{i_2}, \Sigma_j} \right|_{i_1, j}^{[\delta_1, 0, \delta_r]} \\ & \leq \text{const} \frac{j-i+1}{M^{|\beta_1|(j-i_1)}} q(\delta_1, \beta_1, 0; \delta_1 + \beta_1, \delta_1 + \beta_1) |H|_{i_2, j}^{[0, \beta_3, \delta_r]} \\ & \leq \text{const} \frac{j-i_1+1}{M^{j-i_1}} c_F \rho \mathbf{v}_{i_1} \mathfrak{V} \frac{i_{\ell+1}}{i_1} \end{aligned}$$

Setting  $(\delta_1, \delta_r) = (0, \gamma)$ , we get the  $k''$  estimate of (II.11). Setting  $(\delta_1, \delta_r) = (\alpha', \gamma')$  and using Lemma II.17, we get the  $k''$  estimate of (II.12).

d) Part (d) follows from part (c) and (II.12) as in the case  $\ell = 1$ . ■

### Completion of the proof of Theorem I.20:

We prove (II.5). Let  $\vec{\delta} = (\delta_1, \delta_c, \delta_r) \in \vec{\Delta}$ . By parts b) and d) of the Lemma above, for  $\ell \geq 1$ ,

$$\left| K^{(i_1)} \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet K^{(i_2)} \bullet \dots \bullet K^{(i_{\ell+1})} \right|_{j, j}^{[\vec{\delta}]} \leq \text{const}^\ell (c_F \rho)^{\ell+1} \mathbf{v}_{i_2} \cdots \mathbf{v}_{i_\ell} \min \{ \mathbf{v}_{i_1}, \mathbf{v}_{i_{\ell+1}} \}$$

Therefore, by Corollary II.12.i,

$$\begin{aligned} |\mathcal{L}_S^{(j+1)}|_{j, j}^{[\vec{\delta}]} & \leq \sum_{\ell=1}^{\infty} \sum_{i_1, \dots, i_{\ell+1}=2}^j \text{const}^\ell (c_F \rho)^{\ell+1} \mathbf{v}_{i_2} \cdots \mathbf{v}_{i_\ell} \min \{ \mathbf{v}_{i_1}, \mathbf{v}_{i_{\ell+1}} \} \\ & \leq \text{const} c_F^2 \rho^2 \sum_{\ell=1}^{\infty} \left( (\text{const} c_F \rho)^{\ell-1} \sum_{i_2, \dots, i_\ell=2}^{\infty} \mathbf{v}_{i_2} \cdots \mathbf{v}_{i_\ell} \right) \left( \sum_{i_1, i_{\ell+1}=2}^{\infty} \min \{ \mathbf{v}_{i_1}, \mathbf{v}_{i_{\ell+1}} \} \right) \\ & \leq \text{const} c_F^2 \rho^2 \sum_{\ell=1}^{\infty} (\text{const} c_F \rho)^{\ell-1} \left( \sum_{i_1 \geq i_{\ell+1}} \mathbf{v}_{i_1} + \sum_{i_1 < i_{\ell+1}} \mathbf{v}_{i_{\ell+1}} \right) \\ & \leq \text{const} c_F^2 \rho^2 \sum_{i=2}^{\infty} (i-1) \mathbf{v}_i \leq \text{const} c_F^2 \rho^2 = c_{\mathcal{L}} \rho^2 \end{aligned}$$

when  $\rho$  is small enough. This concludes the induction step in the proof of Theorem I.20. ■

### The Infrared Limit

Define, for each  $j \geq 2$ ,  $\ell \geq 1$  and  $i_1, \dots, i_{\ell+1} \geq 2$ , the function

$$\mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}^{(j)} : \mathbb{M}^3 \times \{\uparrow, \downarrow\}^4 \rightarrow \mathbb{C}$$

by

$$\begin{aligned} & \mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t, \sigma_1, \dots, \sigma_4) \\ & = \left[ (F^{(i_1)} + L^{(i_1)} f) \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet (F^{(i_2)} + L^{(i_2)} f) \bullet \dots \right. \\ & \quad \left. \dots \bullet \mathcal{C}^{[\max\{i_\ell, i_{\ell+1}\}, j]} \bullet (F^{(i_{\ell+1})} + L^{(i_{\ell+1})} f) \right]_{i_1, i_2, i_3, i_4=0}^{((q+\frac{t}{2}, \sigma_1), (q-\frac{t}{2}, \sigma_2), (q'+\frac{t}{2}, \sigma_3), (q'-\frac{t}{2}, \sigma_4))} \end{aligned}$$

By Proposition II.3.ii

$$\begin{aligned} \mathcal{L}^{(j+1)} \Big|_{i_1, i_2, i_3, i_4=0}^{((q+\frac{t}{2}, \sigma_1), (q-\frac{t}{2}, \sigma_2), (q'+\frac{t}{2}, \sigma_3), (q'-\frac{t}{2}, \sigma_4))} \\ = \sum_{\ell=1}^{\infty} \sum_{i_1, \dots, i_{\ell+1}=2}^j \mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t, \sigma_1, \dots, \sigma_4) \end{aligned}$$

**Lemma II.28**

$$\sum_{\ell=1}^{\infty} \sum_{i_1, \dots, i_{\ell+1}=2}^{\infty} \sup_{j \geq \max\{i_1, \dots, i_{\ell+1}\}} \sup_{\substack{q, q', t \in \mathbb{M} \\ \sigma_i \in \{\uparrow, \downarrow\}}} \left| \mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t, \sigma_1, \dots, \sigma_4) \right| < \infty$$

**Proof:** By Lemma II.26.ii, with  $\delta_l = \delta_r = 0$ , and the analogous bound for  $\mathcal{L}_C^{(j+1)}$ ,

$$\sup_{\substack{q, q', t \in \mathbb{M} \\ \sigma_i \in \{\uparrow, \downarrow\}}} \left| \mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t, \sigma_1, \dots, \sigma_4) \right| \leq \text{const}^\ell (c_F \rho)^{\ell+1} \mathfrak{v}_{i_2} \cdots \mathfrak{v}_{i_\ell} \min \{ \mathfrak{v}_{i_1}, \mathfrak{v}_{i_{\ell+1}} \}$$

uniformly in  $j$ . Hence, as in the final part of the proof of Theorem I.20,

$$\begin{aligned} \sum_{\ell=1}^{\infty} \sum_{i_1, \dots, i_{\ell+1}=2}^{\infty} \sup_{j \geq \max\{i_1, \dots, i_{\ell+1}\}} \sup_{\substack{q, q', t \in \mathbb{M} \\ \sigma_i \in \{\uparrow, \downarrow\}}} \left| \mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t, \sigma_1, \dots, \sigma_4) \right| \\ \leq \text{const } c_F^2 \rho^2 \sum_{\ell=1}^{\infty} \left( (\text{const } c_F \rho)^{\ell-1} \sum_{i_2, \dots, i_\ell=2}^{\infty} \mathfrak{v}_{i_2} \cdots \mathfrak{v}_{i_\ell} \right) \left( \sum_{i_1, i_{\ell+1}=2}^{\infty} \min \{ \mathfrak{v}_{i_1}, \mathfrak{v}_{i_{\ell+1}} \} \right) \\ \leq \text{const } c_F^2 \rho^2 \sum_{\ell=1}^{\infty} (\text{const } c_F \rho)^{\ell-1} \left( \sum_{i_1 \geq i_{\ell+1}} \mathfrak{v}_{i_1} + \sum_{i_1 < i_{\ell+1}} \mathfrak{v}_{i_{\ell+1}} \right) \\ \leq \text{const } c_F^2 \rho^2 \sum_{i=2}^{\infty} (i-1) \mathfrak{v}_i < \infty \end{aligned}$$

when  $\rho$  is small enough. ■

**Lemma II.29** *For  $t \neq 0$ , the limit*

$$\mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}(q, q', t, \sigma_1, \dots, \sigma_4) = \lim_{j \rightarrow \infty} \mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t, \sigma_1, \dots, \sigma_4)$$

*exists. The limit is continuous in  $(q, q', t)$  for  $t \neq 0$ . The restrictions to  $\mathfrak{t} = 0$  and to  $t_0 = 0$ , namely,  $\mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}(q, q', (t_0, \mathbf{0}), \sigma_1, \dots, \sigma_4)$  and  $\mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}(q, q', (0, \mathfrak{t}), \sigma_1, \dots, \sigma_4)$ , have continuous extensions to  $t = 0$ .*

**Proof:** It suffices to consider separately the spin and charge parts, in the sense of Lemma II.8, of  $\mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t, \sigma_1, \dots, \sigma_4)$ . We denote them  $\mathfrak{L}_{X, \ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t)$  with  $X = S, C$ . The existence and continuity of the limits when  $t \neq 0$  shall be proven in Lemma III.29.

Recall that the bubble propagator  $\mathcal{C}^{[i, j]}$  has momentum space kernel

$$\mathcal{C}^{[i, j]}(p, k) = \frac{\nu^{(\geq i)}(p)\nu^{(\geq i)}(k) - \nu^{(\geq j+1)}(p)\nu^{(\geq j+1)}(k)}{[ip_0 - e'(p)][ik_0 - e'(k)]}$$

where  $e'(k) = e(\mathbf{k}) - v(k)$ . Define the model particle-hole bubble propagators

$$\begin{aligned} \mathcal{A}_{i, j}(p, k) &= \frac{\nu^{(\geq i)}(p)\nu^{(\geq i)}(k)[1 - \nu_j(e(\mathbf{p}))\nu_j(e(\mathbf{k}))]}{[ip_0 - e'(p)][ik_0 - e'(k)]} \\ \mathcal{B}_{i, j}(p, k) &= \frac{\nu^{(\geq i)}(p)\nu^{(\geq i)}(k)[1 - \nu_j(p_0)\nu_j(k_0)]}{[ip_0 - e'(p)][ik_0 - e'(k)]} \end{aligned} \quad (\text{II.13})$$

where

$$\nu_j(\omega) = \sum_{m=j}^{\infty} \nu(M^{2m}\omega^2)$$

with  $\nu$  being the single scale cutoff introduced in Definition I.2. Let

$$\begin{aligned} A_{X, \ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t_0) \\ = [K_X^{(i_{\ell+1})} \bullet \mathcal{A}_{\max\{i_{\ell+1}, i_{\ell}\}, j} \bullet K_X^{(i_{\ell})} \bullet \dots \bullet \mathcal{A}_{\max\{i_2, i_1\}, j} \bullet K_X^{(i_1)}]_{i_1, i_2, i_3, i_4=0} \left( q + \frac{t}{2}, q - \frac{t}{2}, q' + \frac{t}{2}, q' - \frac{t}{2} \right) \Big|_{t=0} \end{aligned}$$

and

$$\begin{aligned} B_{X, \ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', \mathbf{t}) \\ = [K_X^{(i_{\ell+1})} \bullet \mathcal{B}_{\max\{i_{\ell+1}, i_{\ell}\}, j} \bullet K_X^{(i_{\ell})} \bullet \dots \bullet \mathcal{B}_{\max\{i_2, i_1\}, j} \bullet K_X^{(i_1)}]_{i_1, i_2, i_3, i_4=0} \left( q + \frac{t}{2}, q - \frac{t}{2}, q' + \frac{t}{2}, q' - \frac{t}{2} \right) \Big|_{t_0=0} \end{aligned}$$

where  $K_X^{(i)}$  is  $F_S^{(i)} + \frac{1}{2}L_{C\Sigma_i}^{(i)f} - \frac{1}{2}L_{S\Sigma_i}^{(i)f}$  when  $X = S$  and  $F_C^{(i)} + \frac{1}{2}L_{C\Sigma_i}^{(i)f} + \frac{3}{2}L_{S\Sigma_i}^{(i)f}$  when  $X = C$ . By Corollary III.31 the differences

$$\mathcal{L}_{X, \ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', (t_0, \mathbf{0})) - A_{X, \ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t_0)$$

and

$$\mathcal{L}_{X, \ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', (0, \mathbf{t})) - B_{X, \ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', \mathbf{t})$$

both converge to zero for all  $t \neq 0$ . The bounds on the  $|K_X^{(i_m)}|_{i_m, i_m}$ 's required by Corollary III.31 are provided by (II.8) with  $\delta = 0$ .

That  $\lim_{j \rightarrow \infty} A_{X, \ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t_0)$  and  $\lim_{j \rightarrow \infty} B_{X, \ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', \mathbf{t})$  exist and are continuous at  $t = 0$  is proven using Lemma B.3 inductively on  $\ell$ , with  $I = F$ , the full Fermi surface. For the induction step from  $\ell - 1$  to  $\ell$ , set  $z = (q, q')$  and use

$$\begin{aligned} u_j(k, t_0, z) &= K_X^{(i_{\ell+1})} \left( q + \frac{t}{2}, q - \frac{t}{2}, k + \frac{t}{2}, k - \frac{t}{2} \right) \Big|_{t=0} \nu^{(\geq i)}(k+t) \nu^{(\geq i)}(k) A_{\ell-1, i_1, \dots, i_{\ell}}^{(j)}(k, q', t_0) \\ v_j(k, \mathbf{t}, z) &= K_X^{(i_{\ell+1})} \left( q + \frac{t}{2}, q - \frac{t}{2}, k + \frac{t}{2}, k - \frac{t}{2} \right) \Big|_{t_0=0} \nu^{(\geq i)}(k+t) \nu^{(\geq i)}(k) B_{\ell-1, i_1, \dots, i_{\ell}}^{(j)}(k, q', \mathbf{t}) \\ n_j(\omega) &= \nu_{i-1}(\omega) [1 - \nu_j(\omega)^2] \\ i &= \max\{i_{\ell+1}, i_{\ell}\} \end{aligned}$$

Also fix some  $0 < \aleph'' < \aleph$ , and use  $\tilde{\aleph} = \aleph'' + \frac{1}{2^\ell}(\aleph - \aleph'')$  and  $\aleph' = \aleph'' + \frac{1}{2^{\ell+1}}(\aleph - \aleph'')$ . ■

**Proof of Theorem I.22:** By the Lebesgue dominated convergence theorem and the uniform bounds of Lemma II.28, the existence of the limit  $\lim_{j \rightarrow \infty}$  and its continuity for  $t \neq 0$ , as well as the existence and continuity of the limits  $\lim_{t_0 \rightarrow 0} \lim_{j \rightarrow \infty}$  and  $\lim_{t \rightarrow 0} \lim_{j \rightarrow \infty}$  applied to  $\mathcal{L}^{(j+1)} \Big|_{i_1, i_2, i_3, i_4=0}((q+\frac{t}{2}, \sigma_1), (q-\frac{t}{2}, \sigma_2), (q'+\frac{t}{2}, \sigma_3), (q'-\frac{t}{2}, \sigma_4))$  follow from the corresponding properties of  $\mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t, \sigma_1, \dots, \sigma_4)$ , for  $\ell \geq 1$  and  $i_1, \dots, i_{\ell+1} \geq 2$ . These were proven in Lemma II.29. ■

### III. Bubbles

The bulk of this section is devoted to the proof of Theorem II.19. Parts b and c, reformulated as Theorem III.9, are relatively easy to prove. To do so, we fully decompose

$$\mathcal{C}^{[i,j]} = \sum_{m=i}^j \sum_{\substack{m_1, m_2 \in \mathbb{N}_0 \\ \min\{m_1, m_2\} = m}} C_v^{(m_1)} \otimes C_v^{(m_2) t} \quad (\text{III.1})$$

and bound each term naively to achieve ordinary power counting. The factor  $j - i + 1 = \sum_{m=i}^j 1$  in the first statement of part c is a reflection of the marginality of four-legged diagrams in naive power counting. When power counting bubbles with propagator  $D_{\mu, \mu'}^\beta \mathcal{C}^{[i,j]}$ ,  $|\beta| \geq 1$ , the sum  $\sum_{m=i}^j 1$  is replaced by  $\sum_{m=i}^j M^{|\beta|m} \leq \text{const } M^{|\beta|j}$ , which is cancelled by the factors  $\frac{1}{M^{|\beta|j}}$  on the left hand sides of parts b and c. In the  $\beta = 0$  statement of part b, naive power counting gives  $\sum_{i_t=i}^j \sum_{i_b > j} M^{-(i_b - i_t)} \leq \text{const}$ .

The proof of Theorem II.19a, which follows Theorem III.15, relies on two distinct phenomena, volume improvement for large transfer momentum and a sign cancellation in momentum space for small transfer momentum. The mechanism underlying the sign cancellation has been illustrated in the model Lemma I.1 and is fully implemented in Theorem III.15.

We now sketch the idea behind volume improvement. To unravel the sector sums of the  $\bullet$  product of Definition I.8, we define, for any translation invariant functions  $K$  on  $\mathfrak{Y}_\Sigma^2 \times (\mathbb{R} \times \mathbb{R}^2)$ ,  $K'$  on  $(\mathbb{R} \times \mathbb{R}^2) \times \mathfrak{Y}_{\Sigma'}^2$ , and bubble propagator  $P$ ,

$$\begin{aligned} K \circ P(y_1, y_2, x_3, x_4) &= \int dx_1 dx_2 K(y_1, y_2, x_1, x_2) P(x_1, x_2, x_3, x_4) \\ P \circ K'(x_1, x_2, y_3, y_4) &= \int dx_3 dx_4 P(x_1, x_2, x_3, x_4) K'(x_3, x_4, y_3, y_4) \end{aligned}$$

If at least one of  $y_1, y_2, y_3, y_4$  is in  $(\mathbb{R} \times \mathbb{R}^2) \times \Sigma$  or  $(\mathbb{R} \times \mathbb{R}^2) \times \Sigma'$

$$K \circ K'(y_1, y_2, y_3, y_4) = \int dx_1 dx_2 K(y_1, y_2, x_1, x_2) K'(x_1, x_2, y_3, y_4)$$

On the other hand, if all of  $k_1, k_2, k_3, k_4$  are in  $\mathbb{M}$ ,  $K \circ K'(k_1, k_2, k_3, k_4)$  is determined by

$$K \circ K'(k_1, k_2, k_3, k_4) (2\pi)^3 \delta(k_1 - k_2 - k_3 + k_4) = \int dx_1 dx_2 K(k_1, k_2, x_1, x_2) K'(x_1, x_2, k_3, k_4)$$



or equivalently, by

$$K \circ K'(k_1, k_2, k_3, k_4) = \int dx_n \quad K(k_1, k_2, x_1, x_2) K'(x_1, x_2, k_3, k_4) \Big|_{x_3-n=0}$$

for  $n \in \{1, 2\}$ . Then, for the functions  $g$  and  $h$  of the Theorem,

$$(g \bullet \mathcal{C}^{[i,j]} \bullet h)(y_1, y_2, y_3, y_4) = \sum_{\substack{u_1, u_2 \in \Sigma_i \\ v_1, v_2 \in \Sigma_i}} g(y_1, y_2, (\cdot, u_1), (\cdot, u_2)) \circ \mathcal{C}^{[i,j]} \circ h((\cdot, v_1), (\cdot, v_2), y_3, y_4) \quad (\text{III.2})$$

Consider the case in which all external arguments  $y_1, \dots, y_4$  are momenta  $k_1, \dots, k_4$ . Then

$$\begin{aligned} & (g \bullet \mathcal{C}^{[i,j]} \bullet h)(k_1, k_2, k_3, k_4) \\ &= \frac{1}{(2\pi)^3} \int d^3 p d^3 k \sum_{\substack{u_1, u_2 \in \Sigma_i \\ v_1, v_2 \in \Sigma_i}} \delta(k_1 - k_2 - p + k) \check{g}(k_1, k_2, (p, u_1), (k, u_2)) \mathcal{C}^{[i,j]}(p, k) \check{h}((p, v_1), (k, v_2), k_3, k_4) \end{aligned} \quad (\text{III.3})$$

where

$$\mathcal{C}^{[i,j]}(p, k) = \sum_{m=i}^j \sum_{\substack{m_1, m_2 \in \mathbb{N}_0 \\ \min\{m_1, m_2\} = m}} C_v^{(m_1)}(p) C_v^{(m_2)}(k)$$

In order for  $\mathcal{C}^{[i,j]}(p, k)$  to be nonzero, one must have  $p$  and  $k$  in the  $i^{\text{th}}$  neighbourhood. In particular,  $\mathbf{p}$  and  $\mathbf{k}$  must lie within a distance  $\frac{\text{const}}{M^i}$  of the Fermi curve  $F$ . Furthermore, by conservation of momentum at the vertex  $g$ , the “transfer momentum”

$$t = k_1 - k_2$$


is equal to  $p - k$ . Thus, the set of pairs  $(p, k)$  for which the integrand of (III.3) does not vanish is contained in

$$\{ (k, p) \in (\text{supp } C^{(\geq i)})^2 \mid p - k = t \}$$

For each fixed large  $\mathbf{t}$ , the volume of

$$\{ \mathbf{p} \in \text{supp } C^{(\geq i)} \mid \mathbf{p} - \mathbf{t} \in \text{supp } C^{(\geq i)} \} = \text{supp } C^{(\geq i)} \cap (\mathbf{t} + \text{supp } C^{(\geq i)}) \quad (\text{III.4})$$

is very small compared to the volume of  $\text{supp } C^{(\geq i)}$ , as the following figure illustrates.



In naive power counting, the volume of the set (III.4) is bounded by the volume of  $\text{supp } C^{(\geq i)}$ , yielding a relatively loose bound. There is a similar volume improvement, when, for example, the external arguments  $y_1 = (x_1, s_1)$  and  $y_2 = (x_2, s_2)$  lie  $(\mathbb{R} \times \mathbb{R}^2) \times \Sigma_\ell$  and the sectors  $s_1$  and  $s_2$  are widely separated. For a more detailed discussion of this volume improvement in perturbation theory see [FKLT2].

We now give a somewhat more detailed technical outline of the contents of this section. By sector counting and relatively simple propagator estimates, the volume improvement effect can be implemented for all summands

$$\mathcal{C}^{(m)} = \sum_{\substack{m_1, m_2 \in \mathbb{N}_0 \\ \min\{m_1, m_2\} = m}} C_v^{(m_1)} \otimes C_v^{(m_2) t}$$

of (III.1) for which  $\frac{1}{M^m}$  is small compared to the transfer momentum. Sector counting is made precise in Remark III.12.ii and Lemma C.2. The basic propagator estimates are stated in Appendix A and are adapted to the present situation in Lemma III.14. Lemma III.11 shows how one can combine sector counting and propagator estimates on quantities like  $g \bullet \mathcal{C}^{(m)} \bullet h$ . The resulting estimates turn out to be summable over those  $m$ 's for which  $\frac{1}{M^m}$  is smaller than the transfer momentum. This is used to prove parts (b) and (c) of Theorem II.19 (which are reformulated as Theorem III.9) and to reduce the statement of part (a) of Theorem II.19 to the situation of transfer momentum smaller than  $\mathfrak{l}_j$ .

The situation of small transfer momentum is treated in Theorem III.15. To estimate  $g \bullet \mathcal{C}^{[i,j]} \bullet h$  when the transfer momentum is small compared to  $\mathfrak{l}_j$ , we replace  $\mathcal{C}^{[i,j]}$  with a model bubble propagator  $\mathcal{M}$  with a factorized cutoff similar to that of Lemma I.1. In Proposition III.27, we use a position space bound on  $\mathcal{M}$  (which is proven in Appendix B) to estimate  $g \bullet \mathcal{M} \bullet h$ . Propositions III.19, III.22 and III.24 use sector counting and simple propagator estimates as above to bound  $g \bullet (\mathcal{C}^{[i,j]} - \mathcal{M}) \bullet h$ .

The results Lemma III.28 through Corollary III.31, of the final two subsections, are used in the proof, in Lemma II.29, of the existence and continuity properties of the infrared limit of ladders.

Before we implement the program outlined above, we introduce some notation, prove some utility Lemmata and reformulate Theorem II.19 in terms of the new notation.

Let

$$\mathfrak{Y} = \mathbb{M} \sqcup (\mathbb{R} \times \mathbb{R}^2)$$

be the disjoint union of the set,  $\mathbb{M}$ , of all possible momenta and the set,  $\mathbb{R} \times \mathbb{R}^2$ , of all possible positions. We consider  $\mathfrak{Y}$  as the special case of the space  $\mathfrak{Y}_\Sigma$  of the introduction, with the set of sectors  $\Sigma = \Sigma_0$  where  $\Sigma_0$  contains only a single element, namely all momentum space,  $\mathbb{M}$ . In particular, as in (I.2),  $\mathfrak{Y}^4$  is the disjoint union

$$\mathfrak{Y}^4 = \bigcup_{i_1, i_2, i_3, i_4 \in \{0,1\}} \mathfrak{Y}_{i_1} \times \mathfrak{Y}_{i_2} \times \mathfrak{Y}_{i_3} \times \mathfrak{Y}_{i_4}$$

where  $\mathfrak{Y}_0 = \mathbb{M}$  and  $\mathfrak{Y}_1 = \mathbb{R} \times \mathbb{R}^2$ . For a translation invariant function  $f$  on  $\mathfrak{Y}^4$ , we define

$$|||f||| = |f|_{\Sigma_0, \Sigma_0}^{(0,0,0)}$$

using the norm  $|\cdot|_{\Sigma, \Sigma'}^{(0,0,0)}$  of Definition I.14. Concretely,

$$|||f||| = \sum_{i_1, i_2, i_3, i_4 \in \{0,1\}} \sup_{\substack{k_\nu \in \mathbb{M} \\ \nu=1,2,3,4 \\ \text{with } i_\nu=0}} |||f|_{(i_1, \dots, i_4)}|||_{1, \infty}$$

Here, the  $\nu^{\text{th}}$  argument of  $f$  is  $k_\nu$  when  $i_\nu = 0$  and  $x_\nu$  when  $i_\nu = 1$ . The  $|||\cdot|||_{1, \infty}$  norm of Definition I.11 is applied to all spatial arguments of  $f|_{(i_1, \dots, i_4)}$ .

**Definition III.1** We define the bubble operator norm of any translation invariant bubble propagator  $P(x_1, x_2, x_3, x_4)$  by

$$|||P|||_{\text{bubble}} = \sup_{G, H} \frac{|||G \circ P \circ H|||}{|||G||| |||H|||}$$

where the sup is over nonzero, translation invariant functions on  $\mathfrak{Y}^4$ .

**Lemma III.2** *Let  $P$  be a translation invariant bubble propagator. Then*

$$|||P|||_{\text{bubble}} \leq \min \left\{ \min_{n=1,2} \sup_{x_1, x_2} \int dy_n \sup_{y_{\bar{n}}} |P(x_1, x_2, y_1, y_2)|, \right. \\ \left. \min_{n=1,2} \sup_{y_1, y_2} \int dx_n \sup_{x_{\bar{n}}} |P(x_1, x_2, y_1, y_2)| \right\}$$

where  $\bar{n} = 2$  if  $n = 1$  and  $\bar{n} = 1$  if  $n = 2$ .

**Proof:** Let  $c_P$  be the right hand side of the claim. We must prove that

$$|||G \circ P \circ H||| \leq c_P |||G||| |||H|||$$

for all translation invariant functions,  $G, H$  on  $\mathfrak{Y}^4$ . It suffices to consider  $G$  and  $H$  obeying

$$G = G|_{(i_1, i_2, 1, 1)} \quad H = H|_{(1, 1, i_3, i_4)}$$

for some  $i_1, i_2, i_3, i_4 \in \{0, 1\}$ .

First consider the case  $i_1 = i_2 = i_3 = i_4 = 0$ . By definition

$$\begin{aligned} & |G \circ P \circ H(k_1, k_2, k_3, k_4)| \\ & \leq \int d^3 x_2 d^3 y_1 d^3 y_2 |G(k_1, k_2, 0, x_2)| |P(0, x_2, y_1, y_2)| |H(y_1, y_2, k_3, k_4)| \\ & \leq |||G||| \sup_{x_2} \int dy_1 dy_2 |P(0, x_2, y_1, y_2)| |H(y_1, y_2, k_3, k_4)| \quad (\text{III.5}) \\ & \leq |||G||| \sup_{x_2} \int dy_n \left[ \sup_{y_{\bar{n}}} |P(0, x_2, y_1, y_2)| \right] \int dy_{\bar{n}} |H(y_1, y_2, k_3, k_4)| \\ & \leq |||G||| |||H||| \sup_{x_2} \int dy_n \sup_{y_{\bar{n}}} |P(0, x_2, y_1, y_2)| \end{aligned}$$

The other bound is achieved in a similar fashion, starting from

$$\begin{aligned} & |G \circ P \circ H(k_1, k_2, k_3, k_4)| \\ & \leq \int d^3 x_1 d^3 x_2 d^3 y_2 |G(k_1, k_2, x_1, x_2)| |P(x_1, x_2, 0, y_2)| |H(0, y_2, k_3, k_4)| \end{aligned}$$

Now consider the case in which at least one of  $i_1, i_2, i_3, i_4$  is one. Pick any  $\ell \in \{1, 2, 3, 4\}$  with  $i_\ell = 1$ . Then, by translation invariance,

$$\begin{aligned} & \sup_{y_\ell} \sup_{\substack{y_\nu \in \mathbb{M} \\ \nu=1,2,3,4 \\ \text{with } i_\nu=0}} \int \prod_{\substack{\nu=1,2,3,4 \\ \text{with } i_\nu=1 \\ \text{and } \nu \neq \ell}} dy_\nu |G \circ P \circ H(y_1, y_2, y_3, y_4)| \\ & \leq \sup_{y_\ell} \sup_{\substack{y_\nu \in \mathbb{M} \\ \nu=1,2,3,4 \\ \text{with } i_\nu=0}} \int \prod_{\substack{\nu=1,2,3,4 \\ \text{with } i_\nu=1 \\ \text{and } \nu \neq \ell}} dy_\nu \prod_{\nu=1,2,3,4} dx_\nu |G(y_1, y_2, x_1, x_2) P(x_1, x_2, x_3, x_4) H(x_3, x_4, y_3, y_4)| \\ & = \sup_{\substack{y_\nu \in \mathbb{M} \\ \nu=1,2,3,4 \\ \text{with } i_\nu=0}} \int \prod_{\substack{\nu=1,2,3,4 \\ \text{with } i_\nu=1 \\ \text{and } \nu \neq \ell}} dy_\nu \prod_{\nu=1,2,3,4} dx_\nu |G(y_1, y_2, x_1, x_2) P(x_1, x_2, x_3, x_4) H(x_3, x_4, y_3, y_4)|_{y_\ell=0} \\ & = \sup_{\substack{y_\nu \in \mathbb{M} \\ \nu=1,2,3,4 \\ \text{with } i_\nu=0}} \int \prod_{\substack{\nu=1,2,3,4 \\ \text{with } i_\nu=1 \\ \text{and } \nu \neq \ell}} dy_\nu \prod_{\nu=1,2,3,4} dx_\nu |G(y_1, y_2, 0, x_2) P(0, x_2, x_3, x_4) H(x_3, x_4, y_3, y_4)|_{y_\ell=-x_1} \\ & = \sup_{\substack{y_\nu \in \mathbb{M} \\ \nu=1,2,3,4 \\ \text{with } i_\nu=0}} \int \prod_{\substack{\nu=1,2,3,4 \\ \text{with } i_\nu=1}} dy_\nu \prod_{\nu=2,3,4} dx_\nu |G(y_1, y_2, 0, x_2) P(0, x_2, x_3, x_4) H(x_3, x_4, y_3, y_4)| \\ & \leq |||G||| \sup_{\substack{y_\nu \in \mathbb{M} \\ \nu=3,4 \\ \text{with } i_\nu=0}} \sup_{x_2} \int \prod_{\substack{\nu=3,4 \\ \text{with } i_\nu=1}} dy_\nu \prod_{\nu=3,4} dx_\nu |P(0, x_2, x_3, x_4) H(x_3, x_4, y_3, y_4)| \quad (\text{III.6}) \end{aligned}$$

$$\leq \|G\| \sup_{\substack{y_\nu \in \mathbb{M} \\ \nu=3,4 \\ \text{with } i_\nu=0}} \sup_{x_2} \int dx_3 dx_4 |P(0, x_2, x_3, x_4)| \int \prod_{\substack{\nu=3,4 \\ \text{with } i_\nu=1}} dy_\nu |H(x_3, x_4, y_3, y_4)|$$

For the second equality, we made the change of variables  $y_\nu \rightarrow y_\nu + x_1$ , for each  $\nu \neq \ell$  with  $i_\nu = 1$  and the change of variables  $x_\nu \rightarrow x_\nu + x_1$ , for each  $\nu = 2, 3, 4$  and then used translation invariance of the three kernels. This replaces “ $y_\ell = 0$ ” by “ $y_\ell = -x_1$ ”. For the third equality, we made the change of variables  $x_1 \rightarrow -y_\ell$ . Now we may continue as in the case  $i_1 = i_2 = i_3 = i_4 = 0$ . ■

Our bubble propagators are typically of the form  $P = A \otimes B^t$  with translation invariant propagators  $A$  and  $B$ . If  $A$  is a translation invariant propagator, we write  $A(y - x)$  in place of  $A(x, y)$ . With this convention the  $L^1$ – $L^\infty$  norm of Definition I.11 reduces to the  $L^1$  norm  $\|A\|_{L^1} = \int |A(y)| d^3y$ . If  $P = A \otimes B^t$ , then


$$P(x_1, x_2, y_1, y_2) = A(y_1 - x_1)B(x_2 - y_2) = \begin{array}{ccc} x_1 & \xrightarrow{\quad} & y_1 \\ x_2 & \xleftarrow{\quad} & y_2 \end{array}$$

and, by Lemma III.2,

$$\|P\|_{\text{bubble}} \leq \min \{ \|A\|_{L^\infty} \|B\|_{L^1}, \|A\|_{L^1} \|B\|_{L^\infty} \} \quad (\text{III.7})$$

Given any function  $W(p, k)$  on  $\mathbb{M}^2$ , we associate to it the particle–hole bubble propagator

$$\begin{aligned} W(x_1, x_2, y_1, y_2) &= \int \frac{d^3p}{(2\pi)^3} \frac{d^3k}{(2\pi)^3} W(p, k) e^{i\langle p, x_1 - y_1 \rangle -} e^{i\langle k, y_2 - x_2 \rangle -} \\ &= \int \frac{d^3t}{(2\pi)^3} \frac{d^3k}{(2\pi)^3} W(k + t, k) e^{i\langle k, x_1 - y_1 + y_2 - x_2 \rangle -} e^{i\langle t, x_1 - y_1 \rangle -} \end{aligned} \quad (\text{III.8})$$



Here  $k$  is the loop momentum and  $t = p - k$  is the transfer momentum.

Motivated by the introduction to this section, we often treat small and large transfer momenta differently. To isolate a specific set of transfer momenta, we use a function  $R(t)$  on  $\mathbb{M}$  that is supported there.

**Definition III.3** For any function  $W(x_1, x_2, y_1, y_2)$  and any function  $R(t)$ , with Fourier transform  $\hat{R}(z)$ , we set

$$W_R(x_1, x_2, y_1, y_2) = \int dz W(x_1, x_2, y_1 - z, y_2 - z) \hat{R}(z)$$

If  $W(x_1, x_2, y_1, y_2)$  is associated with  $W(p, k)$  as in (III.8), then  $W_R(x_1, x_2, y_1, y_2)$  is associated with

$$W_R(p, k) = W(p, k)R(p - k)$$

**Lemma III.4** Let  $A$  and  $B$  be translation invariant propagators and  $R(t)$  a function on  $\mathbb{M}$ . Then

$$\|(A \otimes B^t)_R\|_{\text{bubble}} \leq \|\hat{R}(x)\|_{L^1} \min \{ \|A(x)\|_{L^\infty} \|B(x)\|_{L^1}, \|A(x)\|_{L^1} \|B(x)\|_{L^\infty} \}$$

**Proof:** By Definition III.3,

$$(A \otimes B^t)_R(x_1, x_2, y_1, y_2) = \int dz A(y_1 - x_1 - z) B(x_2 - y_2 + z) \hat{R}(z)$$

By Lemma III.2,

$$\|(A \otimes B^t)_R\|_{\text{bubble}} \leq \min_{n=1,2} \sup_{x_1, x_2} \int dy_n \sup_{y_{\bar{n}}} |(A \otimes B^t)_R(x_1, x_2, y_1, y_2)|$$

We treat  $n = 1$ . The other case is similar.

$$\begin{aligned} \sup_{x_1, x_2} \int dy_1 \sup_{y_2} |(A \otimes B^t)_R(x_1, x_2, y_1, y_2)| &\leq \int dy_1 \sup_{y_2} \int dz |A(y_1 - z) B(-y_2 - z) \hat{R}(z)| \\ &\leq \|B(x)\|_{L^\infty} \int dy_1 \int dz |A(y_1 - z) \hat{R}(z)| \\ &= \|B(x)\|_{L^\infty} \|A(x)\|_{L^1} \|\hat{R}(z)\|_{L^1} \end{aligned}$$

■

**Remark III.5** Define, for any function  $\hat{R}(x)$ , the bubble operator

$$O_R(x_1, x_2, y_1, y_2) = \hat{R}(y_1 - x_1) \delta(x_2 - y_2 + y_1 - x_1)$$

Then, for any bubble propagator  $W$ ,

$$W \circ O_R = W_R$$

Replacing  $g$  by  $\frac{1}{M^{\ell|\delta_1|}} D_{1,2}^{\delta_1} g$  and  $h$  by  $\frac{1}{M^{j|\delta_r|}} D_{3,4}^{\delta_r} h$  in Theorem II.19 reduces consideration of the norm  $|g \bullet \mathcal{C}^{[i,j]} \bullet h|_{\ell,j}^{[\delta_1,0,\delta_r]}$  to a  $|\cdot|_{\ell,j}^{[0,0,0]}$  norm. Therefore, we introduce the short hand notation

**Definition III.6** For  $f$  a function on  $\mathfrak{Y}_{i_1, i_r}$ , set

$$|f|_{i_1, i_r} = |f|_{i_1, i_r}^{[0,0,0]}$$

With the reduction to  $\delta_1 = \delta_r = 0$ , indicated above, Theorem II.19 becomes bounds on the  $|\cdot|_{\ell, j}$  norm of quantities like  $g \bullet \mathcal{C}^{[i, j]} \bullet h$ . For the rest of this section, we fix  $\ell \geq 1$  and consider, more generally,  $|\cdot|_{\ell, r}$  norms with  $r \geq j$ . The  $|\cdot|_{\ell, r}$  norm of a function  $f$  is obtained by fixing all arguments that lie in  $\mathbb{M}$  and the sectors of all arguments that lie in  $(\mathbb{R} \times \mathbb{R}^2) \times \Sigma_\ell$  or  $(\mathbb{R} \times \mathbb{R}^2) \times \Sigma_r$  and taking the  $\|\cdot\|_{1, \infty}$  of the result. The transfer momentum  $t$  is determined by the momenta and sectors of the last two arguments of  $f$ . This motivates the following

**Definition III.7**

i) Let  $\mathfrak{K}_r = \mathbb{M} \cup \Sigma_r$  be the disjoint union of the set  $\mathbb{M}$  of external momenta and the set  $\Sigma_r$  of sectors of scale  $r$ .

ii) Let  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$ . The subset  $\kappa_1 - \kappa_2$  of  $\mathbb{M}$  is defined by

$$\kappa_1 - \kappa_2 = \begin{cases} \{\kappa_1 - \kappa_2\} & \text{if } \kappa_1, \kappa_2 \in \mathbb{M} \\ \{\kappa_1 - k_2 \mid k_2 \in \kappa_2\} & \text{if } \kappa_1 \in \mathbb{M}, \kappa_2 \in \Sigma_r \\ \{k_1 - \kappa_2 \mid k_1 \in \kappa_1\} & \text{if } \kappa_1 \in \Sigma_r, \kappa_2 \in \mathbb{M} \\ \{k_1 - k_2 \mid k_1 \in \kappa_1, k_2 \in \kappa_2\} & \text{if } \kappa_1, \kappa_2 \in \Sigma_r \end{cases}$$

ii) Let  $f$  be a function on  $\mathfrak{Y}_{\ell, r}$  and  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$ . Then

$$\|f\|_{\kappa_1, \kappa_2} = \begin{cases} \sum_{i_1, i_2 \in \{0, 1\}} \max_{\substack{s_\nu \in \Sigma_\ell \\ \text{if } i_\nu = 1}} \sup_{\substack{k_\nu \in \mathbb{M} \\ \text{if } i_\nu = 0}} \|f|_{(i_1, i_2, 0, 0)}(y_1, y_2, \kappa_1, \kappa_2)\|_{1, \infty} & \text{if } \kappa_1, \kappa_2 \in \mathbb{M} \\ \sum_{i_1, i_2 \in \{0, 1\}} \max_{\substack{s_\nu \in \Sigma_\ell \\ \text{if } i_\nu = 1}} \sup_{\substack{k_\nu \in \mathbb{M} \\ \text{if } i_\nu = 0}} \|f|_{(i_1, i_2, 0, 1)}(y_1, y_2, \kappa_1, (x_4, \kappa_2))\|_{1, \infty} & \text{if } \kappa_1 \in \mathbb{M}, \kappa_2 \in \Sigma_r \\ \sum_{i_1, i_2 \in \{0, 1\}} \max_{\substack{s_\nu \in \Sigma_\ell \\ \text{if } i_\nu = 1}} \sup_{\substack{k_\nu \in \mathbb{M} \\ \text{if } i_\nu = 0}} \|f|_{(i_1, i_2, 1, 0)}(y_1, y_2, (x_3, \kappa_1), \kappa_2)\|_{1, \infty} & \text{if } \kappa_1 \in \Sigma_r, \kappa_2 \in \mathbb{M} \\ \sum_{i_1, i_2 \in \{0, 1\}} \max_{\substack{s_\nu \in \Sigma_\ell \\ \text{if } i_\nu = 1}} \sup_{\substack{k_\nu \in \mathbb{M} \\ \text{if } i_\nu = 0}} \|f|_{(i_1, i_2, 1, 1)}(y_1, y_2, (x_3, \kappa_1), (x_4, \kappa_2))\|_{1, \infty} & \text{if } \kappa_1, \kappa_2 \in \Sigma_r \end{cases}$$

Here, we use the decomposition of (I.2) and, for  $\nu = 1, 2$ ,  $y_\nu = k_\nu$  if  $i_\nu = 0$  and  $y_\nu = (x_\nu, s_\nu)$  if  $i_\nu = 1$ .

**Remark III.8** For a function  $f$  on  $\mathfrak{Y}_{\ell,r}$

$$|f|_{\ell,r} \leq 4 \left\{ \sup_{k_1, k_2 \in \mathbb{M}} \|f\|_{k_1, k_2} + \sup_{\substack{k_1 \in \mathbb{M} \\ \sigma_2 \in \Sigma_r}} \|f\|_{k_1, \sigma_2} + \sup_{\substack{\sigma_1 \in \Sigma_r \\ k_2 \in \mathbb{M}}} \|f\|_{\sigma_1, k_2} + \sup_{\sigma_1, \sigma_2 \in \mathbb{M}} \|f\|_{\sigma_1, \sigma_2} \right\}$$

We now state the reformulation of parts (b) and (c) of Theorem II.19. Recall the decomposition

$$\mathcal{C}^{[i,j]} = \mathcal{C}_{\text{top}}^{[i,j]} + \mathcal{C}_{\text{mid}}^{[i,j]} + \mathcal{C}_{\text{bot}}^{[i,j]}$$

of the particle-hole bubble propagator  $\mathcal{C}^{[i,j]}$  with

$$\mathcal{C}_{\text{top}}^{[i,j]} = \sum_{\substack{i \leq i_t \leq j \\ i_b > j}} C_v^{(i_t)} \otimes C_v^{(i_b)^\dagger}, \quad \mathcal{C}_{\text{mid}}^{[i,j]} = \sum_{\substack{i \leq i_t \leq j \\ i \leq i_b \leq j}} C_v^{(i_t)} \otimes C_v^{(i_b)^\dagger}, \quad \mathcal{C}_{\text{bot}}^{[i,j]} = \sum_{\substack{i_t > j \\ i \leq i_b \leq j}} C_v^{(i_t)} \otimes C_v^{(i_b)^\dagger}$$

and recall from (II.1) that

$$\Delta = \left\{ \delta \in \mathbb{N}_0 \times \mathbb{N}_0^2 \mid \delta_0 \leq r_0, \delta_1 + \delta_2 \leq r_e \right\}$$

where  $r_e + 3$  is the degree of differentiability of the dispersion relation  $e(\mathbf{k})$  and  $r_0$  is the number of  $k_0$  derivatives that we wish to control.

**Theorem III.9** *Let  $1 \leq i \leq j \leq r$  and  $\ell \geq 1$  and let  $g$  and  $h$  be sectorized, translation invariant functions on  $\mathfrak{Y}_{\ell,i}$  and  $\mathfrak{Y}_{i,r}$  respectively. Let  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$ .*

*i) For any  $\beta \in \Delta$*

$$\begin{aligned} \frac{1}{M^{|\beta|j}} \|g \bullet D_{1,3}^\beta \mathcal{C}_{\text{top}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} &\leq \text{const } |g|_{\ell,i} |h|_{i,r} \\ \frac{1}{M^{|\beta|j}} \|g \bullet D_{2,4}^\beta \mathcal{C}_{\text{bot}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} &\leq \text{const } |g|_{\ell,i} |h|_{i,r} \end{aligned}$$

*ii)*

$$\|g \bullet \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |j - i + 1| |g|_{\ell,i} |h|_{i,r}$$

*and for any  $\beta \in \Delta$  with  $|\beta| \geq 1$  and  $(\mu, \mu') = (1, 3), (2, 4)$*

$$\frac{1}{M^{|\beta|j}} \|g \bullet D_{\mu, \mu'}^\beta \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |g|_{\ell,i} |h|_{i,r}$$

*The constant const depends on  $e(\mathbf{k})$ ,  $M$  and  $\Delta$ , but not on  $i, \ell, j, r, g, h, \kappa_1$  or  $\kappa_2$ .*

The proof of Theorem III.9 follows Lemma III.14.

**Proof of Theorem II.19b,c** (assuming Theorem III.9):

As pointed out above, we may assume without loss of generality that  $\delta_l = \delta_r = 0$ . Then parts (b) and (c) of Theorem II.19 follow directly from Remark III.8 and parts (i) and (ii) of Theorem III.9, with  $r = j$ , respectively. ■



**Definition III.10** For any subset  $d \subset \mathbb{M}$ , let  $\mathcal{R}(d)$  be the set of all functions  $R(t)$  that are identically one on  $d$ .


**Lemma III.11** Let  $1 \leq i \leq j$  and  $\ell, r \geq 1$ . Let  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$  and  $g$  and  $h$  be sectorized, translation invariant functions on  $\mathfrak{Y}_{\ell,i}$  and  $\mathfrak{Y}_{i,r}$  respectively. Let  $W$  be a particle-hole bubble propagator whose total Fourier transform is of the form

$$\check{W}(p_1, k_1, p_2, k_2) = \sum_{m=i}^j \sum_{s_1, s_2 \in \Sigma_m} W_{s_1, s_2}^{(m)}(p_1, k_1, p_2, k_2) \quad \text{if } p_2 - k_2 \in \kappa_1 - \kappa_2$$

with  $W_{s_1, s_2}^{(m)}(p_1, k_1, p_2, k_2)$  vanishing unless  $p_1, p_2 \in s_1$  and  $k_1, k_2 \in s_2$ . Then

$$\begin{aligned} \|g \bullet W \bullet h\|_{\kappa_1, \kappa_2} &\leq 81 |g|_{\ell, i} |h|_{i, r} \sum_{m=i}^j \sum_{\substack{s_1, s_2 \in \Sigma_m \\ (s_1 - s_2) \cap (\kappa_1 - \kappa_2) \neq \emptyset}} \inf_{R \in \mathcal{R}(\kappa_1 - \kappa_2)} \|W_{s_1, s_2, R}^{(m)}\|_{\text{bubble}} \\ &\leq 81 |g|_{\ell, i} |h|_{i, r} \sum_{m=i}^j \sum_{\substack{s_1, s_2 \in \Sigma_m \\ (s_1 - s_2) \cap (\kappa_1 - \kappa_2) \neq \emptyset}} \|W_{s_1, s_2}^{(m)}\|_{\text{bubble}} \end{aligned}$$

**Proof:** Consider the case in which all of the external arguments of  $g \bullet W \bullet h$  are (position, sector)'s. Fix (external) sectors  $\sigma'_1, \sigma'_2 \in \Sigma_\ell$  and call  $\sigma_1 = \kappa_1, \sigma_2 = \kappa_2 \in \Sigma_r$  and  $d = \sigma_1 - \sigma_2$ . With the sector names



we have

$$g \bullet W \bullet h = \sum_{m=i}^j \sum_{\substack{u_1, v_1 \in \Sigma_i \\ u_2, v_2 \in \Sigma_i \\ s_1, s_2 \in \Sigma_m}} g((\cdot, \sigma'_1), (\cdot, \sigma'_2), (\cdot, u_1), (\cdot, u_2)) \circ W_{s_1, s_2}^{(m)} \circ h((\cdot, v_1), (\cdot, v_2), (\cdot, \sigma_1), (\cdot, \sigma_2))$$

For each choice of  $u_1, v_1, u_2, v_2, s_1, s_2$ , by conservation of momentum at the vertex  $h$ ,

$$\begin{aligned} &g((\cdot, \sigma'_1), (\cdot, \sigma'_2), (\cdot, u_1), (\cdot, u_2)) \circ W_{s_1, s_2}^{(m)} \circ h((\cdot, v_1), (\cdot, v_2), (\cdot, \sigma_1), (\cdot, \sigma_2)) \\ &= g((\cdot, \sigma'_1), (\cdot, \sigma'_2), (\cdot, u_1), (\cdot, u_2)) \circ W_{s_1, s_2, R}^{(m)} \circ h((\cdot, v_1), (\cdot, v_2), (\cdot, \sigma_1), (\cdot, \sigma_2)) \end{aligned}$$

for all  $R \in \mathcal{R}(d)$  and the convolution vanishes identically unless  $(s_1 - s_2) \cap d \neq \emptyset$ . The convolution also vanishes identically unless

$$\begin{aligned} u_1 \cap s_1 &\neq \emptyset & s_1 \cap v_1 &\neq \emptyset \\ u_2 \cap s_2 &\neq \emptyset & s_2 \cap v_2 &\neq \emptyset \end{aligned}$$

For each fixed  $s_1, s_2$  there are only 81 quadruples  $(u_1, u_2, v_1, v_2)$  satisfying these conditions. The same is true, by a similar argument, if one or more of the external arguments of  $g \bullet W \bullet h$  are momenta. Just replace, for example,  $\sigma'_1$  by  $\{k'\}$ . Hence

$$\|g \bullet W \bullet h\|_{\kappa_1, \kappa_2} \leq 81 |g|_{\ell, i} |h|_{i, r} \sum_{m=i}^j \sum_{\substack{s_1, s_2 \in \Sigma_m \\ (s_1 - s_2) \cap d \neq \emptyset}} \inf_{R \in \mathcal{R}(d)} \|W_{s_1, s_2, R}^{(m)}\|_{\text{bubble}}$$

The second inequality follows by choosing an  $R(t)$  that is identically one on a large enough ball. ■

**Remark III.12** Let  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$ .

i) The set  $\kappa_1 - \kappa_2$  is contained in a ball of radius  $2\mathfrak{l}_r$ .

ii) Let  $m \leq r$ . Then,

$$\#\{ (s_1, s_2) \in \Sigma_m \times \Sigma_m \mid (s_1 - s_2) \cap (\kappa_1 - \kappa_2) \neq \emptyset \} \leq \frac{\text{const}}{\mathfrak{l}_m}$$

iii) The set  $\{ t_0 \in \mathbb{R} \mid (t_0, \mathbf{t}) \in \kappa_1 - \kappa_2 \text{ for some } \mathbf{t} \in \mathbb{R}^2 \}$  is contained in an interval of length  $\frac{4\sqrt{2M}}{M^r}$ .

**Proof:** Part (i) is an immediate consequence of the facts that  $\kappa_1$  and  $\kappa_2$  are each contained in a ball of radius  $\mathfrak{l}_r$ . Given any fixed  $s_1 \in \Sigma_m$ ,  $(s_1 - s_2) \cap (\kappa_1 - \kappa_2) \neq \emptyset$  only if  $s_2$  intersects  $s_1 - \kappa_1 + \kappa_2$ . As  $s_1 - \kappa_1 + \kappa_2$  is contained in a ball of radius at most  $3\mathfrak{l}_m$ , there are at most eight sectors  $s_2 \in \Sigma_m$  that intersect it. This proves part (ii). Part (iii) follows from the fact that, for  $\nu = 1, 2$ ,  $\{ k_0 \in \mathbb{R} \mid (k_0, \mathbf{k}) \in \kappa_\nu \text{ for some } \mathbf{k} \in \mathbb{R}^2 \}$  is contained in an interval of length  $\frac{2\sqrt{2M}}{M^r}$ . ■

**Remark III.13** Let  $\pi : k = (k_0, \mathbf{k}) \mapsto \mathbf{k}$  be the projection of  $\mathbb{M} = \mathbb{R} \times \mathbb{R}^2$  onto its second factor. If we retain all of the hypotheses of Lemma III.11, except that we only require  $\check{W}_{s_1, s_2}^{(m)}(p_1, k_1, p_2, k_2)$  to vanish unless  $\pi(p_1), \pi(p_2) \in \pi(s_1)$  and  $\pi(k_1), \pi(k_2) \in \pi(s_2)$ , then we still have

$$\begin{aligned} \|g \bullet W \bullet h\|_{\kappa_1, \kappa_2} &\leq 81 |g|_{\ell, i} |h|_{i, r} \sum_{m=i}^j \sum_{\substack{s_1, s_2 \in \Sigma_m \\ \pi(s_1 - s_2) \cap \pi(\kappa_1 - \kappa_2) \neq \emptyset}} \inf_{R \in \mathcal{R}(\kappa_1 - \kappa_2)} \|W_{s_1, s_2, R}^{(m)}\|_{\text{bubble}} \\ &\leq 81 |g|_{\ell, i} |h|_{i, r} \sum_{m=i}^j \sum_{\substack{s_1, s_2 \in \Sigma_m \\ \pi(s_1 - s_2) \cap \pi(\kappa_1 - \kappa_2) \neq \emptyset}} \|W_{s_1, s_2}^{(m)}\|_{\text{bubble}} \end{aligned}$$

We are particularly interested in the particle–hole bubble propagator

$$\mathcal{C}^{[i,j]}(p, k) = \mathcal{C}_{\text{top}}^{[i,j]}(p, k) + \mathcal{C}_{\text{mid}}^{[i,j]}(p, k) + \mathcal{C}_{\text{bot}}^{[i,j]}(p, k)$$

where

$$\begin{aligned}\mathcal{C}_{\text{top}}^{[i,j]}(p, k) &= \sum_{\substack{i \leq m_1 \leq j \\ m_2 > j}} C_v^{(m_1)}(p) C_v^{(m_2)}(k) \\ \mathcal{C}_{\text{mid}}^{[i,j]}(p, k) &= \sum_{\substack{i \leq m_1 \leq j \\ i \leq m_2 \leq j}} C_v^{(m_1)}(p) C_v^{(m_2)}(k) \\ \mathcal{C}_{\text{bot}}^{[i,j]}(p, k) &= \sum_{\substack{m_1 > j \\ i \leq m_2 \leq j}} C_v^{(m_1)}(p) C_v^{(m_2)}(k)\end{aligned}$$

We split  $\mathcal{C}_{\text{top}}^{[i,j]}$ ,  $\mathcal{C}_{\text{mid}}^{[i,j]}$  and  $\mathcal{C}_{\text{bot}}^{[i,j]}$  into scales and we split each scale contribution into pieces with additional sector restrictions on the momenta  $p$  and  $k$  and the transfer momentum  $p - k$ . Recall, from just before Definition I.17, that  $\sum_{s \in \Sigma_m} \chi_s(k)$  is a partition of unity of the  $m^{\text{th}}$  neighbourhood subordinate to  $\Sigma_m$ . For any scale  $i \leq m \leq j$  and sectors  $s_1, s_2 \in \Sigma_m$ , set

$$\begin{aligned}\mathcal{C}_{\text{top},j,s_1,s_2}^{(m)}(p, k) &= \sum_{m_2 > j} C_v^{(m)}(p) \chi_{s_1}(p) C_v^{(m_2)}(k) \chi_{s_2}(k) \\ \mathcal{C}_{\text{mid},j,s_1,s_2}^{(m)}(p, k) &= \sum_{\substack{m_1, m_2 \leq j \\ \min(m_1, m_2) = m}} C_v^{(m_1)}(p) \chi_{s_1}(p) C_v^{(m_2)}(k) \chi_{s_2}(k) \\ \mathcal{C}_{\text{bot},j,s_1,s_2}^{(m)}(p, k) &= \sum_{m_1 > j} C_v^{(m_1)}(p) \chi_{s_1}(p) C_v^{(m)}(k) \chi_{s_2}(k)\end{aligned}$$

Then, for each of  $\text{loc} = \text{top}, \text{mid}, \text{bot}$

$$\mathcal{C}_{\text{loc}}^{[i,j]} = \sum_{m=i}^j \sum_{s_1, s_2 \in \Sigma_m} \mathcal{C}_{\text{loc},j,s_1,s_2}^{(m)}$$

**Lemma III.14** *Let  $1 \leq m \leq j$  and  $s_1, s_2 \in \Sigma_m$ . If  $\beta \in \Delta$  and  $(\mu, \mu') \in \{(1, 3), (2, 4)\}$ , then*

$$\begin{aligned}\|D_{1,3}^\beta \mathcal{C}_{\text{top},j,s_1,s_2}^{(m)}\|_{\text{bubble}} &\leq \text{const } \mathfrak{l}_m \frac{M^m}{M^j} M^{|\beta|m} \\ \|D_{\mu,\mu'}^\beta \mathcal{C}_{\text{mid},j,s_1,s_2}^{(m)}\|_{\text{bubble}} &\leq \text{const } \mathfrak{l}_m \begin{cases} M^m M^{(|\beta|-1)j} & \text{if } |\beta| \geq 2 \\ M^m (j - m + 1) & \text{if } |\beta| = 1 \\ 1 & \text{if } |\beta| = 0 \end{cases} \\ \|D_{2,4}^\beta \mathcal{C}_{\text{bot},j,s_1,s_2}^{(m)}\|_{\text{bubble}} &\leq \text{const } \mathfrak{l}_m \frac{M^m}{M^j} M^{|\beta|m}\end{aligned}$$

**Proof:** Set, for  $s \in \Sigma_m$  and  $n \geq 1$ ,

$$c_s^{(n)}(k) = C_v^{(n)}(k)\chi_s(k) \quad (\text{III.9})$$

and denote by  $c_s^{(n)}(x)$  its Fourier transform. By Lemma A.2, for all  $\beta \in \Delta$ ,

$$\|x^\beta c_s^{(m)}(x)\|_{L^1} \leq \text{const } M^{(1+|\beta|)m} \quad (\text{III.10})$$

$$\|c_s^{(n)}(x)\|_{L^\infty} \leq \text{const } \frac{\mathfrak{l}_m}{M^n} \quad (\text{III.11})$$

$$\|x^\beta c_s^{(n)}(x)\|_{L^\infty} \leq \text{const } \mathfrak{l}_m M^{(|\beta|-1)n} \quad \text{if } n \geq m \quad (\text{III.12})$$

Recall that

$$D_{1,3}^\beta \mathcal{C}_{\text{top},j,s_1,s_2}^{(m)}(x_1, x_2, y_1, y_2) = \sum_{n>j} (y_1 - x_1)^\beta c_{s_1}^{(m)}(y_1 - x_1) c_{s_2}^{(n)}(x_2 - y_2)$$

Hence, by the triangle inequality and (III.7),

$$\begin{aligned} \|D_{1,3}^\beta \mathcal{C}_{\text{top},j,s_1,s_2}^{(m)}\|_{\text{bubble}} &\leq \sum_{n>j} \|x^\beta c_{s_1}^{(m)}\|_{L^1} \|c_{s_2}^{(n)}\|_{L^\infty} \\ &\leq \sum_{n>j} \text{const } M^{(1+|\beta|)m} \frac{\mathfrak{l}_m}{M^n} \\ &\leq \text{const } \frac{M^m}{M^j} \mathfrak{l}_m M^{|\beta|m} \end{aligned}$$

The bound on  $\|D_{2,4}^\beta \mathcal{C}_{\text{bot},j,s_1,s_2}^{(m)}\|_{\text{bubble}}$  is proven similarly. As

$$\begin{aligned} D_{1,3}^\beta \mathcal{C}_{\text{mid},j,s_1,s_2}^{(m)}(x_1, x_2, y_1, y_2) &= \sum_{m \leq n \leq j} (y_1 - x_1)^\beta c_{s_1}^{(m)}(y_1 - x_1) c_{s_2}^{(n)}(x_2 - y_2) \\ &\quad + \sum_{m < n \leq j} (y_1 - x_1)^\beta c_{s_1}^{(n)}(y_1 - x_1) c_{s_2}^{(m)}(x_2 - y_2) \end{aligned}$$

we have

$$\begin{aligned} \|D_{1,3}^\beta \mathcal{C}_{\text{mid},j,s_1,s_2}^{(m)}\|_{\text{bubble}} &\leq \sum_{m \leq n \leq j} \|x^\beta c_{s_1}^{(m)}\|_{L^1} \|c_{s_2}^{(n)}\|_{L^\infty} + \sum_{m < n \leq j} \|x^\beta c_{s_1}^{(n)}\|_{L^\infty} \|c_{s_2}^{(m)}\|_{L^1} \\ &\leq \sum_{m \leq n \leq j} \text{const } M^{(1+|\beta|)m} \frac{\mathfrak{l}_m}{M^n} + \sum_{m < n \leq j} \text{const } \frac{\mathfrak{l}_m}{M^n} M^{|\beta|n} M^m \\ &\leq \text{const } \mathfrak{l}_m \sum_{m \leq n \leq j} \frac{M^m}{M^n} M^{|\beta|n} \end{aligned}$$

To bound  $\|x^\beta c_{s_1}^{(n)}\|_{L^\infty}$ , we used (III.12). ■

**Proof of Theorem III.9.i:** We prove the bound for  $\mathcal{C}_{\text{top}}^{[i,j]}$ . The proof for  $\mathcal{C}_{\text{bot}}^{[i,j]}$  is virtually identical. By Lemma III.11, Remark III.12.ii and Lemma III.14,

$$\begin{aligned}
\|g \bullet D_{1,3}^\beta \mathcal{C}_{\text{top}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} &\leq \text{const } |g|_{\ell,i} |h|_{i,r} \sum_{m=i}^j \frac{\text{const}}{\mathfrak{l}_m} \max_{s_1, s_2 \in \Sigma_m} \|D_{1,3}^\beta \mathcal{C}_{\text{top},j,s_1,s_2}^{(m)}\|_{\text{bubble}} \\
&\leq \text{const } |g|_{\ell,i} |h|_{i,r} \sum_{m=i}^j \frac{\text{const}}{\mathfrak{l}_m} \mathfrak{l}_m \frac{M^m}{M^j} M^{|\beta|m} \\
&\leq \text{const } M^{|\beta|j} |g|_{\ell,i} |h|_{i,r}
\end{aligned}$$

■

**Proof of Theorem III.9.ii:** By Lemma III.11, followed by Lemma III.14 and Remark III.12.ii, we have

$$\begin{aligned}
\|g \bullet \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} &\leq \text{const } |g|_{\ell,i} |h|_{i,r} \sum_{m=i}^j \frac{\text{const}}{\mathfrak{l}_m} \max_{s_1, s_2 \in \Sigma_m} \|\mathcal{C}_{\text{mid},s_1,s_2}^{(m)}\|_{\text{bubble}} \\
&\leq \text{const } |g|_{\ell,i} |h|_{i,r} \sum_{m=i}^j \frac{\text{const}}{\mathfrak{l}_m} \mathfrak{l}_m \\
&\leq \text{const } |j - i + 1| |g|_{\ell,i} |h|_{i,r}
\end{aligned}$$

For  $|\beta| \geq 1$  and  $(\mu, \mu') = (1, 3), (2, 4)$ , by Lemma III.14,

$$\begin{aligned}
\|g \bullet D_{\mu,\mu'}^\beta \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} &\leq \text{const } |g|_{\ell,i} |h|_{i,r} \sum_{m=i}^j \frac{\text{const}}{\mathfrak{l}_m} \max_{s_1, s_2 \in \Sigma_m} \|D_{\mu,\mu'}^\beta \mathcal{C}_{\text{mid},j,s_1,s_2}^{(m)}\|_{\text{bubble}} \\
&\leq \text{const } |g|_{\ell,i} |h|_{i,r} \sum_{m=i}^j \frac{1}{\mathfrak{l}_m} \mathfrak{l}_m M^m \begin{cases} M^{(|\beta|-1)j} & |\beta| \geq 2 \\ (j - m + 1) & |\beta| = 1 \end{cases} \\
&\leq \text{const } M^{|\beta|j} |g|_{\ell,i} |h|_{i,r}
\end{aligned}$$

since, for  $|\beta| \geq 2$ ,

$$\sum_{m=i}^j M^m M^{(|\beta|-1)j} \leq \text{const } M^{|\beta|j}$$

and, for  $|\beta| = 1$ ,

$$\sum_{m=i}^j M^m (j - m + 1) = M^j \sum_{m=i}^j M^{-(j-m)} (j - m + 1) \leq \text{const } M^j$$

■

We now start the proof of part (a) of Theorem II.19. We shall prove, at the end of this subsection, the following bound on the small transfer momentum contributions to  $g \bullet \mathcal{C}^{[i,j]} \bullet h$ .

**Theorem III.15** *Let  $1 \leq i \leq j \leq r$  and  $\ell \geq 1$  and let  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$ . Set  $d = \kappa_1 - \kappa_2$  and denote by  $\mathbf{d}$  the projection of  $d$  onto  $\{0\} \times \mathbb{R}^2$  identified with  $\mathbb{R}^2$ . By Remark III.12, the set  $\mathbf{d}$  is contained in a disc of radius  $2\mathfrak{l}_r$ . Fix such a disk and denote by  $\boldsymbol{\tau}$  its centre. Furthermore, set  $\tau_0 = \inf \{ |t_0| \mid (t_0, \mathbf{t}) \in d \text{ for some } \mathbf{t} \in \mathbb{R}^2 \}$ . Assume that*

$$\tau_0 \leq \frac{1}{M^{j-1}} \quad |\boldsymbol{\tau}| \leq \max \left\{ \frac{1}{M^j}, r^3 \mathfrak{l}_r \right\} \quad M^i \leq \mathfrak{l}_j M^j$$

*Also assume that  $p^{(i)}$  vanishes for all  $i > j + 1$ . For any sectorized, translation invariant functions  $g$  and  $h$  on  $\mathfrak{Y}_{\ell,i}$  and  $\mathfrak{Y}_{i,r}$  respectively,*

$$\|g \bullet \mathcal{C}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const} \max_{\substack{\alpha_r, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_r| + |\alpha_1| \leq 3}} |g|_{\ell,i}^{[0,0,\alpha_r]} |h|_{i,r}^{[\alpha_1,0,0]}$$

*The constant const depends on  $e(\mathbf{k})$ ,  $M$  and  $\Delta$ , but not on  $i, \ell, j, r, g, h, \kappa_1$  or  $\kappa_2$ .*

Theorem III.15 is proven following Proposition III.27.

**Proof of Theorem II.19a** (assuming Theorem III.15):

As pointed out above, we may assume without loss of generality that  $\delta_l = \delta_r = 0$ . Fix  $0 \leq i, \ell \leq j$  and sectorized, translation invariant functions  $g$  and  $h$  on  $\mathfrak{Y}_{\ell,i}$  and  $\mathfrak{Y}_{i,j}$  as in Theorem II.19. By Remark III.8, it suffices to prove that

$$\|g \bullet \mathcal{C}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const} \max_{\substack{\alpha_r, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_r| + |\alpha_1| \leq 3}} |g|_{\ell,i}^{[0,0,\alpha_r]} |h|_{i,j}^{[\alpha_1,0,0]} \quad (\text{III.13})$$

for all  $\kappa_1, \kappa_2 \in \mathfrak{K}_j$ . Fix  $\kappa_1, \kappa_2 \in \mathfrak{K}_j$ . Set  $d = \kappa_1 - \kappa_2$  and denote by  $\mathbf{d}$  the projection of  $d$  onto  $\{0\} \times \mathbb{R}^2$  identified with  $\mathbb{R}^2$ . By Remark III.12, the set  $\mathbf{d}$  is contained in a disc of radius  $2\mathfrak{l}_j$ . We fix such a disk and denote by  $\boldsymbol{\tau}$  its centre. Furthermore, we define  $\tau_0 = \inf \{ |t_0| \mid (t_0, \mathbf{t}) \in d \text{ for some } \mathbf{t} \in \mathbb{R}^2 \}$ . Define

$$\begin{aligned} j_0 &= \begin{cases} \max \{ n \in \mathbb{N}_0 \mid \tau_0 \leq \frac{1}{M^{n-1}} \} & \text{if } 0 < \tau_0 \leq M \\ 0 & \text{if } \tau_0 \geq M \\ \infty & \text{if } \tau_0 = 0 \end{cases} \\ j_1 &= \begin{cases} \max \{ n \in \mathbb{N}_0 \mid |\boldsymbol{\tau}| \leq \frac{1}{M^n} \} & \text{if } j^3 \mathfrak{l}_j < |\boldsymbol{\tau}| \leq 1 \\ 0 & \text{if } |\boldsymbol{\tau}| \geq 1 \\ \infty & \text{if } |\boldsymbol{\tau}| \leq j^3 \mathfrak{l}_j \end{cases} \\ \bar{j} &= \max \{ i - 1, \min \{ j, j_0, j_1 \} \} \end{aligned}$$

One of the tools that we use in the proof that Theorem III.15 implies Theorem II.19.a is

**Proposition III.16 (Large Transfer Momentum)**

$$\|g \bullet \mathcal{C}^{[\bar{j}+1, j]} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |g|_{\ell, i} |h|_{i, j}$$

**Proof:** If  $\min\{j, j_0, j_1\} = j$ , then  $\bar{j} = j$  and  $\mathcal{C}^{[\bar{j}+1, j]} = 0$  so that there is nothing to prove. So we may assume that  $\min\{j_0, j_1\} < j$ .

*Case 1:*  $j_0 \leq j_1$ . In this case,  $\|g \bullet \mathcal{C}^{[\bar{j}+1, j]} \bullet h\|_{\kappa_1, \kappa_2} = 0$ , because  $\mathcal{C}^{[\bar{j}+1, j]}(p, k)$  vanishes unless  $|p_0|, |k_0| \leq \frac{\sqrt{2M}}{M^{\bar{j}+1}}$  and hence unless  $|p_0 - k_0| \leq \frac{2\sqrt{2M}}{M^{\bar{j}+1}} < \frac{1}{M^{\bar{j}}} < \tau_0$ , while  $|t_0| \geq \tau_0$  for all  $t \in d$ .

*Case 2:*  $j_1 < j_0$ . In this case  $|\tau| > j^3 \mathfrak{l}_j$ . Let  $\delta_F$  be the constant of Lemma C.2. By Lemma C.2.a, with  $\epsilon = 2\mathfrak{l}_j$  and  $m \leq j$ ,

$$\#\{ (s_1, s_2) \in \Sigma_m \times \Sigma_m \mid (s_1 - s_2) \cap d \neq \emptyset \} \leq \text{const} \begin{cases} \frac{1}{\sqrt{\mathfrak{l}_m}} & \text{if } |\tau| \geq \delta_F \\ 1 + \frac{1}{|\tau| \mathfrak{l}_m} \left( \frac{1}{M^m} + \mathfrak{l}_j \right) & \text{otherwise} \end{cases}$$

Hence, by Lemma III.11 and Lemma III.14,

$$\begin{aligned} \|g \bullet \mathcal{C}^{[\bar{j}+1, j]} \bullet h\|_{\kappa_1, \kappa_2} &\leq \text{const } |g|_{\ell, i} |h|_{i, j} \sum_{m=\bar{j}+1}^j \mathfrak{l}_m \#\{ (s_1, s_2) \in \Sigma_m \times \Sigma_m \mid (s_1 - s_2) \cap d \neq \emptyset \} \\ &\leq \text{const } |g|_{\ell, i} |h|_{i, j} \begin{cases} \sum_{m=\bar{j}+1}^j \sqrt{\mathfrak{l}_m} & \text{if } |\tau| \geq \delta_F \\ 1 + \frac{1}{|\tau|} \sum_{m=j_1+1}^j \left( \frac{1}{M^m} + \mathfrak{l}_j \right) & \text{otherwise} \end{cases} \\ &\leq \text{const } |g|_{\ell, i} |h|_{i, j} \end{aligned}$$

since, by the definition of  $j_1$ ,

$$\frac{1}{|\tau|} \sum_{m=j_1+1}^j \left( \frac{1}{M^m} + \mathfrak{l}_j \right) \leq \frac{1}{|\tau|} \left( \frac{1}{M^{j_1}} + j \mathfrak{l}_j \right) \leq \text{const}$$

■

**Continuation of the proof of Theorem II.19a** (assuming Theorem III.15):

When  $M^i \geq \mathfrak{l}_{\bar{j}} M^{\bar{j}} = M^{(1-\aleph)\bar{j}}$ , we have  $|\bar{j} - i + 1| \leq \text{const } i$ . In this case Theorem III.9, with  $r = j$  and  $j = \bar{j}$ , gives

$$\|g \bullet \mathcal{C}^{[i, \bar{j}]} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } i |g|_{\ell, i} |h|_{i, j}$$

This together with Proposition III.16 yields (III.13). Therefore, we may assume that

$$M^i \leq \mathfrak{l}_{\bar{j}} M^{\bar{j}} \tag{III.14}$$

Furthermore, if  $\bar{j} = i - 1$ ,  $\mathcal{C}^{[i,\bar{j}]} = 0$  and there is nothing more to prove. So we may also assume that  $j_0, j_1 \geq i$  and  $\bar{j} \leq j, j_0, j_1$ .

Set  $v' = \sum_{i=2}^{\bar{j}+1} p^{(i)}$ . Recall that  $\mathcal{C}^{[i,\bar{j}]} = \mathcal{C}_{\text{top}}^{[i,\bar{j}]} + \mathcal{C}_{\text{mid}}^{[i,\bar{j}]} + \mathcal{C}_{\text{bot}}^{[i,\bar{j}]}$  with

$$\mathcal{C}_{\text{top}}^{[i,\bar{j}]} = \sum_{\substack{i \leq i_t \leq \bar{j} \\ i_b > \bar{j}}} C_v^{(i_t)} \otimes C_v^{(i_b) t}, \quad \mathcal{C}_{\text{mid}}^{[i,\bar{j}]} = \sum_{\substack{i \leq i_t \leq \bar{j} \\ i \leq i_b \leq \bar{j}}} C_v^{(i_t)} \otimes C_v^{(i_b) t}, \quad \mathcal{C}_{\text{bot}}^{[i,\bar{j}]} = \sum_{\substack{i_t > \bar{j} \\ i \leq i_b \leq \bar{j}}} C_v^{(i_t)} \otimes C_v^{(i_b) t}$$

and set  $\mathcal{C}'^{[i,\bar{j}]} = \mathcal{C}'_{\text{top}}^{[i,\bar{j}]} + \mathcal{C}'_{\text{mid}}^{[i,\bar{j}]} + \mathcal{C}'_{\text{bot}}^{[i,\bar{j}]}$  with

$$\mathcal{C}'_{\text{top}}^{[i,\bar{j}]} = \sum_{\substack{i \leq i_t \leq \bar{j} \\ i_b > \bar{j}}} C_{v'}^{(i_t)} \otimes C_{v'}^{(i_b) t}, \quad \mathcal{C}'_{\text{mid}}^{[i,\bar{j}]} = \sum_{\substack{i \leq i_t \leq \bar{j} \\ i \leq i_b \leq \bar{j}}} C_{v'}^{(i_t)} \otimes C_{v'}^{(i_b) t}, \quad \mathcal{C}'_{\text{bot}}^{[i,\bar{j}]} = \sum_{\substack{i_t > \bar{j} \\ i \leq i_b \leq \bar{j}}} C_{v'}^{(i_t)} \otimes C_{v'}^{(i_b) t}$$

As  $v - v'$  is supported on the  $(\bar{j} + 2)^{\text{nd}}$  extended neighbourhood,  $\mathcal{C}_{\text{mid}}^{[i,\bar{j}]} = \mathcal{C}'_{\text{mid}}^{[i,\bar{j}]}$ . Hence, by Theorem III.9.i, with  $\beta = 0$ ,  $r = j$  and  $j = \bar{j}$ ,

$$\|g \bullet [\mathcal{C}^{[i,\bar{j}]} - \mathcal{C}'^{[i,\bar{j}]}] \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |g|_{\ell, i} |h|_{i, j} \quad (\text{III.15})$$

By (III.14) and the Definitions of  $\bar{j}$  and  $\mathcal{C}'^{[i,\bar{j}]}$ , the hypotheses of Theorem III.15, with  $r = j$  and  $j = \bar{j}$ , apply to  $g \bullet \mathcal{C}'^{[i,\bar{j}]} \bullet h$ . Hence

$$\|g \bullet \mathcal{C}'^{[i,\bar{j}]} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } \max_{\substack{\alpha_r, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_r| + |\alpha_1| \leq 3}} |g|_{\ell, i}^{[0, 0, \alpha_r]} |h|_{i, j}^{[\alpha_1, 0, 0]}$$

This together with (III.15) and Proposition III.16 yields (III.13). This completes the proof that Theorem III.15 implies Theorem II.19.a.  $\blacksquare$

The rest of this subsection is devoted to the proof of Theorem III.15. So we fix  $\ell \geq 1$ ,  $1 \leq i \leq j \leq r$  and sectorized, translation invariant functions,  $g$  and  $h$ , on  $\mathfrak{Y}_{\ell, i}$  and  $\mathfrak{Y}_{i, j}$  respectively. We also fix  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$  and assume that

$$\tau_0 \leq \frac{1}{M^{j-1}} \quad |\tau| \leq \max \left\{ \frac{1}{M^j}, r^3 \mathfrak{l}_r \right\} \quad M^i \leq \mathfrak{l}_j M^j \quad (\text{III.16})$$

and that  $p^{(i)}$  vanishes for all  $i > j + 1$ .

We shall not need to decompose  $\mathcal{C}^{[i, j]} = \mathcal{C}_{\text{top}}^{[i, j]} + \mathcal{C}_{\text{mid}}^{[i, j]} + \mathcal{C}_{\text{bot}}^{[i, j]}$  but we still split  $\mathcal{C}^{[i, j]}$  into scales and split each scale contribution into pieces with additional sector restrictions. For any scale  $i \leq m \leq j$  and sectors  $s_1, s_2 \in \Sigma_m$ , set

$$\mathcal{C}_{s_1, s_2}^{(m)}(p, k) = \sum_{\substack{m_1, m_2 \geq 0 \\ \min(m_1, m_2) = m}} c_{s_1}^{(m_1)}(p) c_{s_2}^{(m_2)}(k)$$



where  $c_s^{(n)}$  was defined in (III.9). Then

$$\mathcal{C}^{[i,j]} = \sum_{m=i}^j \sum_{s_1, s_2 \in \Sigma_m} \mathcal{C}_{s_1, s_2}^{(m)}$$

By Lemmas III.4 and III.14,

$$\begin{aligned} \|\mathcal{C}_{s_1, s_2, R}^{(m)}\|_{\text{bubble}} &\leq \text{const } \mathfrak{l}_m \|\hat{R}(x)\|_{L^1} \\ \|\mathcal{C}_{s_1, s_2}^{(m)}\|_{\text{bubble}} &\leq \text{const } \mathfrak{l}_m \end{aligned} \quad (\text{III.17})$$

### *Reduction to the Model Bubble Propagator*

The above argument for large transfer momentum implicitly exploited the fact that the particle–hole bubble is Hölder continuous in the transfer momentum  $t$  when  $t$  is nonzero. As was pointed out in the introduction, this is false for  $t = 0$ . However, if one restricts to transfer momenta with  $t_0 = 0$  then, at least for the delta function interaction,  $C^\infty$  dispersion relation and a model propagator with suitable cutoff procedure, the particle–hole bubble is in fact  $C^\infty$  for  $\mathbf{t}$  near zero. This was seen in Lemma I.1.

Lemma I.1 applied to the particle–hole bubble with a delta function interaction and choice of cutoff different from that used in this paper. In the present situation, we have general interaction kernels  $g$  and  $h$  rather than delta functions and cutoffs that do not treat  $k_0$  and  $e(\mathbf{k})$  independently. Furthermore, the time component  $t_0$  of the transfer momentum need not be zero. We now perform three reduction steps leading to a situation similar to that of Lemma I.1.

#### *Step 1 (Decoupling of the $k_0$ integral.)*

Define the zero component localization operator

$$\mathcal{Z}(x_1, x_2, y_1, y_2) = \delta(x_1 - y_1) \delta(\mathbf{x}_2 - \mathbf{y}_2) \delta(y_{1,0} - y_{2,0}) \quad (\text{III.18})$$

where  $x_i = (x_{i,0}, \mathbf{x}_i)$  and  $y_i = (y_{i,0}, \mathbf{y}_i)$ . The transpose of this operator has kernel

$$\mathcal{Z}^t(x_1, x_2, y_1, y_2) = \delta(x_1 - y_1) \delta(\mathbf{x}_2 - \mathbf{y}_2) \delta(x_{1,0} - x_{2,0})$$

**Remark III.17** If  $W(x_1, x_2, y_1, y_2)$  is a particle–hole propagator

$$(\mathcal{Z} \circ W \circ \mathcal{Z}^t)(x_1, x_2, y_1, y_2) = W(x_1, (x_{1,0}, \mathbf{x}_2), y_1, (y_{1,0}, \mathbf{y}_2))$$

If  $W(x_1, x_2, y_1, y_2)$  is associated to  $W(p, k)$  as in (III.8), then

$$(\mathcal{Z} \circ W \circ \mathcal{Z}^t)(x_1, x_2, y_1, y_2) = \int \frac{d^3 t}{(2\pi)^3} \frac{d^2 \mathbf{k}}{(2\pi)^3} e^{i\mathbf{k} \cdot (\mathbf{x}_1 - \mathbf{y}_1 + \mathbf{y}_2 - \mathbf{x}_2)} e^{i\langle t, x_1 - y_1 \rangle} \int dk_0 W(k + t, k)$$

That is,  $(\mathcal{Z} \circ W \circ \mathcal{Z}^t)$  is associated to  $\delta(k_0) \int d\omega W((\omega, \mathbf{0}) + p, (\omega, \mathbf{k}))$ .

**Lemma III.18** *Let  $W$  be a particle-hole bubble propagator.*

*i) Let  $R(t)$  be any cutoff function for the transfer momentum. Then,*

$$(\mathcal{Z} \circ W \circ \mathcal{Z}^t)_R = \mathcal{Z} \circ W_R \circ \mathcal{Z}^t$$

*ii) For any translation invariant kernels  $G$  on  $\mathfrak{Y}^2 \times (\mathbb{R} \times \mathbb{R}^2)^2$  and  $H$  on  $(\mathbb{R} \times \mathbb{R}^2)^2 \times \mathfrak{Y}^2$ ,*

$$\|G \circ \mathcal{Z}\| \leq \|G\| \quad \text{and} \quad \|\mathcal{Z}^t \circ H\| \leq \|H\|$$

*iii)*

$$\|\mathcal{Z} \circ W\|_{\text{bubble}} \leq \|W\|_{\text{bubble}} \quad \text{and} \quad \|W \circ \mathcal{Z}^t\|_{\text{bubble}} \leq \|W\|_{\text{bubble}}$$

*iv)*

$$\begin{aligned} \|\mathcal{Z} \circ W \circ \mathcal{Z}^t\|_{\text{bubble}} \leq \min \Big\{ & \min_{n=1,2} \sup_{x_1, x_2} \int d\mathbf{y}_n dy_{1,0} \sup_{\mathbf{y}_{\bar{n}}} |W(x_1, (x_{1,0}, \mathbf{x}_2), y_1, (y_{1,0}, \mathbf{y}_2))|, \\ & \min_{n=1,2} \sup_{y_1, y_2} \int d\mathbf{x}_n dx_{1,0} \sup_{\mathbf{x}_{\bar{n}}} |W(x_1, (x_{1,0}, \mathbf{x}_2), y_1, (y_{1,0}, \mathbf{y}_2))| \Big\} \end{aligned}$$

where  $\bar{n} = 2$  if  $n = 1$  and  $\bar{n} = 1$  if  $n = 2$ .

**Proof:** i) This is obvious from Remark III.5, since  $\mathcal{Z}^t \circ O_R = O_R \circ \mathcal{Z}^t$ .

ii) This is obvious since

$$\begin{aligned} (G \circ \mathcal{Z})(\cdot, \cdot, x_3, x_4) &= \delta(x_{3,0} - x_{4,0}) \int d\omega G(\cdot, \cdot, x_3, (\omega, \mathbf{x}_4)) \\ (\mathcal{Z}^t \circ H)(x_1, x_2, \cdot, \cdot) &= \delta(x_{1,0} - x_{2,0}) \int d\omega H(x_1, (\omega, \mathbf{x}_2), \cdot, \cdot) \end{aligned}$$

iii) By part (ii), for any translation invariant  $G, H$

$$\|G \circ \mathcal{Z} \circ W \circ H\| \leq \|G \circ \mathcal{Z}\| \|W\|_{\text{bubble}} \|H\| \leq \|G\| \|W\|_{\text{bubble}} \|H\|$$

and similarly for  $\|G \circ W \circ \mathcal{Z}^t \circ H\|$ .

iv) The bounds with  $n = 1, \bar{n} = 2$  are direct consequences of Remark III.17 and Lemma III.2.

We prove

$$\|\mathcal{Z} \circ W \circ \mathcal{Z}^t\|_{\text{bubble}} \leq \sup_{x_1, x_2} \int d\mathbf{y}_2 dy_{1,0} \sup_{\mathbf{y}_1} |W(x_1, (x_{1,0}, \mathbf{x}_2), y_1, (y_{1,0}, \mathbf{y}_2))|$$

The remaining case is similar. Let  $G(y_1, y_2, u_1, u_2)$  and  $H(v_1, v_2, y_3, y_4)$  be translation invariant four-legged kernels obeying

$$G = G|_{(i_1, i_2, 1, 1)} \quad H = H|_{(1, 1, i_3, i_4)}$$

for some  $i_1, i_2, i_3, i_4 \in \{0, 1\}$ . By (III.5) and (III.6), with  $P$  replaced by  $\mathcal{Z} \circ W \circ \mathcal{Z}^t$ ,

$$\begin{aligned} & \|G \circ \mathcal{Z} \circ W \circ \mathcal{Z}^t \circ H\| \\ & \leq \|G\| \sup_{\substack{y_\nu \in \mathbb{M} \\ \nu=3,4 \\ \text{with } i_\nu=0}} \sup_{u_1, u_2} \int \prod_{\substack{\nu=3,4 \\ \text{with } i_\nu=1}} dy_\nu dv_1 dv_2 |W(u_1, (u_{1,0}, \mathbf{u}_2), v_1, (v_{1,0}, \mathbf{v}_2)) H(v_1, v_2, y_3, y_4)| \\ & \leq \|G\| \sup_{\substack{y_\nu \in \mathbb{M} \\ \nu=3,4 \\ \text{with } i_\nu=0}} \sup_{u_1, u_2} \int dv_{1,0} d\mathbf{v}_2 \left\{ \sup_{\mathbf{v}_1} |W(u_1, (u_{1,0}, \mathbf{u}_2), v_1, (v_{1,0}, \mathbf{v}_2))| \right. \\ & \quad \left. \int \prod_{\substack{\nu=3,4 \\ \text{with } i_\nu=1}} dy_\nu dv_{2,0} d\mathbf{v}_1 |H(v_1, v_2, y_3, y_4)| \right\} \\ & \leq \|G\| \left[ \sup_{u_1, u_2} \int dv_{1,0} d\mathbf{v}_2 \sup_{\mathbf{v}_1} |W(u_1, (u_{1,0}, \mathbf{u}_2), v_1, (v_{1,0}, \mathbf{v}_2))| \right] \\ & \quad \sup_{\substack{y_\nu \in \mathbb{M} \\ \nu=3,4 \\ \text{with } i_\nu=0}} \sup_{v_{1,0}, \mathbf{v}_2} \int \prod_{\substack{\nu=3,4 \\ \text{with } i_\nu=1}} dy_\nu dv_{2,0} d\mathbf{v}_1 |H(v_1, v_2, y_3, y_4)| \end{aligned}$$

By translation invariance

$$\begin{aligned} & \sup_{v_{1,0}, \mathbf{v}_2} \int \prod_{\substack{\nu=3,4 \\ \text{with } i_\nu=1}} dy_\nu dv_{2,0} d\mathbf{v}_1 |H(v_1, v_2, y_3, y_4)| \\ & = \sup_{v_1} \int \prod_{\substack{\nu=3,4 \\ \text{with } i_\nu=1}} dy_\nu dv_2 |H((v_{1,0}, \mathbf{v}_2), (v_{2,0}, \mathbf{v}_1), y_3, y_4)| \\ & = \sup_{v_1} \int \prod_{\substack{\nu=3,4 \\ \text{with } i_\nu=1}} dy_\nu dv_2 |H(v_1, (v_{2,0}, 2\mathbf{v}_1 - \mathbf{v}_2), y_3, y_4)| \\ & = \sup_{v_1} \int \prod_{\substack{\nu=3,4 \\ \text{with } i_\nu=1}} dy_\nu dv_2 |H(v_1, v_2, y_3, y_4)| \leq \|H\| \end{aligned}$$

■

### Proposition III.19

$$\|g \bullet (\mathcal{C}^{[i,j]} - \mathcal{Z} \bullet \mathcal{C}^{[i,j]} \bullet \mathcal{Z}^t) \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const} \max_{\substack{\alpha_1, \alpha_r \in \Delta \\ \alpha_1 + \alpha_r = (1, 0, 0)}} |g|_{\ell, i}^{[0, 0, \alpha_r]} |h|_{i, r}^{[\alpha_1, 0, 0]}$$

In preparation for the proof, which follows Lemma III.21, we define

$$\begin{aligned} g_r((x_1, \sigma'_1), (x_2, \sigma'_2), (x_3, u_1), (x_4, u_2)) &= (x_{4,0} - x_{3,0})g((x_1, \sigma'_1), (x_2, \sigma'_2), (x_3, u_1), (x_4, u_2)) \\ h_l((x_1, v_1), (x_2, v_2), (x_3, \sigma_1), (x_4, \sigma_2)) &= (x_{2,0} - x_{1,0})h((x_1, v_1), (x_2, v_2), (x_3, \sigma_1), (x_4, \sigma_2)) \end{aligned}$$

For a particle-hole bubble propagator  $W(x_1, x_2, y_1, y_2)$  set

$$\begin{aligned} (D_l W)(x_1, x_2, y_1, y_2) &= \int_0^1 d\omega \frac{\partial W}{\partial x_{2,0}}(x_1, (\omega x_{2,0} + (1-\omega)x_{1,0}, \mathbf{x}_2), y_1, y_2) \\ (D_r W)(x_1, x_2, y_1, y_2) &= \int_0^1 d\omega \frac{\partial W}{\partial y_{2,0}}(x_1, x_2, y_1, (\omega y_{2,0} + (1-\omega)y_{1,0}, \mathbf{y}_2)) \end{aligned}$$

**Lemma III.20**

$$g \bullet (W - \mathcal{Z}W\mathcal{Z}^t) \bullet h = g_r \bullet D_l W \bullet h + g \circ \mathcal{Z} \bullet D_r W \bullet h_l$$

**Proof:** By Remark III.17,

$$\begin{aligned} g \bullet (W - \mathcal{Z}W\mathcal{Z}^t) \bullet h &= g \bullet \{W(x_1, x_2, y_1, y_2) - W(x_1, (x_{1,0}, \mathbf{x}_2), y_1, (y_{1,0}, \mathbf{y}_2))\} \bullet h \\ &= g_r \bullet \left[ \frac{1}{x_{2,0} - x_{1,0}} \{W(x_1, x_2, y_1, y_2) - W(x_1, (x_{1,0}, \mathbf{x}_2), y_1, y_2)\} \right] \bullet h \\ &\quad + g \bullet \left[ \{W(x_1, (x_{1,0}, \mathbf{x}_2), y_1, y_2) - W(x_1, (x_{1,0}, \mathbf{x}_2), y_1, (y_{1,0}, \mathbf{y}_2))\} \frac{1}{y_{2,0} - y_{1,0}} \right] \bullet h_l \\ &= g_r \bullet D_l W \bullet h + g \circ \mathcal{Z} \bullet D_r W \bullet h_l \end{aligned}$$

by the Fundamental Theorem of Calculus. ■

**Lemma III.21** *Let  $i \leq m \leq j$  and  $s_1, s_2 \in \Sigma_m$ . Then*

$$\begin{aligned} \|D_l \mathcal{C}_{s_1, s_2}^{(m)}\|_{\text{bubble}} &\leq \text{const} \frac{l_m}{M^m} \\ \|D_r \mathcal{C}_{s_1, s_2}^{(m)}\|_{\text{bubble}} &\leq \text{const} \frac{l_m}{M^m} \end{aligned}$$

**Proof:** We treat  $D_l \mathcal{C}_{s_1, s_2}^{(m)}$ . The other case is similar. For each fixed  $0 \leq \omega \leq 1$

$$\begin{aligned} \left( \frac{\partial}{\partial x_{2,0}} \mathcal{C}_{s_1, s_2}^{(m)} \right) (x_1, (\omega x_{2,0} + (1-\omega)x_{1,0}, \mathbf{x}_2), y_1, y_2) \\ = \sum_{\substack{m_1, m_2 \geq 0 \\ \min\{m_1, m_2\} = m}} c_{s_1}^{(m_1)}(y_1 - x_1) \left( \frac{\partial}{\partial x_{2,0}} c_{s_2}^{(m_2)} \right) ((\omega x_{2,0} + (1-\omega)x_{1,0} - y_{2,0}, \mathbf{x}_2 - \mathbf{y}_2)) \end{aligned}$$

We bound the bubble norm of each term separately. For  $m_1 \geq m_2 = m$ , by Lemma III.2,

$$\begin{aligned}
& \left\| c_{s_1}^{(m_1)} \left( \frac{\partial}{\partial x_{2,0}} c_{s_2}^{(m_2)} \right) \right\|_{\text{bubble}} \\
& \leq \|c_{s_1}^{(m_1)}\|_{L^\infty} \sup_{x_1, x_2} \int dy_2 \left| \left( \frac{\partial}{\partial x_{2,0}} c_{s_2}^{(m_2)} \right) ((\omega x_{2,0} + (1-\omega)x_{1,0} - y_{2,0}, \mathbf{x}_2 - \mathbf{y}_2)) \right| \\
& = \|c_{s_1}^{(m_1)}\|_{L^\infty} \left\| \frac{\partial}{\partial x_{2,0}} c_{s_2}^{(m_2)} \right\|_{L^1} \\
& \leq \text{const} \frac{l_m}{M^{m_1}} \frac{1}{M^{m_2}} M^{m_2} \leq \text{const} \frac{l_m}{M^{m_1}}
\end{aligned}$$

by parts (iii) and (iv) of Lemma A.2. For  $m = m_1 \leq m_2$

$$\left\| c_{s_1}^{(m_1)} \left( \frac{\partial}{\partial x_{2,0}} c_{s_2}^{(m_2)} \right) \right\|_{\text{bubble}} \leq \left\| \frac{\partial}{\partial x_{2,0}} c_{s_2}^{(m_2)} \right\|_{L^\infty} \|c_{s_1}^{(m_1)}\|_{L^1} \leq \text{const} \frac{l_m}{M^{2m_2}} M^{m_1}$$

Hence

$$\|D_1 \mathcal{C}_{s_1, s_2}^{(m)}\|_{\text{bubble}} \leq \text{const} \sum_{n=m}^{\infty} \left[ \frac{l_m}{M^n} + \frac{l_m}{M^{2n}} M^n \right] \leq \text{const} \frac{l_m}{M^m}$$

■

**Proof of Proposition III.19:** By Lemma III.20 followed by Remark III.13<sup>(1)</sup>

$$\begin{aligned}
& \|g \bullet (\mathcal{C}^{[i,j]} - \mathcal{Z} \bullet \mathcal{C}^{[i,j]} \bullet \mathcal{Z}^t) \bullet h\|_{\kappa_1, \kappa_2} \\
& \leq \|g_r \bullet D_1 \mathcal{C}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} + \|g \circ \mathcal{Z} \bullet D_r \mathcal{C}^{[i,j]} \bullet h_l\|_{\kappa_1, \kappa_2} \\
& \leq \text{const} |g_r|_{\ell, i} |h|_{i, r} \sum_{m=i}^j \sum_{\substack{s_1, s_2 \in \Sigma_m \\ \pi(s_1 - s_2) \cap \pi(d) \neq \emptyset}} \|D_1 \mathcal{C}_{s_1, s_2}^{(m)}\|_{\text{bubble}} \\
& \quad + \text{const} |g \circ \mathcal{Z}|_{\ell, i} |h_l|_{i, r} \sum_{m=i}^j \sum_{\substack{s_1, s_2 \in \Sigma_m \\ \pi(s_1 - s_2) \cap \pi(d) \neq \emptyset}} \|D_r \mathcal{C}_{s_1, s_2}^{(m)}\|_{\text{bubble}}
\end{aligned}$$

By the spatial projection of Remark III.12.ii,

$$\#\{ (s_1, s_2) \in \Sigma_m \times \Sigma_m \mid \pi(s_1 - s_2) \cap \pi(d) \neq \emptyset \} \leq \frac{\text{const}}{l_m} \quad (\text{III.19})$$

Using this, Lemma III.21, Lemma III.18.ii and the definitions of  $g_r$ ,  $h_l$ , we have

$$\begin{aligned}
& \|g \bullet (\mathcal{C}^{[i,j]} - \mathcal{Z} \bullet \mathcal{C}^{[i,j]} \bullet \mathcal{Z}^t) \bullet h\|_{\kappa_1, \kappa_2} \\
& \leq \text{const} M^i |g|_{\ell, i}^{[0,0,(1,0,0)]} |h|_{i, r} \sum_{m=i}^j \frac{1}{l_m} \frac{l_m}{M^m} + \text{const} |g|_{\ell, i} M^i |h|_{i, r}^{[(1,0,0),0,0]} \sum_{m=i}^j \frac{1}{l_m} \frac{l_m}{M^m}
\end{aligned}$$

■

---

<sup>(1)</sup> The operators  $D_l, D_r$  can enlarge supports in the  $k_0$  direction. So we cannot apply Lemma III.11 directly.

*Step 2 (Reduction to  $t_0 = 0$ .)*

For any particle-hole bubble propagator  $W(x_1, x_2, y_1, y_2)$  set

$$\widetilde{W}(x_1, x_2, y_1, y_2) = \delta(y_{1,0} - x_{1,0}) \int dz_0 W(x_1, (x_{1,0}, \mathbf{x}_2), (z_0, \mathbf{y}_1), (z_0, \mathbf{y}_2)) \quad (\text{III.20})$$

If  $W(x_1, x_2, y_1, y_2)$  is associated to  $W(p, k)$  as in (III.8), then  $\widetilde{W}(x_1, x_2, y_1, y_2)$  is associated to

$$\widetilde{W}(p, k) = \delta(k_0) \int d\omega W((\omega, \mathbf{p}), (\omega, \mathbf{k}))$$

By Remark III.17,  $\widetilde{W} = \mathcal{Z} \circ \widetilde{W} \circ \mathcal{Z}^t$  for all particle-hole bubble propagators  $W$ .

**Proposition III.22**

$$\|g \bullet (\mathcal{Z} \bullet \mathcal{C}^{[i,j]} \bullet \mathcal{Z}^t - \widetilde{\mathcal{C}}^{[i,j]}) \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |g|_{\ell, i} |h|_{i, r}$$

**Proof:** Choose a  $C_0^\infty$  function  $\phi(t_0)$  that takes values in  $[0, 1]$ , is supported in the interval  $|t_0| \leq 2\frac{M+4\sqrt{2M}}{M^j}$ , is identically one for  $|t_0| \leq \frac{M+4\sqrt{2M}}{M^j}$  and obeys  $|\frac{d^n}{dt^n} \phi(t_0)| \leq \text{const } M^{jn}$  for  $n \leq 2$ . By Remark III.12.iii,

$$\{ |t_0| \mid (t_0, \mathbf{t}) \in d \text{ for some } \mathbf{t} \in \mathbb{R}^2 \} \subset \left[ \tau_0, \tau_0 + \frac{4\sqrt{2M}}{M^j} \right]$$

Hence, by (III.16),  $\phi$  is in  $\mathcal{R}(d)$ . By Remark III.13 and (III.19),

$$\begin{aligned} & \|g \bullet (\mathcal{Z} \circ \mathcal{C}^{[i,j]} \circ \mathcal{Z}^t - \widetilde{\mathcal{C}}^{[i,j]}) \bullet h\|_{\kappa_1, \kappa_2} \\ & \leq \text{const } |g|_{\ell, i} |h|_{i, r} \sum_{m=i}^j \frac{1}{l_m} \max_{s_1, s_2 \in \Sigma_m} \left\| \mathcal{Z} \circ \mathcal{C}_{s_1, s_2, \phi}^{(m)} \circ \mathcal{Z}^t - (\widetilde{\mathcal{C}}_{s_1, s_2}^{(m)})_\phi \right\|_{\text{bubble}} \end{aligned}$$

Here, we used  $(\mathcal{Z} \circ W \circ \mathcal{Z}^t)_R = \mathcal{Z} \circ W_R \circ \mathcal{Z}^t$ , which was proven in Lemma III.18.i. The proposition follows from the next Lemma. ■

**Lemma III.23** *Let  $m \leq j$  and  $s_1, s_2 \in \Sigma_m$ . Then*

$$\left\| \mathcal{Z} \circ \mathcal{C}_{s_1, s_2, \phi}^{(m)} \circ \mathcal{Z}^t - (\widetilde{\mathcal{C}}_{s_1, s_2}^{(m)})_\phi \right\|_{\text{bubble}} \leq \text{const } l_m \frac{M^m}{M^j}$$

**Proof:** For any  $m_1, m_2 \geq 0$ , set

$$\begin{aligned} W^{(m_1, m_2)} &= \mathcal{Z} \circ (c_{s_1}^{(m_1)} \otimes c_{s_2}^{(m_2)} t) \circ \mathcal{Z}^t - (c_{s_1}^{(m_1)} \otimes c_{s_2}^{(m_2)} t) \widetilde{\circ} \mathcal{Z}^t \\ &= \mathcal{Z} \circ (c_{s_1}^{(m_1)} \otimes c_{s_2}^{(m_2)} t) \circ \mathcal{Z}^t - \mathcal{Z} \circ (c_{s_1}^{(m_1)} \otimes c_{s_2}^{(m_2)} t) \widetilde{\circ} \mathcal{Z}^t \end{aligned}$$

Observe that

$$\mathcal{Z} \circ \mathcal{C}_{s_1, s_2, \phi}^{(m)} \circ \mathcal{Z}^t - (\widetilde{\mathcal{C}}_{s_1, s_2}^{(m)})_{\phi} = \sum_{\substack{m_1, m_2 \in \mathbb{N}_0 \\ \min\{m_1, m_2\} = m}} W_{\phi}^{(m_1, m_2)} \quad (\text{III.21})$$

We now fix any  $m_1, m_2 \geq 0$  with  $\min\{m_1, m_2\} = m$  and bound  $\|W_{\phi}^{(m_1, m_2)}\|_{\text{bubble}}$ . By definition

$$\begin{aligned} W^{(m_1, m_2)}(x_1, x_2, y_1, y_2) &= c_{s_1}^{(m_1)}(y_1 - x_1) c_{s_2}^{(m_2)}((x_{1,0} - y_{1,0}, \mathbf{x}_2 - \mathbf{y}_2)) \\ &\quad - \delta(y_{1,0} - x_{1,0}) \int du c_{s_1}^{(m_1)}((u - x_{1,0}, \mathbf{y}_1 - \mathbf{x}_1)) c_{s_2}^{(m_2)}((x_{1,0} - u, \mathbf{x}_2 - \mathbf{y}_2)) \end{aligned}$$

and

$$\begin{aligned} W_{\phi}^{(m_1, m_2)}(x_1, x_2, y_1, y_2) &= \int dz_0 \hat{\phi}(z_0) \left[ c_{s_1}^{(m_1)}((y_{1,0} - x_{1,0} - z_0, \mathbf{y}_1 - \mathbf{x}_1)) c_{s_2}^{(m_2)}((x_{1,0} - y_{1,0} + z_0, \mathbf{x}_2 - \mathbf{y}_2)) \right. \\ &\quad \left. - \delta(y_{1,0} - x_{1,0} - z_0) \int du c_{s_1}^{(m_1)}((u - x_{1,0}, \mathbf{y}_1 - \mathbf{x}_1)) c_{s_2}^{(m_2)}((x_{1,0} - u, \mathbf{x}_2 - \mathbf{y}_2)) \right] \\ &= \int dz_0 c_{s_1}^{(m_1)}((y_{1,0} - x_{1,0} - z_0, \mathbf{y}_1 - \mathbf{x}_1)) c_{s_2}^{(m_2)}((x_{1,0} - y_{1,0} + z_0, \mathbf{x}_2 - \mathbf{y}_2)) \hat{\phi}(z_0) \\ &\quad - \int du c_{s_1}^{(m_1)}((u - x_{1,0}, \mathbf{y}_1 - \mathbf{x}_1)) c_{s_2}^{(m_2)}((x_{1,0} - u, \mathbf{x}_2 - \mathbf{y}_2)) \hat{\phi}(y_{1,0} - x_{1,0}) \\ &= \int dz_0 c_{s_1}^{(m_1)}((z_0 - x_{1,0}, \mathbf{y}_1 - \mathbf{x}_1)) c_{s_2}^{(m_2)}((x_{1,0} - z_0, \mathbf{x}_2 - \mathbf{y}_2)) [\hat{\phi}(y_{1,0} - z_0) - \hat{\phi}(y_{1,0} - x_{1,0})] \end{aligned}$$

The last factor

$$\hat{\phi}(y_{1,0} - z_0) - \hat{\phi}(y_{1,0} - x_{1,0}) = (x_{1,0} - z_0) \int_0^1 dt \hat{\phi}'(y_{1,0} - x_{1,0} + t(x_{1,0} - z_0))$$

Observe that

$$\int dy_{1,0} \int_0^1 dt |\hat{\phi}'(y_{1,0} - x_{1,0} + t(x_{1,0} - z_0))| = \int dy_{1,0} |\hat{\phi}'(y_{1,0})| \leq \frac{\text{const}}{M^j}$$

since

$$|\hat{\phi}'(y_{1,0})| \leq \text{const} \frac{1/M^{2j}}{[1 + |y_{1,0}/M^j|]^2}$$

If  $m_2 \geq m_1 = m$ , we apply Lemma III.18.iv<sup>(2)</sup>, (III.11) and (III.10), giving

$$\begin{aligned}
& \|W_\phi^{(m_1, m_2)}\|_{\text{bubble}} \\
& \leq \text{const} \sup_{x_1, x_2} \int dy_1 \sup_{\mathbf{y}_2} \int dz_0 |x_{1,0} - z_0| \\
& \quad \left| c_{s_1}^{(m_1)}((z_0 - x_{1,0}, \mathbf{y}_1 - \mathbf{x}_1)) \right| \left| c_{s_2}^{(m_2)}((x_{1,0} - z_0, \mathbf{x}_2 - \mathbf{y}_2)) \right| \\
& \quad \int_0^1 dt \left| \hat{\phi}'(y_{1,0} - x_{1,0} + t(x_{1,0} - z_0)) \right| \\
& \leq \text{const} \|c_{s_2}^{(m_2)}\|_{L^\infty} \sup_{x_1} \int d\mathbf{y}_1 dz_0 |x_{1,0} - z_0| \left| c_{s_1}^{(m_1)}((z_0 - x_{1,0}, \mathbf{y}_1 - \mathbf{x}_1)) \right| \\
& \quad \int dy_{1,0} \int_0^1 dt \left| \hat{\phi}'(y_{1,0} - x_{1,0} + t(x_{1,0} - z_0)) \right| \\
& \leq \text{const} \frac{1}{M^j} \frac{\mathfrak{l}_m}{M^{m_2}} \left[ \sup_{x_1} \int d\mathbf{y}_1 dz_0 |x_{1,0} - z_0| \left| c_{s_1}^{(m_1)}((z_0 - x_{1,0}, \mathbf{y}_1 - \mathbf{x}_1)) \right| \right] \\
& = \text{const} \frac{1}{M^j} \frac{\mathfrak{l}_m}{M^{m_2}} \|x_0 c_{s_1}^{(m_1)}(x)\|_{L^1} \\
& \leq \text{const} \frac{1}{M^j} \frac{\mathfrak{l}_m}{M^{m_2}} M^{2m}
\end{aligned}$$

Similarly, if  $m_1 \geq m_2 = m$ ,

$$\begin{aligned}
& \|W_\phi^{(m_1, m_2)}\|_{\text{bubble}} \\
& \leq \text{const} \sup_{x_1, x_2} \int dy_{1,0} d\mathbf{y}_2 \sup_{\mathbf{y}_1} \int dz_0 |x_{1,0} - z_0| \\
& \quad \left| c_{s_1}^{(m_1)}((z_0 - x_{1,0}, \mathbf{y}_1 - \mathbf{x}_1)) \right| \left| c_{s_2}^{(m_2)}((x_{1,0} - z_0, \mathbf{x}_2 - \mathbf{y}_2)) \right| \\
& \quad \int_0^1 dt \left| \phi'(y_{1,0} - x_{1,0} + t(x_{1,0} - z_0)) \right| \\
& \leq \text{const} \frac{1}{M^j} \|c_{s_1}^{(m_1)}\|_{L^\infty} \sup_{x_{1,0}, \mathbf{x}_2} \int d\mathbf{y}_2 dz_0 |x_{1,0} - z_0| \left| c_{s_2}^{(m_2)}((x_{1,0} - z_0, \mathbf{x}_2 - \mathbf{y}_2)) \right| \\
& \leq \text{const} \frac{1}{M^j} \frac{\mathfrak{l}_m}{M^{m_1}} M^{2m}
\end{aligned}$$

Consequently, by (III.21),

$$\begin{aligned}
\|\mathcal{Z} \circ \mathcal{C}_{s_1, s_2, \phi}^{(m)} \circ \mathcal{Z}^t - (\tilde{\mathcal{C}}_{s_1, s_2}^{(m)})_\phi\|_{\text{bubble}} & \leq \text{const} \sum_{\substack{m_1, m_2 \geq 0 \\ \min\{m_1, m_2\} = m}} \frac{M^{2m}}{M^j} \mathfrak{l}_m \min\left\{\frac{1}{M^{m_1}}, \frac{1}{M^{m_2}}\right\} \\
& \leq \text{const} \frac{M^{2m}}{M^j} \mathfrak{l}_m \sum_{m' \geq m} \frac{1}{M^{m'}} \\
& \leq \text{const} \frac{M^m}{M^j} \mathfrak{l}_m
\end{aligned}$$

■

---

<sup>(2)</sup> Note that in the bound on the right hand side of Lemma III.18.iv,  $W$  only appears in the form  $W(x_1, (x_{1,0}, \mathbf{x}_2), y_1, (y_{1,0}, \mathbf{y}_2)) = (\mathcal{Z} \circ W \circ \mathcal{Z}^t)(x_1, x_2, y_1, y_2)$ .



Step 3 (Introduction of Factorized Cutoff.)

Define

$$\begin{aligned}\nu_0(\omega) &= \sum_{m=i+1}^{j-1} \nu(M^{2m}\omega^2) \\ \nu_1(\mathbf{p}, \mathbf{k}) &= \left[ \sum_{m_1=i+1}^{\infty} \nu(M^{2m_1}e(\mathbf{p})^2) \right] \left[ \sum_{m_2=i+1}^{\infty} \nu(M^{2m_2}e(\mathbf{k})^2) \right]\end{aligned}$$

where  $\nu$  is the single scale cutoff introduced in Definition I.2. Recall that  $\nu(x)$  is identically one on  $[\frac{2}{M}, M]$  and is supported on  $[\frac{1}{M}, 2M]$ . Define

$$e'(k) = e(\mathbf{k}) - v(k)$$

and the model particle-hole bubble propagator

$$\mathcal{M}(p, k) = \delta(k_0) \int d\omega \frac{\nu_0(\omega)\nu_1(\mathbf{p}, \mathbf{k})}{[i\omega - e'(\omega, \mathbf{p})][i\omega - e'(\omega, \mathbf{k})]} \quad (\text{III.22})$$

Observe that  $\mathcal{Z} \circ \mathcal{M} \circ \mathcal{Z}^t = \mathcal{M}$ .

**Proposition III.24**

$$\|g \bullet (\tilde{\mathcal{C}}^{[i,j]} - \mathcal{M}) \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |g|_{\ell, i} |h|_{i, r}$$

**Proof:** The cutoff  $\nu_0(\omega)\nu_1(\mathbf{p}, \mathbf{k})$  is supported on  $\left\{(\omega, \mathbf{p}, \mathbf{k}) \mid |\omega|, |e(\mathbf{p})|, |e(\mathbf{k})| \leq \sqrt{\frac{2}{M} \frac{1}{M^i}}\right\}$ . Since  $\nu^{(m)}(k)$  vanishes for all  $|ik_0 - e(\mathbf{k})| \leq \frac{1}{\sqrt{M}} \frac{1}{M^{i-1}}$  when  $m \leq i-1$

$$\nu_0(\omega)\nu_1(\mathbf{p}, \mathbf{k}) = \sum_{m_1, m_2 \geq i} \nu_0(\omega)\nu_1(\mathbf{p}, \mathbf{k})\nu^{(m_1)}((\omega, \mathbf{p}))\nu^{(m_2)}((\omega, \mathbf{k}))$$

if  $M$  is large enough. Since every  $k$  in the support of  $\nu^{(m)}(k)$  obeys  $|ik_0 - e(\mathbf{k})| \leq \sqrt{2M} \frac{1}{M^{j+1}}$  for all  $m \geq j+1$  and since  $\nu_0(\omega)$  is supported on  $|\omega| \geq \sqrt{M} \frac{1}{M^j}$

$$\nu_0(\omega)\nu_1(\mathbf{p}, \mathbf{k}) = \sum_{i \leq \min\{m_1, m_2\} \leq j} \nu_0(\omega)\nu_1(\mathbf{p}, \mathbf{k})\nu^{(m_1)}((\omega, \mathbf{p}))\nu^{(m_2)}((\omega, \mathbf{k}))$$

Recall that

$$\begin{aligned}\tilde{\mathcal{C}}^{[i,j]}(p, k) &= \delta(k_0) \int d\omega \mathcal{C}^{[i,j]}((\omega, \mathbf{p}), (\omega, \mathbf{k})) \\ &= \delta(k_0) \int d\omega \frac{\nu^{(\geq i)}((\omega, \mathbf{p}))\nu^{(\geq i)}((\omega, \mathbf{k})) - \nu^{(\geq j+1)}((\omega, \mathbf{p}))\nu^{(\geq j+1)}((\omega, \mathbf{k}))}{[i\omega - e'(\omega, \mathbf{p})][i\omega - e'(\omega, \mathbf{k})]}\end{aligned}$$

The difference of cutoff functions

$$\begin{aligned} \nu^{(\geq i)}((\omega, \mathbf{p}))\nu^{(\geq i)}((\omega, \mathbf{k})) - \nu^{(\geq j+1)}((\omega, \mathbf{p}))\nu^{(\geq j+1)}((\omega, \mathbf{k})) - \nu_0(\omega)\nu_1(\mathbf{p}, \mathbf{k}) \\ = \sum_{m=i}^j \sum_{\substack{m_1, m_2 \geq 1 \\ \min\{m_1, m_2\}=m}} \mu^{(m_1, m_2)}(\omega, \mathbf{k}, \mathbf{p}) \end{aligned}$$

where

$$\mu^{(m_1, m_2)}(\omega, \mathbf{k}, \mathbf{p}) = \left[1 - \nu_0(\omega)\nu_1(\mathbf{p}, \mathbf{k})\right] \nu^{(m_1)}((\omega, \mathbf{p}))\nu^{(m_2)}((\omega, \mathbf{k}))$$

Define, for  $i \leq m \leq j$ ,  $m_1, m_2 \geq m$  and  $s_1, s_2 \in \Sigma_m$

$$\mathcal{D}_{s_1, s_2}^{m_1, m_2}(p, k) = \delta(k_0)\phi(p_0)\Delta_{s_1, s_2}^{m_1, m_2}(\mathbf{p}, \mathbf{k})$$

where

$$\Delta_{s_1, s_2}^{m_1, m_2}(\mathbf{p}, \mathbf{k}) = \int d\omega \frac{\mu^{(m_1, m_2)}(\omega, \mathbf{k}, \mathbf{p})\chi_{s_1}(0, \mathbf{p})\chi_{s_2}(0, \mathbf{k})}{[i\omega - e'(\omega, \mathbf{p})][i\omega - e'(\omega, \mathbf{k})]}$$

and  $\phi$  was defined at the beginning of the proof of Proposition III.22. Define

$$\mathcal{D}_{s_1, s_2}^{(m)}(p, k) = \sum_{\substack{m_1, m_2 \geq 1 \\ \min\{m_1, m_2\}=m}} \mathcal{D}_{s_1, s_2}^{m_1, m_2}(p, k) \quad (\text{III.23})$$

As  $\phi(p_0) = 1$  for all  $|p_0| \leq \frac{M+4\sqrt{2M}}{M^j}$ ,

$$\tilde{\mathcal{C}}^{[i, j]} - \mathcal{M} = \sum_{m=i}^j \sum_{s_1, s_2 \in \Sigma_m} \mathcal{D}_{s_1, s_2}^{(m)} \quad \text{if } |p_0| \leq \frac{M+4\sqrt{2M}}{M^j}$$

Observe that the kernels of  $\tilde{\mathcal{C}}^{[i, j]}$ ,  $\mathcal{M}$  and  $\mathcal{D}_{s_1, s_2}^{(m)}(p, k)$  each contain a factor of  $\delta(k_0)$ . Hence, in the product  $g \bullet (\tilde{\mathcal{C}}^{[i, j]} - \mathcal{M} - \sum_{m=i}^j \mathcal{D}_{s_1, s_2}^{(m)}) \bullet h$ ,  $\check{h}$  is restricted to  $k_0 = 0$ , so that  $p_0 = t_0$ , where  $t$  is the transfer momentum. But  $t \in d$ , so that, by Remark III.12 and (III.16),  $|t_0| \leq \tau_0 + \frac{4\sqrt{2M}}{M^r} \leq \frac{M+4\sqrt{2M}}{M^j}$ . Hence, by Remark III.13 and (III.19),

$$\|g \bullet (\tilde{\mathcal{C}}^{[i, j]} - \mathcal{M}) \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |g|_{\ell, i} |h|_{i, r} \sum_{m=i}^j \frac{1}{l_m} \max_{s_1, s_2 \in \Sigma_m} \|\mathcal{D}_{s_1, s_2}^{(m)}\|_{\text{bubble}}$$

The Proposition follows from Lemma III.25 below. ■

**Lemma III.25** *Let  $i \leq m \leq j$  and  $s_1, s_2 \in \Sigma_m$ . Then*

$$\|\mathcal{D}_{s_1, s_2}^{(m)}\|_{\text{bubble}} \leq \text{const} \begin{cases} \mathfrak{l}_m & \text{if } m = i, i+1 \\ (j-m+1) \frac{M^m}{M^j} \mathfrak{l}_m & \text{if } m \geq i+2 \end{cases}$$

**Proof:** Fix any  $m_1, m_2$  with  $\min\{m_1, m_2\} = m$ . If  $\omega$  is in the support of  $\mu^{(m_1, m_2)}(\omega, \mathbf{k}, \mathbf{p})$  for some  $\mathbf{k}, \mathbf{p}$  then  $|\omega| \leq \frac{\text{const}}{M^{\max\{m_1, m_2\}}}$ . In the case when  $m > i+1$ ,  $|\omega|$  is restricted even farther. Then, in the support of  $\mu^{(m_1, m_2)}(\omega, \mathbf{k}, \mathbf{p})$ , both  $|i\omega - e(\mathbf{p})| \leq \frac{\sqrt{2M}}{M^{m_1}} \leq \frac{\sqrt{2M}}{M^{i+2}}$  and  $|i\omega - e(\mathbf{k})| \leq \frac{\sqrt{2M}}{M^{m_2}} \leq \frac{\sqrt{2M}}{M^{i+2}}$  and hence  $|\omega|, |e(\mathbf{p})|, |e(\mathbf{k})| \leq \frac{\sqrt{2M}}{M^{i+2}}$ . But  $\nu_1(\mathbf{p}, \mathbf{k}) = 1$  whenever  $|e(\mathbf{p})|, |e(\mathbf{k})| \leq \frac{1}{M^{i+1/2}}$ . Hence on the support of  $\mu^{(m_1, m_2)}(\omega, \mathbf{k}, \mathbf{p})$ ,  $|\omega| \leq \frac{\sqrt{2M}}{M^{i+2}}$  and  $\nu_0(\omega) \neq 1$ . This forces  $|\omega| \leq \frac{\sqrt{2}}{M^{j-1/2}} \leq \frac{1}{M^{j-3/4}}$ . Set

$$b(m_1, m_2) = \begin{cases} \frac{\text{const}}{M^{\max\{m_1, m_2\}}} & \text{if } m = i, i+1 \\ \min \left\{ \frac{1}{M^{j-3/4}}, \frac{\text{const}}{M^{\max\{m_1, m_2\}}} \right\} & \text{if } m \geq i+2 \end{cases}$$

Thus

$$\Delta_{s_1, s_2}^{m_1, m_2}(\mathbf{p}, \mathbf{k}) = \int_{-b(m_1, m_2)}^{b(m_1, m_2)} d\omega \frac{\mu^{(m_1, m_2)}(\omega, \mathbf{k}, \mathbf{p}) \chi_{s_1}(\mathbf{p}) \chi_{s_2}(\mathbf{k})}{[i\omega - e'(\omega, \mathbf{p})][i\omega - e'(\omega, \mathbf{k})]}$$

By Lemma A.3, the Fourier transform of  $\Delta_{s_1, s_2}^{m_1, m_2}(\mathbf{p}, \mathbf{k})$  obeys

$$\begin{aligned} \int d\mathbf{z}_1 \sup_{\mathbf{z}_2} |\hat{\Delta}_{s_1, s_2}^{m_1, m_2}(\mathbf{z}_1, \mathbf{z}_2)| &\leq \text{const } b(m_1, m_2) \mathfrak{l}_m^2 \frac{M^{m_1}}{\mathfrak{l}_{m_1}} \\ \int d\mathbf{z}_2 \sup_{\mathbf{z}_1} |\hat{\Delta}_{s_1, s_2}^{m_1, m_2}(\mathbf{z}_1, \mathbf{z}_2)| &\leq \text{const } b(m_1, m_2) \mathfrak{l}_m^2 \frac{M^{m_2}}{\mathfrak{l}_{m_2}} \end{aligned}$$

As the particle-hole bubble propagator associated to  $\mathcal{D}_{s_1, s_2}^{(m)}(p, k)$  by (III.8), namely

$$\mathcal{D}_{s_1, s_2}^{m_1, m_2}(x_1, x_2, y_1, y_2) = \hat{\phi}(y_{1,0} - x_{1,0}) \hat{\Delta}_{s_1, s_2}^{m_1, m_2}(\mathbf{y}_1 - \mathbf{x}_1, \mathbf{x}_2 - \mathbf{y}_2),$$

is independent of  $x_{2,0}$  and  $y_{2,0}$ ,  $\mathcal{Z} \circ \mathcal{D}_{s_1, s_2}^{m_1, m_2} \circ \mathcal{Z}^t = \mathcal{D}_{s_1, s_2}^{m_1, m_2}$  and we have, by Lemma III.18.iv,

$$\begin{aligned} \|\mathcal{D}_{s_1, s_2}^{m_1, m_2}\|_{\text{bubble}} &\leq \text{const} \min \left\{ \sup_{x_1, x_2} \int dy_1 \sup_{\mathbf{y}_2} |\hat{\phi}(y_{1,0} - x_{1,0}) \hat{\Delta}_{s_1, s_2}^{m_1, m_2}(\mathbf{y}_1 - \mathbf{x}_1, \mathbf{x}_2 - \mathbf{y}_2)|, \right. \\ &\quad \left. \sup_{x_1, x_2} \int dy_{1,0} d\mathbf{y}_2 \sup_{\mathbf{y}_1} |\hat{\phi}(y_{1,0} - x_{1,0}) \hat{\Delta}_{s_1, s_2}^{m_1, m_2}(\mathbf{y}_1 - \mathbf{x}_1, \mathbf{x}_2 - \mathbf{y}_2)| \right\} \\ &\leq \text{const} \min \left\{ \int d\mathbf{z}_1 \sup_{\mathbf{z}_2} |\hat{\Delta}_{s_1, s_2}^{m_1, m_2}(\mathbf{z}_1, \mathbf{z}_2)|, \int d\mathbf{z}_2 \sup_{\mathbf{z}_1} |\hat{\Delta}_{s_1, s_2}^{m_1, m_2}(\mathbf{z}_1, \mathbf{z}_2)| \right\} \\ &\leq \text{const } b(m_1, m_2) \mathfrak{l}_m^2 \min \left\{ \frac{M^{m_1}}{\mathfrak{l}_{m_1}}, \frac{M^{m_2}}{\mathfrak{l}_{m_2}} \right\} \\ &= \text{const } b(m_1, m_2) \mathfrak{l}_m M^m \end{aligned}$$

If  $m \in \{i, i+1\}$

$$\|\mathcal{D}_{s_1, s_2}^{(m)}\|_{\text{bubble}} \leq \sum_{\min\{m_1, m_2\}=m} \|\mathcal{D}_{s_1, s_2}^{m_1, m_2}\|_{\text{bubble}} \leq \text{const} \sum_{m' \geq m} \frac{1}{M^{m'}} \mathfrak{l}_m M^m \leq \text{const} \mathfrak{l}_m$$

and if  $m \geq i+2$

$$\begin{aligned} \|\mathcal{D}_{s_1, s_2}^{(m)}\|_{\text{bubble}} &\leq \sum_{\min\{m_1, m_2\}=m} \|\mathcal{D}_{s_1, s_2}^{m_1, m_2}\|_{\text{bubble}} \\ &\leq \text{const} \sum_{m'=m}^j \frac{1}{M^j} \mathfrak{l}_m M^m + \text{const} \sum_{m'>j} \frac{1}{M^{m'}} \mathfrak{l}_m M^m \\ &\leq \text{const} (j - m + 1) \frac{M^m}{M^j} \mathfrak{l}_m \end{aligned}$$

■

### Model Propagator in Position Space

**Lemma III.26** *The set  $\mathbf{d} = \{ \mathbf{p}_1 - \mathbf{p}_2 \mid \mathbf{p}_1 \in \kappa_1, \mathbf{p}_2 \in \kappa_2 \}$  is contained in a rectangle with two sides of length  $\text{const} \mathfrak{l}_j$  and two sides of length  $\text{const} \frac{1}{M^j}$ .*

**Proof:** For every sector  $\sigma \in \Sigma_r$ , let  $\mathbf{k}_\sigma$  be the centre of  $\sigma \cap F$ ,  $\mathbf{t}_\sigma$  a unit tangent vector to  $F$  at  $\mathbf{k}_\sigma$  and

$$\sigma = \{ \mathbf{k} \in \mathbb{R}^2 \mid (k_0, \mathbf{k}) \in \sigma \text{ for some } k_0 \in \mathbb{R} \}$$

Then  $\sigma$  is contained in a rectangle  $R_\sigma$  centered at  $\mathbf{k}_\sigma$  with two sides parallel to  $\mathbf{t}_\sigma$  of length  $\text{const} \mathfrak{l}_r$  and two sides perpendicular to  $\mathbf{t}_\sigma$  of length  $\text{const} \frac{1}{M^r}$ . If at least one of  $\kappa_1$  and  $\kappa_2$  are in  $\mathbb{M}$ , the claim follows since  $j \leq r$ . So assume that  $\kappa_1, \kappa_2 \in \Sigma_r$ . Then the distance between  $\mathbf{k}_{\kappa_1}$  and  $\mathbf{k}_{\kappa_2}$  is at most  $|\tau| + 5\mathfrak{l}_r$  and therefore the angle between  $\mathbf{t}_{\kappa_1}$  and  $\mathbf{t}_{\kappa_2}$  is at most  $\text{const} (|\tau| + \mathfrak{l}_r)$ . Consequently  $\mathbf{d}$  is contained in a rectangle with two sides parallel to  $\mathbf{t}_{\kappa_1}$  of length  $\text{const} \mathfrak{l}_r$  and two sides perpendicular to  $\mathbf{t}_{\kappa_1}$  of length

$$\text{const} \left( \frac{1}{M^r} + (|\tau| + \mathfrak{l}_r) \mathfrak{l}_r \right) \leq \text{const} \left( \frac{1}{M^r} + |\tau| \mathfrak{l}_r \right)$$

By (III.16),  $|\tau| \leq \max\{\frac{1}{M^j}, r^3 \mathfrak{l}_r\}$ , so that  $|\tau| \mathfrak{l}_r \leq \text{const} \frac{1}{M^j}$ .

■

Fix two mutually perpendicular unit vector  $\mathbf{t}$  and  $\mathbf{n}$  and a rectangle  $R$ , with two sides parallel to  $\mathbf{t}$  of length  $\text{const} \mathfrak{l}_j$  and two sides parallel to  $\mathbf{n}$  of length  $\text{const} \frac{1}{M^j}$ , such that

$\mathbf{d} \subset R$ . By Lemma III.26, such a rectangle exists. Let  $\rho(\mathbf{t})$  be identically one on  $R$ , be supported on a set of area twice that of  $R$  and obey

$$\left| (\mathbf{n} \cdot \partial_{\mathbf{t}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{t}})^{\alpha_2} \rho(\mathbf{t}) \right| \leq \text{const } M^{\alpha_1 j} \frac{1}{l_j^{\alpha_2}}$$

for all  $\alpha_1, \alpha_2 \leq 2$ . Define  $\mathcal{M}_\rho$  as in Definition III.3. Then

$$\|g \bullet \mathcal{M} \bullet h\|_{\kappa_1, \kappa_2} = \|g \bullet \mathcal{M}_\rho \bullet h\|_{\kappa_1, \kappa_2} \quad (\text{III.24})$$

### Proposition III.27

$$\|g \bullet \mathcal{M} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const} \max_{\substack{\alpha_r, \alpha_l \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_r| + |\alpha_l| \leq 3}} |g|_{\ell, i}^{[0, 0, \alpha_r]} |h|_{i, r}^{[\alpha_l, 0, 0]}$$

**Proof:** Write

$$\mathcal{M}_\rho(p, k) = \sum_{s_1, s_2 \in \Sigma_i} \mathcal{M}_{s_1, s_2}(p, k)$$

where, for  $s_1, s_2 \in \Sigma_i$

$$\mathcal{M}_{s_1, s_2}(p, k) = \mathcal{M}(p, k) \rho(\mathbf{p} - \mathbf{k}) \chi_{s_1}(\mathbf{p}) \chi_{s_2}(\mathbf{k})$$

By (III.19),

$$\begin{aligned} \|g \bullet \mathcal{M}_\rho \bullet h\|_{\kappa_1, \kappa_2} &\leq \sum_{\substack{s_1, s_2 \in \Sigma_i \\ (s_1 - s_2) \cap d \neq \emptyset}} \|g \bullet \mathcal{M}_{s_1, s_2} \bullet h\|_{\kappa_1, \kappa_2} \\ &\leq \frac{\text{const}}{l_i} \max_{s_1, s_2 \in \Sigma_i} \|g \bullet \mathcal{M}_{s_1, s_2} \bullet h\|_{\kappa_1, \kappa_2} \end{aligned} \quad (\text{III.25})$$

Define, for  $v_1, v_2 \in \Sigma_i$ ,

$$h_{v_1, v_2}(x_1, x_2, y_3, y_4) = h((x_1, v_1), (x_2, v_2), y_3, y_4) \quad \text{where } y_\nu = \begin{cases} \kappa_{\nu-2} & \text{if } \kappa_{\nu-2} \in \mathbb{M} \\ (x_\nu, \kappa_{\nu-2}) & \text{if } \kappa_{\nu-2} \in \Sigma_r \end{cases}$$

Observe that  $h_{v_1, v_2}$  is a function on  $(\mathbb{R} \times \mathbb{R}^2)^{2+n_r}$  where  $n_r = \#\{ \nu \in \{1, 2\} \mid \kappa_\nu \in \Sigma_r \}$ . Also define, for  $u_3, u_4 \in \Sigma_i$  and  $\lambda_1, \lambda_2 \in \mathbb{M} \cup \Sigma_\ell$ ,

$$g_{\lambda_1, \lambda_2; u_3, u_4}(y_1, y_2, x_3, x_4) = g(y_1, y_2, (x_3, u_3), (x_4, u_4)) \quad \text{where } y_\nu = \begin{cases} \lambda_\nu & \text{if } \lambda_\nu \in \mathbb{M} \\ (x_\nu, \lambda_\nu) & \text{if } \lambda_\nu \in \Sigma_r \end{cases}$$

Observe that  $g_{\lambda_1, \lambda_2; u_3, u_4}$  is a function on  $(\mathbb{R} \times \mathbb{R}^2)^{2+n_1}$  where  $n_1 = \#\{ \nu \in \{1, 2\} \mid \lambda_\nu \in \Sigma_\ell \}$ . Then, for all  $s_1, s_2 \in \Sigma_i$ ,

$$\|g \bullet \mathcal{M}_{s_1, s_2} \bullet h\|_{\kappa_1, \kappa_2} \leq \sup_{\lambda_1, \lambda_2} \sum_{\substack{u_3, u_4 \in \Sigma_i \\ v_1, v_2 \in \Sigma_i}} \|g_{\lambda_1, \lambda_2; u_3, u_4} \circ \mathcal{M}_{s_1, s_2} \circ h_{v_1, v_2}\|_{1, \infty} \quad (\text{III.26})$$

By conservation of momentum, the convolution  $g_{\lambda_1, \lambda_2; u_3, u_4} \circ \mathcal{M}_{s_1, s_2} \circ h_{v_1, v_2}$  vanishes identically unless

$$\begin{aligned} u_3 \cap s_1 &\neq \emptyset & s_1 \cap v_1 &\neq \emptyset \\ u_4 \cap s_2 &\neq \emptyset & s_2 \cap v_2 &\neq \emptyset \end{aligned}$$

There only  $3^4$  quadruples  $(u_3, u_4, v_1, v_2)$  satisfying these conditions, so that, by (III.24), (III.25) and (III.26),

$$\|g \bullet \mathcal{M} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } \frac{1}{\ell_i} \sup_{\lambda_1, \lambda_2} \max_{\substack{s_1, s_2 \in \Sigma_i \\ u_3, u_4 \in \Sigma_i \\ v_1, v_2 \in \Sigma_i}} \|g_{\lambda_1, \lambda_2; u_3, u_4} \circ \mathcal{M}_{s_1, s_2} \circ h_{v_1, v_2}\|_{1, \infty} \quad (\text{III.27})$$

Fix  $\lambda_1, \lambda_2, s_1, s_2, u_3, u_4, v_1, v_2$  and denote  $g' = g_{\lambda_1, \lambda_2; u_3, u_4}$  and  $h' = h_{v_1, v_2}$ . Write the convolution

$$\begin{aligned} g' \circ \mathcal{M}_{s_1, s_2} \circ h' &= \int d^3 z_1 d^3 z_2 d^3 y_1 d^3 y_2 \frac{d^3 \mathbf{p}}{(2\pi)^3} \frac{d^3 \mathbf{k}}{(2\pi)^3} d\omega e^{i\langle \mathbf{p}, \mathbf{z}_1 - \mathbf{y}_1 \rangle} e^{i\langle \mathbf{k}, \mathbf{y}_2 - \mathbf{z}_2 \rangle} g'(\cdot, \cdot, z_1, z_2) \\ &\quad \delta(k_0) \frac{\nu_0(\omega) \nu_1(\mathbf{p}, \mathbf{k}) \chi_{s_1}(\mathbf{p}) \chi_{s_2}(\mathbf{k})}{[i\omega - e'(\omega, \mathbf{p})][i\omega - e'(\omega, \mathbf{k})]} \rho(\mathbf{p} - \mathbf{k}) h'(y_1, y_2, \cdot, \cdot) \\ &= \int d^3 z_1 d^3 z_2 d^2 \mathbf{y}_1 d^3 y_2 \frac{d^2 \mathbf{p}}{(2\pi)^2} \frac{d^2 \mathbf{k}}{(2\pi)^2} \frac{d\omega}{2\pi} e^{i\mathbf{p} \cdot (\mathbf{z}_1 - \mathbf{y}_1)} e^{i\mathbf{k} \cdot (\mathbf{y}_2 - \mathbf{z}_2)} g'(\cdot, \cdot, z_1, z_2) \\ &\quad \frac{\nu_0(\omega) \nu_1(\mathbf{p}, \mathbf{k}) \chi_{s_1}(\mathbf{p}) \chi_{s_2}(\mathbf{k})}{[i\omega - e'(\omega, \mathbf{p})][i\omega - e'(\omega, \mathbf{k})]} \rho(\mathbf{p} - \mathbf{k}) h'((z_{1,0}, \mathbf{y}_1), y_2, \cdot, \cdot) \\ &= \int d^3 z_1 d^3 z_2 d^2 \mathbf{y}_1 d^3 y_2 \frac{d^2 \mathbf{t}}{(2\pi)^2} \frac{d^2 \mathbf{k}}{(2\pi)^2} \frac{d\omega}{2\pi} e^{i\mathbf{t} \cdot (\mathbf{z}_1 - \mathbf{y}_1)} e^{i\mathbf{k} \cdot (\mathbf{z}_1 - \mathbf{z}_2 + \mathbf{y}_2 - \mathbf{y}_1)} g'(\cdot, \cdot, z_1, z_2) \\ &\quad \frac{\nu_0(\omega) \nu_1(\mathbf{k} + \mathbf{t}, \mathbf{k}) \chi_{s_1}(\mathbf{k} + \mathbf{t}) \chi_{s_2}(\mathbf{k})}{[i\omega - e'(\omega, \mathbf{k} + \mathbf{t})][i\omega - e'(\omega, \mathbf{k})]} \rho(\mathbf{t}) h'((z_{1,0}, \mathbf{y}_1), y_2, \cdot, \cdot) \\ &= \int d^3 z_1 d^3 z_2 d^2 \mathbf{y}_1 d^3 y_2 g'(\cdot, \cdot, z_1, z_2) \widehat{B}_{s_1, s_2}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{y}_1, \mathbf{y}_2) h'((z_{1,0}, \mathbf{y}_1), y_2, \cdot, \cdot) \end{aligned}$$

where

$$\widehat{B}_{s_1, s_2}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{y}_1, \mathbf{y}_2) = \int \frac{d^2 \mathbf{t}}{(2\pi)^2} e^{i\mathbf{t} \cdot (\mathbf{z}_1 - \mathbf{y}_1)} B_{s_1, s_2}(\mathbf{t}, \mathbf{z}_1 - \mathbf{z}_2 + \mathbf{y}_2 - \mathbf{y}_1) \rho(\mathbf{t})$$

with

$$B_{s_1, s_2}(\mathbf{t}, \mathbf{w}) = \int \frac{d^2 \mathbf{k}}{(2\pi)^2} \frac{d\omega}{2\pi} e^{i\mathbf{k} \cdot \mathbf{w}} \frac{\nu_0(\omega) \nu_1(\mathbf{k} + \mathbf{t}, \mathbf{k}) \chi_{s_1}(\mathbf{k} + \mathbf{t}) \chi_{s_2}(\mathbf{k})}{[i\omega - e'(\omega, \mathbf{k} + \mathbf{t})][i\omega - e'(\omega, \mathbf{k})]}$$

Recall that  $M^i \leq \mathfrak{l}_j M^j$  and that  $p^{(i)} = 0$  for  $i > j + 1$ . By Theorem B.2, with  $u(\mathbf{k}, \mathbf{t}) = e^{i\mathbf{k} \cdot \mathbf{w}} \nu_1(\mathbf{k} + \mathbf{t}, \mathbf{k}) \chi_{s_1}(\mathbf{k} + \mathbf{t}) \chi_{s_2}(\mathbf{k})$ , there is an  $a > 1$  such that

$$|\widehat{B}_{s_1, s_2}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{y}_1, \mathbf{y}_2)| \leq \text{const} \frac{\mathfrak{l}_i \mathfrak{l}_j}{M^j} \frac{1 + |\mathbf{z}_2 - \mathbf{z}_1 + \mathbf{y}_1 - \mathbf{y}_2|^3 / M^{3i}}{[1 + (\mathfrak{n} \cdot (\mathbf{z}_1 - \mathbf{y}_1) / M^j)^{3/2}][1 + |\mathfrak{l}_j \mathbf{t} \cdot (\mathbf{z}_1 - \mathbf{y}_1)|^a]}$$

Since  $\sup_{\mathbf{z}_1} \int d^2 \mathbf{y}_1 \frac{\mathfrak{l}_j}{M^j} \frac{1}{[1 + (\mathfrak{n} \cdot (\mathbf{z}_1 - \mathbf{y}_1) / M^j)^{3/2}][1 + |\mathfrak{l}_j \mathbf{t} \cdot (\mathbf{z}_1 - \mathbf{y}_1)|^a]} \leq \text{const}$ , we have

$$\|g' \circ \mathcal{M}_{s_1, s_2} \circ h'\|_{1, \infty} \leq \text{const} \mathfrak{l}_i \max_{\substack{\alpha_r, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_r| + |\alpha_1| \leq 3}} |g|_{\ell, i}^{[0, 0, \alpha_r]} |h|_{i, r}^{[\alpha_1, 0, 0]} \quad (\text{III.28})$$

and the Proposition follows by (III.27). ■

**Proof of Theorem III.15:** By Propositions III.19, III.22, III.24 and III.27

$$\begin{aligned} & \|g \bullet \mathcal{C}^{[i, j]} \bullet h\|_{\kappa_1, \kappa_2} \\ & \leq \|g \bullet (\mathcal{C}^{[i, j]} - \mathcal{Z} \bullet \mathcal{C}^{[i, j]} \bullet \mathcal{Z}^t) \bullet h\|_{\kappa_1, \kappa_2} + \|g \bullet (\mathcal{Z} \bullet \mathcal{C}^{[i, j]} \bullet \mathcal{Z}^t - \widetilde{\mathcal{C}}^{[i, j]}) \bullet h\|_{\kappa_1, \kappa_2} \\ & \quad + \|g \bullet (\widetilde{\mathcal{C}}^{[i, j]} - \mathcal{M}) \bullet h\|_{\kappa_1, \kappa_2} + \|g \bullet \mathcal{M} \bullet h\|_{\kappa_1, \kappa_2} \\ & \leq \text{const} \max_{\substack{\alpha_r, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_r| + |\alpha_1| \leq 3}} |g|_{\ell, i}^{[0, 0, \alpha_r]} |h|_{i, r}^{[\alpha_1, 0, 0]} \end{aligned}$$

as desired. ■

## The Infrared Limit – Nonzero Transfer Momentum

**Lemma III.28** *Let  $\kappa_1, \kappa_2 \in \mathbb{M}$  and  $g$  and  $h$  be sectorized, translation invariant functions on  $\mathfrak{Y}_{\Sigma_j}^4$  and  $\mathfrak{Y}_{\Sigma_j}^2 \times \mathbb{M}^2$ , respectively. Then*

$$\|g \bullet \mathcal{C}^{(j)} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const} |g|_{j, j} \|h\|_{\kappa_1, \kappa_2} \left\{ \sqrt{\mathfrak{l}_j} + \min \left\{ 1, \frac{1}{|\kappa_1 - \kappa_2| M^j} \right\} \right\}$$

Furthermore, for  $(\text{loc}, \mu, \mu') \in \{(\text{top}, 1, 3), (\text{bot}, 2, 4), (\text{mid}, 1, 3), (\text{mid}, 2, 4)\}$  and  $\beta \in \Delta$ ,

$$\|g \bullet \mathbf{D}_{\mu, \mu'}^\beta \mathcal{C}_{\text{loc}}^{[j, j]} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const} |g|_{j, j} \|h\|_{\kappa_1, \kappa_2} M^{|\beta|j} \left\{ \sqrt{\mathfrak{l}_j} + \min \left\{ 1, \frac{1}{|\kappa_1 - \kappa_2| M^j} \right\} \right\}$$

**Proof:** Write  $\mathcal{C}^{(j)} = \sum_{s_1, s_2 \in \Sigma_j} [\mathcal{C}_{\text{top}, j, s_1, s_2}^{(j)} + \mathcal{C}_{\text{mid}, j, s_1, s_2}^{(j)} + \mathcal{C}_{\text{bot}, j, s_1, s_2}^{(j)}]$ . As in Lemma III.11,

$$\|g \bullet D_{\mu, \mu'}^\beta \mathcal{C}_{\text{loc}}^{[j, j]} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |g|_{j, j} \|h\|_{\kappa_1, \kappa_2} \sum_{\substack{s_1, s_2 \in \Sigma_j \\ \kappa_1 - \kappa_2 \in s_1 - s_2}} \|D_{\mu, \mu'}^\beta \mathcal{C}_{\text{loc}, j, s_1, s_2}^{(j)}\|_{\text{bubble}}$$

By Lemma III.14, for  $(\text{loc}, \mu, \mu') \in \{(\text{top}, 1, 3), (\text{bot}, 2, 4), (\text{mid}, 1, 3), (\text{mid}, 2, 4)\}$  and  $\beta \in \Delta$ ,

$$\|D_{\mu, \mu'}^\beta \mathcal{C}_{\text{loc}, j, s_1, s_2}^{(j)}\|_{\text{bubble}} \leq \text{const } l_j M^{|\beta|j}$$

and by Lemma C.2,

$$\#\{ (s_1, s_2) \in \Sigma_j \mid \kappa_1 - \kappa_2 \in s_1 - s_2 \} \leq \text{const } \left\{ \frac{1}{\sqrt{l_j}} + \frac{1}{l_j} \min \left\{ 1, \frac{1}{|\kappa_1 - \kappa_2| M^j} \right\} \right\}$$

so that the desired bounds follow. ■

Recall, from just before Lemma II.28, that

$$\begin{aligned} & \mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t, \sigma_1, \dots, \sigma_4) \\ &= \left[ (F^{(i_1)} + L^{(i_1)} f) \bullet \mathcal{C}^{[\max\{i_1, i_2\}, j]} \bullet (F^{(i_2)} + L^{(i_2)} f) \bullet \dots \right. \\ & \quad \left. \dots \bullet \mathcal{C}^{[\max\{i_\ell, i_{\ell+1}\}, j]} \bullet (F^{(i_{\ell+1})} + L^{(i_{\ell+1})} f) \right]_{i_1, i_2, i_3, i_4=0}^{((q+\frac{t}{2}, \sigma_1), (q-\frac{t}{2}, \sigma_2), (q'+\frac{t}{2}, \sigma_3), (q'-\frac{t}{2}, \sigma_4))} \end{aligned}$$

**Lemma III.29** *For  $t \neq 0$ , the limit*

$$\mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}(q, q', t, \sigma_1, \dots, \sigma_4) = \lim_{j \rightarrow \infty} \mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t, \sigma_1, \dots, \sigma_4)$$

*exists. The limit is continuous in  $(q, q', t)$  for  $t \neq 0$ .*

**Proof:** It suffices to consider separately the spin and charge parts, in the sense of Lemma II.8, of  $\mathfrak{L}_{\ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t, \sigma_1, \dots, \sigma_4)$ . We denote them  $\mathfrak{L}_{X, \ell, i_1, \dots, i_{\ell+1}}^{(j)}(q, q', t)$  with  $X = S, C$ . Define  $L_{X, j_1, \dots, j_\ell}(q, q', t)$  to be the  $X$  part of

$$\begin{aligned} & \left[ (F^{(i_1)} + L^{(i_1)} f) \bullet \mathcal{C}^{(j_1)} \bullet (F^{(i_2)} + L^{(i_2)} f) \bullet \dots \right. \\ & \quad \left. \dots \bullet \mathcal{C}^{(j_\ell)} \bullet (F^{(i_{\ell+1})} + L^{(i_{\ell+1})} f) \right]_{i_1, i_2, i_3, i_4=0}^{((q+\frac{t}{2}, \sigma_1), (q-\frac{t}{2}, \sigma_2), (q'+\frac{t}{2}, \sigma_3), (q'-\frac{t}{2}, \sigma_4))} \end{aligned}$$

and

$$\Delta \mathfrak{L}_X^{(p)}(q, q', t) = \sum_{\substack{\max\{i_1, i_2\} \leq j_1 \leq j \\ \vdots \\ \max\{i_\ell, i_{\ell+1}\} \leq j_\ell \leq j \\ \max\{j_1, \dots, j_\ell\} = p}} L_{X, j_1, \dots, j_\ell}(q, q', t)$$



Then

$$\mathfrak{L}_{X,\ell,i_1,\dots,i_{\ell+1}}^{(j)}(q,q',t,\sigma_1,\dots,\sigma_4) = \sum_{p=2}^j \Delta \mathfrak{L}_X^{(p)}(q,q',t,\sigma_1,\dots,\sigma_4)$$

By repeated use of Lemma III.28, (II.8) and its  $X = C$  analog,

$$\|L_{X,j_1,\dots,j_\ell}\|_{\kappa_1,\kappa_2} \leq \text{const} \prod_{m=1}^{\ell} \left\{ \sqrt{\mathfrak{l}_{j_m}} + \min \left\{ 1, \frac{1}{|\kappa_1 - \kappa_2| M^{j_m}} \right\} \right\}$$

with the constant depending on  $\ell$  and  $i_1, \dots, i_{\ell+1}$ . Summing this bound over  $j_1, \dots, j_\ell$  with  $\max\{j_1, \dots, j_\ell\} = p$

$$\|\Delta \mathfrak{L}_X^{(p)}\|_{\kappa_1,\kappa_2} \leq \text{const} \left\{ \sqrt{\mathfrak{l}_p} + \min \left\{ 1, \frac{1}{|\kappa_1 - \kappa_2| M^p} \right\} \right\}$$

The constant now depends on  $|\kappa_1 - \kappa_2|$  as well but remains finite as long as  $\kappa_1 \neq \kappa_2$ . The existence of the limit  $j \rightarrow \infty$  is now a consequence of the Lebesgue dominated convergence theorem. Similarly, if  $D$  is any first order differential operator in  $q, q'$  or  $t$ ,

$$\begin{aligned} \|DL_{X,j_1,\dots,j_\ell}\|_{\kappa_1,\kappa_2} &\leq \text{const} M^{\max\{j_1,\dots,j_\ell\}} \prod_{m=1}^{\ell} \left\{ \sqrt{\mathfrak{l}_{j_m}} + \min \left\{ 1, \frac{1}{|\kappa_1 - \kappa_2| M^{j_m}} \right\} \right\} \\ \|D\Delta \mathfrak{L}_X^{(p)}\|_{\kappa_1,\kappa_2} &\leq \text{const} M^p \left\{ \sqrt{\mathfrak{l}_p} + \min \left\{ 1, \frac{1}{|\kappa_1 - \kappa_2| M^p} \right\} \right\} \end{aligned}$$

Continuity now follows from

$$\frac{|g(x) - g(y)|}{|x - y|^\epsilon} = |g(x) - g(y)|^{1-\epsilon} \left[ \frac{|g(x) - g(y)|}{|x - y|} \right]^\epsilon \leq [2 \sup |g(x)|]^{1-\epsilon} [\sup |\nabla g(x)|]^\epsilon$$

and the observation that

$$\sum_{p=1}^{\infty} M^{\epsilon p} \left\{ \sqrt{\mathfrak{l}_p} + \min \left\{ 1, \frac{1}{|\kappa_1 - \kappa_2| M^p} \right\} \right\} < \infty \quad \text{if } \epsilon < \frac{\aleph}{2} \text{ and } \kappa_1 \neq \kappa_2$$

■

## The Infrared Limit – Reduction to Factorized Cutoffs

Recall from (II.13) that

$$\begin{aligned} \mathcal{C}^{[i,j]}(p,k) &= \frac{\nu^{(\geq i)}(p) \nu^{(\geq i)}(k) - \nu^{(\geq j+1)}(p) \nu^{(\geq j+1)}(k)}{[ip_0 - e'(p)][ik_0 - e'(k)]} \\ \mathcal{A}_{i,j}(p,k) &= \frac{\nu^{(\geq i)}(p) \nu^{(\geq i)}(k) [1 - \nu_j(e(\mathbf{p})) \nu_j(e(\mathbf{k}))]}{[ip_0 - e'(p)][ik_0 - e'(k)]} \\ \mathcal{B}_{i,j}(p,k) &= \frac{\nu^{(\geq i)}(p) \nu^{(\geq i)}(k) [1 - \nu_j(p_0) \nu_j(k_0)]}{[ip_0 - e'(p)][ik_0 - e'(k)]} \end{aligned}$$

where  $e'(k) = e(\mathbf{k}) - v(k)$  and

$$\nu_j(\omega) = \sum_{m=j}^{\infty} \nu(M^{2m} \omega^2)$$

with  $\nu$  being the single scale cutoff introduced in Definition I.2.

**Proposition III.30** *Let  $1 \leq i \leq j$  and  $\ell, r \geq 1$ . Let  $\kappa_1, \kappa_2 \in \mathbb{M}$  with and  $g$  and  $h$  be sectorized, translation invariant functions on  $\mathfrak{Y}_{\ell,i}$  and  $\mathfrak{Y}_{i,r}$  respectively. If  $|\kappa_1 - \kappa_2| > 0$ , then*

$$\|g \bullet (\mathcal{C}^{[i,j]} - \mathcal{A}_{i,j}) \bullet h\|_{\kappa_1, \kappa_2}, \|g \bullet (\mathcal{C}^{[i,j]} - \mathcal{B}_{i,j}) \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |g|_{\ell,i} \|h\|_{\kappa_1, \kappa_2} j \left\{ \sqrt{\mathfrak{l}_j} + \frac{1}{|\kappa_1 - \kappa_2| M^j \mathfrak{l}_j} \right\}$$

**Proof:** Let

$$\mathcal{D}(p, k) = \frac{\nu^{(\geq j+1)}(p) \nu^{(\geq j+1)}(k)}{[ip_0 - e'(p)][ik_0 - e'(k)]} \text{ or } \frac{\nu^{(\geq i)}(p) \nu^{(\geq i)}(k) \nu_j(e(\mathbf{p})) \nu_j(e(\mathbf{k}))}{[ip_0 - e'(p)][ik_0 - e'(k)]} \text{ or } \frac{\nu^{(\geq i)}(p) \nu^{(\geq i)}(k) \nu_j(p_0) \nu_j(k_0)}{[ip_0 - e'(p)][ik_0 - e'(k)]}$$

It suffices to prove that

$$\|g \bullet \mathcal{D} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |g|_{\ell,i} \|h\|_{\kappa_1, \kappa_2} j \left\{ \sqrt{\mathfrak{l}_j} + \frac{1}{|\kappa_1 - \kappa_2| M^j \mathfrak{l}_j} \right\}$$

Define, for  $m \geq i$ ,  $m_1, m_2 \geq m$  and  $s_1, s_2 \in \Sigma_m$

$$\mathcal{D}_{s_1, s_2}^{m_1, m_2}(p, k) = \frac{\nu^{(m_1)}(p) \nu^{(m_2)}(k) \chi_{s_1}(p) \chi_{s_2}(k)}{[ip_0 - e'(p)][ik_0 - e'(k)]} \begin{cases} \nu^{(\geq j+1)}(p) \nu^{(\geq j+1)}(k) & \text{or} \\ \nu_j(e(\mathbf{p})) \nu_j(e(\mathbf{k})) & \text{or} \\ \nu_j(p_0) \nu_j(k_0) \end{cases}$$

and

$$\mathcal{D}_{s_1, s_2}^{(m)}(p, k) = \sum_{\substack{m_1, m_2 \geq 1 \\ \min\{m_1, m_2\} = m}} \mathcal{D}_{s_1, s_2}^{m_1, m_2}(p, k)$$

We have

$$\mathcal{D} = \sum_{m=i}^{\infty} \sum_{s_1, s_2 \in \Sigma_m} \mathcal{D}_{s_1, s_2}^{(m)}$$

Hence, as in Lemma III.11,

$$\|g \bullet \mathcal{D} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |g|_{\ell,i} \|h\|_{\kappa_1, \kappa_2} \sum_{m=i}^{\infty} \sum_{\substack{s_1, s_2 \in \Sigma_m \\ \kappa_1 - \kappa_2 \in s_1 - s_2}} \max_{s_1, s_2 \in \Sigma_m} \|\mathcal{D}_{s_1, s_2}^{(m)}\|_{\text{bubble}}$$

To bound  $\|\mathcal{D}_{s_1, s_2}^{(m)}\|_{\text{bubble}}$ , set, for  $s \in \Sigma_m$  and  $n \geq m$ ,

$$c_s^{(n,j)}(k) = \frac{\nu^{(n)}(k) \chi_s(k)}{ik_0 - e'(k)} \begin{cases} \nu^{(\geq j+1)}(k) & \text{or} \\ \nu_j(e(\mathbf{k})) & \text{or} \\ \nu_j(k_0) \end{cases}$$

and denote by  $c_s^{(n,j)}(x)$  its Fourier transform. As in Lemma A.2,

$$\begin{aligned} \|c_s^{(m,j)}(x)\|_{L^1} &\leq \text{const } M^m \frac{\mathfrak{l}_m}{\mathfrak{l}_{\max\{m,j\}}} \\ \|c_s^{(n,j)}(x)\|_{L^\infty} &\leq \text{const } \mathfrak{l}_m M^n \frac{1}{M^n M^{\max\{n,j\}}} \leq \text{const } \frac{\mathfrak{l}_m}{M^{\max\{n,j\}}} \end{aligned}$$

The factor  $\frac{\mathfrak{l}_m}{M^{\max\{n,j\}}}$  in the first inequality arises, when  $j > m$ , from splitting  $s$  up into sectors of length  $\mathfrak{l}_j$ . Hence, by the triangle inequality and (III.7),

$$\begin{aligned}
\|\mathcal{D}_{s_1, s_2}^{(m)}\|_{\text{bubble}} &\leq \sum_{n \geq m} \left[ \|c_{s_1}^{(m,j)}\|_{L^1} \|c_{s_2}^{(n,j)}\|_{L^\infty} + \|c_{s_1}^{(n,j)}\|_{L^\infty} \|c_{s_2}^{(m,j)}\|_{L^1} \right] \\
&\leq \sum_{n > m} \text{const } M^m \frac{\mathfrak{l}_m}{\mathfrak{l}_{\max\{m,j\}}} \frac{\mathfrak{l}_m}{M^{\max\{n,j\}}} \\
&\leq \text{const } \frac{M^m \mathfrak{l}_m^2}{\mathfrak{l}_{\max\{m,j\}}} \sum_{n > m} \frac{1}{M^{\max\{n,j\}}} \\
&\leq \text{const } \frac{M^m \mathfrak{l}_m^2}{\mathfrak{l}_{\max\{m,j\}}} \frac{j}{M^{\max\{m,j\}}}
\end{aligned}$$

By Lemma C.2,

$$\#\{ (s_1, s_2) \in \Sigma_m \mid \kappa_1 - \kappa_2 \in s_1 - s_2 \} \leq \text{const} \left\{ \frac{1}{\sqrt{\mathfrak{l}_m}} + \frac{1}{\mathfrak{l}_m} \min \left\{ 1, \frac{1}{|\kappa_1 - \kappa_2| M^m} \right\} \right\}$$

so that

$$\begin{aligned}
\|g \bullet \mathcal{D} \bullet h\|_{\kappa_1, \kappa_2} &\leq \text{const } |g|_{\ell, i} \|h\|_{\kappa_1, \kappa_2} \sum_{m=i}^{\infty} \left\{ \frac{1}{\sqrt{\mathfrak{l}_m}} + \frac{1}{\mathfrak{l}_m} \min \left\{ 1, \frac{1}{|\kappa_1 - \kappa_2| M^m} \right\} \right\} \frac{M^m \mathfrak{l}_m^2}{\mathfrak{l}_{\max\{m,j\}}} \frac{j}{M^{\max\{m,j\}}} \\
&\leq \text{const } |g|_{\ell, i} \|h\|_{\kappa_1, \kappa_2} \sum_{m=1}^j \left\{ \frac{1}{\sqrt{\mathfrak{l}_m}} + \frac{1}{|\kappa_1 - \kappa_2| M^m \mathfrak{l}_m} \right\} \frac{M^m \mathfrak{l}_m^2}{\mathfrak{l}_j} \frac{j}{M^j} \\
&\quad + \text{const } |g|_{\ell, i} \|h\|_{\kappa_1, \kappa_2} j \sum_{m=j}^{\infty} \left\{ \frac{1}{\sqrt{\mathfrak{l}_m}} + \frac{1}{|\kappa_1 - \kappa_2| M^m \mathfrak{l}_m} \right\} \mathfrak{l}_m \\
&\leq \text{const } |g|_{\ell, i} \|h\|_{\kappa_1, \kappa_2} j \left\{ \frac{1}{M^j \mathfrak{l}_j} M^j \mathfrak{l}_j^{3/2} + \frac{1}{M^j \mathfrak{l}_j} \frac{1}{|\kappa_1 - \kappa_2|} + \sqrt{\mathfrak{l}_j} + \frac{1}{|\kappa_1 - \kappa_2| M^j} \right\} \\
&\leq \text{const } |g|_{\ell, i} \|h\|_{\kappa_1, \kappa_2} j \left\{ \sqrt{\mathfrak{l}_j} + \frac{1}{|\kappa_1 - \kappa_2| M^j \mathfrak{l}_j} \right\}
\end{aligned}$$

■

**Corollary III.31** *Let  $\ell \geq 1$ ,  $i_1, \dots, i_{\ell+1} \geq 2$ ,  $j \geq \max\{i_1, \dots, i_{\ell+1}\}$  and  $|\kappa_1 - \kappa_2| > 0$ . Define  $I_0 = i_1$ ,  $I_{\ell+1} = i_{\ell+1}$  and, for  $1 \leq m \leq \ell$ ,  $I_m = \max\{i_m, i_{m+1}\}$ . For each  $1 \leq m \leq \ell + 1$ , let  $g_m$  be a sectorized, translation invariant function on  $\mathfrak{Y}_{I_{m-1}, I_m}$  with  $|g_m|_{I_{m-1}, I_m} < \infty$ . Then*

$$\|g_1 \bullet \mathcal{C}^{[I_1, j]} \bullet g_2 \bullet \dots \bullet \mathcal{C}^{[I_\ell, j]} \bullet g_{\ell+1} - g_1 \bullet \mathcal{A}_{I_1, j} \bullet g_2 \bullet \dots \bullet \mathcal{A}_{I_\ell, j} \bullet g_{\ell+1}\|_{\kappa_1, \kappa_2}$$

and

$$\|g_1 \bullet \mathcal{C}^{[I_1, j]} \bullet g_2 \bullet \dots \bullet \mathcal{C}^{[I_\ell, j]} \bullet g_{\ell+1} - g_1 \bullet \mathcal{B}_{I_1, j} \bullet g_2 \bullet \dots \bullet \mathcal{B}_{I_\ell, j} \bullet g_{\ell+1}\|_{\kappa_1, \kappa_2}$$

both converge to zero as  $j \rightarrow \infty$ .

**Proof:** Use Proposition III.30 and the bounds

$$\|g \bullet \mathcal{A}_{i,j} \bullet h\|_{\kappa_1, \kappa_2}, \|g \bullet \mathcal{B}_{i,j} \bullet h\|_{\kappa_1, \kappa_2}, \|g \bullet \mathcal{C}^{i,j} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |j - i + 1| |g|_{\ell, i} \|h\|_{\kappa_1, \kappa_2}$$

which follow from Proposition III.30 and Theorem III.9, with  $\beta = 0$ , to prove that


$$\begin{aligned} & \|g_1 \bullet \mathcal{A}_{I_1, j} \bullet g_2 \bullet \dots \bullet \mathcal{A}_{I_{m-1}, j} \bullet g_m \bullet [\mathcal{C}^{[I_m, j]} - \mathcal{A}_{I_m, j}] \bullet g_{m+1} \bullet \mathcal{C}^{[I_{m+1}, j]} \bullet \dots \bullet \mathcal{C}^{[I_\ell, j]} \bullet g_{\ell+1}\|_{\kappa_1, \kappa_2}, \\ & \|g_1 \bullet \mathcal{B}_{I_1, j} \bullet g_2 \bullet \dots \bullet \mathcal{B}_{I_{m-1}, j} \bullet g_m \bullet [\mathcal{C}^{[I_m, j]} - \mathcal{B}_{I_m, j}] \bullet g_{m+1} \bullet \mathcal{C}^{[I_{m+1}, j]} \bullet \dots \bullet \mathcal{C}^{[I_\ell, j]} \bullet g_{\ell+1}\|_{\kappa_1, \kappa_2} \\ & \leq \text{const}^\ell |j|^\ell \left\{ \sqrt{\mathfrak{l}_j} + \frac{1}{|\kappa_1 - \kappa_2| M^j \mathfrak{l}_j} \right\} \prod_{m=1}^{\ell+1} |g_m|_{I_{m-1}, I_m} \end{aligned}$$

■

## IV. Double Bubbles

In this section we prove the “double bubble bound”, Theorem II.20. The techniques we use are essentially those of §IV with one additional wrinkle – volume improvement due to overlapping loops.

To illustrate the effect of overlapping loops we consider one double bubble, namely



with the kernels of all vertices being identically one in momentum space and all lines having propagator  $C^{(j)}$ . By the Feynman rules

$$D(q, q', t) = \int dk dp \ C^{(j)}(p+q) C^{(j)}(k+p) C^{(j)}(k+t) C^{(j)}(k)$$

The naive power counting bound is, for each  $q, q', t \in \mathbb{R} \times \mathbb{R}^2$ ,

$$\begin{aligned} |D(q, q', t)| &\leq \int dk dp \ |C^{(j)}(p+q)| |C^{(j)}(k+p)| |C^{(j)}(k+t)| |C^{(j)}(k)| \\ &\leq \|C^{(j)}\|_\infty^2 \int dk dp \ |C^{(j)}(k)| |C^{(j)}(p+q)| \\ &= \|C^{(j)}\|_\infty^2 \|C^{(j)}\|_1^2 \\ &\leq \text{const} \end{aligned}$$

since, denoting the  $j^{\text{th}}$  shell by  $S_j$  (see Definition I.2),


$$\begin{aligned} \|C^{(j)}\|_\infty &= \sup_{k \in S_j} \frac{1}{|ik_o - e(\mathbf{k})|} \leq \sqrt{M} M^j \\ \|C^{(j)}\|_1 &\leq \|C^{(j)}\|_\infty (\text{volume of } S_j) \leq \text{const } M^j \frac{1}{M^{2j}} = \frac{\text{const}}{M^j} \end{aligned}$$

In the naive bound, we ignored the constraint that  $|e(\mathbf{k}+\mathbf{p})| \leq \frac{\sqrt{2M}}{M^j}$ . Taking it into account, one has the better estimate

$$\begin{aligned}
|D(q, q', t)| &\leq \int dk dp |C^{(j)}(p+q)| |C^{(j)}(k+p)| |C^{(j)}(k+t)| |C^{(j)}(k)| \\
&\leq \text{const } M^{4j} \int dk dp \nu^{(j)}(p+q) \nu^{(j)}(k+q) \nu^{(j)}(k) \\
&\leq \text{const } M^{4j} \text{vol} \left\{ (k, p) \in (\mathbb{R} \times \mathbb{R}^2)^2 \mid |\imath(p_0 + q_0) - e(\mathbf{p} + \mathbf{q})| \leq \frac{\sqrt{2M}}{M^j}, \right. \\
&\quad \left. |\imath(k_0 + p_0) - e(\mathbf{k} + \mathbf{p})| \leq \frac{\sqrt{2M}}{M^j}, |\imath k_0 - e(\mathbf{k})| \leq \frac{\sqrt{2M}}{M^j} \right\} \\
&\leq \text{const } M^{4j} \frac{2M}{M^{2j}} \text{vol} \left\{ (\mathbf{k}, \mathbf{p}) \in \mathbb{R}^2 \times \mathbb{R}^2 \mid |e(\mathbf{k})|, |e(\mathbf{k} + \mathbf{p})|, |e(\mathbf{p} + \mathbf{q})| \leq \frac{\sqrt{2M}}{M^j} \right\} \\
&\leq \text{const } M^{2j} \int_{|e(\mathbf{p} + \mathbf{q})| \leq \frac{\sqrt{2M}}{M^j}} d\mathbf{p} \text{vol} \left\{ \mathbf{k} \in \mathbb{R}^2 \mid |e(\mathbf{k})|, |e(\mathbf{k} + \mathbf{p})| \leq \frac{\sqrt{2M}}{M^j} \right\}
\end{aligned}$$

There is  $\varepsilon > 0$  such that for  $\mathbf{p}$  outside a ball of radius  $\frac{\text{const}}{M^{j(1-\varepsilon)}}$  around the origin

$$\text{vol} \left( \left\{ \mathbf{k} \in \mathbb{R}^2 \mid |e(\mathbf{k})| \leq \frac{\sqrt{2M}}{M^j} \right\} \cap \left\{ \mathbf{k} \in \mathbb{R}^2 \mid |e(\mathbf{k} + \mathbf{p})| \leq \frac{\sqrt{2M}}{M^j} \right\} \right) \leq \frac{\text{const}}{M^{(1+\varepsilon)j}}$$



because, roughly speaking,  $\{\mathbf{k} \in \mathbb{R}^2 \mid |e(\mathbf{k})| \leq \frac{\sqrt{2M}}{M^j}\}$  and  $\{\mathbf{k} \in \mathbb{R}^2 \mid |e(\mathbf{k} + \mathbf{p})| \leq \frac{\sqrt{2M}}{M^j}\}$  cross at an angle of about  $\text{const } |\mathbf{p}| \geq \frac{\text{const}}{M^{j(1-\varepsilon)}}$ . Therefore

$$\begin{aligned}
\|D\|_\infty &\leq \text{const } M^{2j} \left( \frac{1}{M^{2j(1-\varepsilon)}} \frac{\sqrt{2M}}{M^j} + \frac{\sqrt{2M}}{M^j} \frac{1}{M^{(1+\varepsilon)j}} \right) \\
&\leq \text{const } \frac{1}{M^{\varepsilon j}}
\end{aligned}$$

This “volume improvement” is realized in terms of sector counting in Lemma C.2. Sector counting and simple propagator estimates (Lemma III.14 and Lemma IV.2) are combined using Corollary IV.3 (an analog of Lemma III.11) to prove Theorem IV.4 (which is essentially a reformulation of Theorem II.20 parts b and c in terms of the  $\|\cdot\|_{\kappa_1, \kappa_2}$  norm of Definition III.7) and to treat the large transfer momentum part of the reformulation, Theorem IV.5, of Theorem II.20a (Proposition IV.6). Theorem II.20 parts b and c are proven following Theorem IV.4. The treatment of the small transfer momentum part of Theorem IV.5 closely parallels the corresponding argument in §IV. Theorem II.20a is proven following Theorem IV.5.

We first prove a general bound on  $(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet W \bullet h$  similar to Lemma III.11.

**Lemma IV.1** *Let  $1 \leq \ell \leq i \leq j$  and  $r \geq 1$ . Let  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$  and  $g_1, g_2$  and  $h$  be sectorized, translation invariant functions on  $\mathfrak{Y}_{\ell,\ell}$ ,  $\mathfrak{Y}_{\ell,\ell}$  and  $\mathfrak{Y}_{i,r}$  respectively. Let  $W$  be a particle-hole bubble propagator whose total Fourier transform is of the form*

$$\check{W}(p_1, k_1, p_2, k_2) = \sum_{m=i}^j \sum_{s_1, s_2 \in \Sigma_m} W_{s_1, s_2}^{(m)}(p_1, k_1, p_2, k_2) \quad \text{if } p_2 - k_2 \in \kappa_1 - \kappa_2$$


with  $W_{s_1, s_2}^{(m)}(p_1, k_1, p_2, k_2)$  vanishing unless  $\pi(p_1), \pi(p_2) \in \pi(s_1)$  and  $\pi(k_1), \pi(k_2) \in \pi(s_2)$ . Here  $\pi : k = (k_0, \mathbf{k}) \mapsto \mathbf{k}$  is the projection of  $\mathbb{M} = \mathbb{R} \times \mathbb{R}^2$  onto its second factor. Let  $\mathcal{V}$  be another particle-hole bubble propagator with

$$\check{\mathcal{V}}(p_1, k_1, p_2, k_2) = \sum_{u_1, u_2 \in \Sigma_\ell} \mathcal{V}_{u_1, u_2}(p_1, k_1, p_2, k_2)$$

and  $\mathcal{V}_{u_1, u_2}(p_1, k_1, p_2, k_2) = 0$  unless  $\pi(p_1), \pi(p_2) \in \pi(u_1)$  and  $\pi(k_1), \pi(k_2) \in \pi(u_2)$ . Then

$$\begin{aligned} \|(g_1 \bullet \mathcal{V} \bullet g_2)^f \bullet W \bullet h\|_{\kappa_1, \kappa_2} &\leq 3^8 |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, r} \max_{(u_1, u_2) \in \Sigma_\ell} \|\mathcal{V}_{u_1, u_2}\|_{\text{bubble}} \\ &\sup_{\kappa' \in \mathfrak{K}_\ell} \sum_{m=i}^j \sum_{\substack{s_1, s_2 \in \Sigma_m \\ \pi(s_1 - s_2) \cap \pi(\kappa_1 - \kappa_2) \neq \emptyset}} \inf_{R \in \mathcal{R}(\kappa_1 - \kappa_2)} \|W_{s_1, s_2, R}^{(m)}\|_{\text{bubble}} \\ &\# \{ (u_1, u_2) \in \Sigma_\ell \times \Sigma_\ell \mid \pi(u_1 - u_2) \cap \pi(\kappa' - s_1) \neq \emptyset \} \end{aligned}$$

**Proof:** Consider the case in which all of the external arguments of  $(g_1 \bullet \mathcal{V} \bullet g_2)^f \bullet W \bullet h$  are (position, sector)'s. Set  $\sigma'_1 = \kappa'$  and fix an external sector  $\sigma'_2 \in \Sigma_\ell$ . With the sector names



we have

$$\begin{aligned} &(g_1 \bullet \mathcal{V} \bullet g_2)^f \bullet W \bullet h \\ &= \sum_{\substack{u_1, u_2 \in \Sigma_\ell \\ w_{1,1}, w_{1,2} \in \Sigma_\ell \\ w_{2,1}, w_{2,2} \in \Sigma_\ell}} \sum_{m=i}^j \sum_{\substack{s_1, s_2 \in \Sigma_m \\ v_1, v_2 \in \Sigma_i}} \left( g_1((\cdot, \sigma'_1), (\cdot, w_{1,3}), (\cdot, w_{1,1}), (\cdot, w_{1,2})) \circ \mathcal{V}_{u_1, u_2} \circ g_2((\cdot, w_{2,1}), (\cdot, w_{2,2}), (\cdot, \sigma'_2), (\cdot, w_{2,3})) \right)^f \\ &\quad \circ W_{s_1, s_2}^{(m)} \circ h((\cdot, v_1), (\cdot, v_2), (\cdot, \kappa_1), (\cdot, \kappa_2)) \end{aligned}$$

For each choice of sectors, by conservation of momentum at the vertex  $h$ , we may replace the  $W_{s_1, s_2}^{(m)}$  above by  $W_{s_1, s_2, R}^{(m)}$  with any  $R \in \mathcal{R}(\kappa_1 - \kappa_2)$ . Furthermore the multiple convolution vanishes unless

$$\pi(s_1 - s_2) \cap \pi(\kappa_1 - \kappa_2) \neq \emptyset \quad \pi(u_1 - u_2) \cap \pi(\sigma'_1 - s_1) \neq \emptyset \quad (\text{IV.1})$$

and

$$\begin{aligned} \pi(w_{1,1}) \cap \pi(u_1) &\neq \emptyset & \pi(w_{2,1}) \cap \pi(u_1) &\neq \emptyset & \pi(w_{1,2}) \cap \pi(u_2) &\neq \emptyset & \pi(w_{2,2}) \cap \pi(u_2) &\neq \emptyset \\ \pi(w_{1,3}) \cap \pi(s_1) &\neq \emptyset & \pi(v_1) \cap \pi(s_1) &\neq \emptyset & \pi(w_{2,3}) \cap \pi(s_2) &\neq \emptyset & \pi(v_2) \cap \pi(s_2) &\neq \emptyset \end{aligned} \quad (\text{IV.2})$$

For each fixed  $(u_1, u_2, s_1, s_2)$ , at most  $3^8$  8-tuples  $(w_{1,1}, w_{2,1}, w_{1,2}, w_{2,2}, w_{1,3}, v_1, w_{2,3}, v_2)$  can satisfy (IV.2). Hence

$$\begin{aligned} \|(g_1 \bullet \mathcal{V} \bullet g_2)^f \bullet W \bullet h\|_{\kappa_1, \kappa_2} &\leq 3^8 |h|_{i,r} \max_{\sigma'_1 \in \Sigma_\ell} \sum_{m=i}^j \sum_{\substack{s_1, s_2 \in \Sigma_m \\ \pi(s_1 - s_2) \cap \pi(\kappa_1 - \kappa_2) \neq \emptyset}} \inf_{R \in \mathcal{R}(\kappa_1 - \kappa_2)} \|W_{s_1, s_2, R}^{(m)}\|_{\text{bubble}} \\ &\quad \sum_{\substack{u_1, u_2 \in \Sigma_\ell \\ \pi(u_1 - u_2) \cap \pi(\sigma'_1 - s_1) \neq \emptyset}} |(g_1 \bullet \mathcal{V}_{u_1, u_2} \bullet g_2)^f|_{\ell, \ell} \end{aligned}$$

By definition of the  $\|\cdot\|_{\text{bubble}}$  norm

$$|(g_1 \bullet \mathcal{V}_{u_1, u_2} \bullet g_2)^f|_{\ell, \ell} \leq |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} \|\mathcal{V}_{u_1, u_2}\|_{\text{bubble}}$$

Inserting this gives the Lemma. ■

Recall, for the statement of Theorem II.20, that, for  $\nu \in \mathbb{N}_0 \times \mathbb{N}_0^2$ ,

$$\begin{aligned} \mathcal{D}_{\nu, \text{up}}^{(\ell)}(x_1, x_2, x_3, x_4) &= \frac{1}{M^{|\nu|_\ell}} \sum_{m=\ell}^{\infty} D_{1;3}^\nu C_v^{(\ell)}(x_1, x_3) C_v^{(m)}(x_4, x_2) \\ \mathcal{D}_{\nu, \text{dn}}^{(\ell)}(x_1, x_2, x_3, x_4) &= \frac{1}{M^{|\nu|_\ell}} \sum_{m=\ell+1}^{\infty} C_v^{(m)}(x_1, x_3) D_{2;4}^\nu C_v^{(\ell)}(x_4, x_2) \end{aligned} \quad (\text{IV.3})$$

Express  $\mathcal{D}_{\nu, \text{up}}^{(\ell)} = \sum_{u_1, u_2 \in \Sigma_\ell} \mathcal{D}_{\nu, \text{up}, u_1, u_2}^{(\ell)}$  with

$$\mathcal{D}_{\nu, \text{up}, u_1, u_2}^{(\ell)}(x_1, x_2, x_3, x_4) = \frac{1}{M^{|\nu|_\ell}} \sum_{m \geq \ell} (x_1 - x_3)^\nu c_{u_1}^{(\ell)}(x_3 - x_1) c_{u_2}^{(m)}(x_2 - x_4)$$

where  $c_u^{(n)}(x)$  was defined in (III.9). Do the same for  $\mathcal{D}_{\nu, \text{up}}^{(\ell)}$ .

**Lemma IV.2** *Let  $\ell \geq 1$ ,  $u_1, u_2 \in \Sigma_\ell$  and  $\nu \in \Delta$ . Then*

$$\begin{aligned} \|\mathcal{D}_{\nu, \text{up}, u_1, u_2}^{(\ell)}\|_{\text{bubble}} &\leq \text{const } \mathfrak{l}_\ell \\ \|\mathcal{D}_{\nu, \text{dn}, u_1, u_2}^{(\ell)}\|_{\text{bubble}} &\leq \text{const } \mathfrak{l}_\ell \end{aligned}$$



**Proof:** By the triangle inequality, Lemma III.2 and Lemma A.2,

$$\begin{aligned} \|\mathcal{D}_{\nu,\text{up},u_1,u_2}^{(\ell)}\|_{\text{bubble}} &\leq \frac{1}{M^{|\nu|\ell}} \sum_{m \geq \ell} \|x^\nu c_{u_1}^{(\ell)}\|_{L^1} \|c_{u_2}^{(m)}\|_{L^\infty} \leq \sum_{m \geq \ell} \text{const} \frac{1}{M^{|\nu|\ell}} M^{(1+|\nu|)\ell} \frac{\mathfrak{l}_\ell}{M^m} \\ &\leq \text{const} \mathfrak{l}_\ell \end{aligned}$$

The bound on  $\|\mathcal{D}_{\nu,\text{dn},u_1,u_2}^{(\ell)}\|_{\text{bubble}}$  is proven similarly. ■

**Corollary IV.3** *Let  $1 \leq \ell \leq i \leq j$  and  $r \geq 1$ . Let  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$  and  $g_1, g_2$  and  $h$  be sectorized, translation invariant functions on  $\mathfrak{Y}_{\ell,\ell}$ ,  $\mathfrak{Y}_{\ell,\ell}$  and  $\mathfrak{Y}_{i,r}$  respectively. Let  $W$  be a particle-hole bubble propagator of the form*

$$W(p_1, k_1, p_2, k_2) = \sum_{m=i}^j \sum_{s_1, s_2 \in \Sigma_m} W_{s_1, s_2}^{(m)}(p_1, k_1, p_2, k_2) \quad \text{if } p_2 - k_2 \in \kappa_1 - \kappa_2$$

with  $W_{s_1, s_2}^{(m)}(p_1, k_1, p_2, k_2)$  vanishing unless  $\pi(p_1), \pi(p_2) \in \pi(s_1)$  and  $\pi(k_1), \pi(k_2) \in \pi(s_2)$ . Let  $\mathcal{D}$  be either  $\mathcal{D}_{\nu,\text{up}}^{(\ell)}$  or  $\mathcal{D}_{\nu,\text{dn}}^{(\ell)}$ , with  $\nu \in \Delta$ . Then

$$\begin{aligned} &\|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet W \bullet h\|_{\kappa_1, \kappa_2} \\ &\leq \text{const} |g_1|_{\ell,\ell} |g_2|_{\ell,\ell} |h|_{i,r} \mathfrak{l}_\ell \sup_{\kappa' \in \mathfrak{K}_\ell} \sum_{m=i}^j \sum_{\substack{s_1, s_2 \in \Sigma_m \\ \pi(s_1 - s_2) \cap \pi(\kappa_1 - \kappa_2) \neq \emptyset}} \inf_{R \in \mathcal{R}(\kappa_1 - \kappa_2)} \|W_{s_1, s_2, R}^{(m)}\|_{\text{bubble}} \\ &\quad \# \{ (u_1, u_2) \in \Sigma_\ell \times \Sigma_\ell \mid \pi(u_1 - u_2) \cap \pi(\kappa' - s_1) \neq \emptyset \} \\ &\leq \text{const} |g_1|_{\ell,\ell} |g_2|_{\ell,\ell} |h|_{i,r} \mathfrak{l}_\ell \sum_{m=i}^j \sup_{s_1, s_2 \in \Sigma_m} \|W_{s_1, s_2}^{(m)}\|_{\text{bubble}} \\ &\quad \sup_{\kappa' \in \mathfrak{K}_\ell} \# \left\{ (u_1, u_2, s_1, s_2) \in \Sigma_\ell^2 \times \Sigma_m^2 \mid \begin{array}{l} \pi(u_1 - u_2) \cap \pi(\kappa' - s_1) \neq \emptyset \\ \pi(s_1 - s_2) \cap \pi(\kappa_1 - \kappa_2) \neq \emptyset \end{array} \right\} \end{aligned}$$

**Proof:** The first inequality follows directly from Lemmas IV.1 and IV.2. The second inequality follows by choosing an  $R$  which is one on a large ball. ■

**Theorem IV.4** *Let  $1 \leq \ell \leq i \leq j \leq r$ ,  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$  and let  $g_1, g_2$  and  $h$  be sectorized, translation invariant functions on  $\mathfrak{Y}_{\ell,\ell}$ ,  $\mathfrak{Y}_{\ell,\ell}$  and  $\mathfrak{Y}_{i,r}$  respectively. Let  $\nu \in \Delta$  and  $\mathcal{D}$  be either  $\mathcal{D}_{\nu,\text{up}}^{(\ell)}$  or  $\mathcal{D}_{\nu,\text{dn}}^{(\ell)}$ .*

*i) For any  $\beta \in \Delta$*

$$\begin{aligned} \frac{1}{M^{|\beta|\ell}} \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet D_{1;3}^\beta \mathcal{C}_{\text{top}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} &\leq \text{const} \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell,\ell} |g_2|_{\ell,\ell} |h|_{i,r} \\ \frac{1}{M^{|\beta|\ell}} \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet D_{2;4}^\beta \mathcal{C}_{\text{bot}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} &\leq \text{const} \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell,\ell} |g_2|_{\ell,\ell} |h|_{i,r} \end{aligned}$$

ii)

$$\|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } |j - i + 1| \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, r}$$

and for any  $\beta \in \Delta$  with  $|\beta| \geq 1$  and  $(\mu, \mu') = (1, 3), (2, 4)$

$$\frac{1}{M^{|\beta|j}} \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{D}_{\mu; \mu'}^\beta \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const } \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, r}$$

**Proof:** i) We treat  $\mathcal{C}_{\text{top}}^{[i,j]}$ . The proof for  $\mathcal{C}_{\text{bot}}^{[i,j]}$  is similar. By Corollary IV.3, followed by Lemma III.14 and Lemma C.3

$$\begin{aligned} & \frac{1}{M^{|\beta|j}} \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{D}_{1;3}^\beta \mathcal{C}_{\text{top}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} \\ & \leq \text{const } |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, r} \mathfrak{l}_\ell \sum_{m=i}^j \sup_{s_1, s_2 \in \Sigma_m} \frac{1}{M^{|\beta|j}} \|\mathcal{D}_{1;3}^\beta \mathcal{C}_{\text{top}, j, s_1, s_2}^{(m)}\|_{\text{bubble}} \\ & \quad \sup_{\kappa' \in \mathfrak{K}_\ell} \# \left\{ (u_1, u_2, s_1, s_2) \in \Sigma_\ell^2 \times \Sigma_m^2 \left| \begin{array}{l} \pi(u_1 - u_2) \cap \pi(\kappa' - s_1) \neq \emptyset \\ \pi(s_1 - s_2) \cap \pi(\kappa_1 - \kappa_2) \neq \emptyset \end{array} \right. \right\} \\ & \leq \text{const } |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, r} \mathfrak{l}_\ell \sum_{m=i}^j \frac{1}{M^{|\beta|j}} \mathfrak{l}_m \frac{M^m}{M^j} M^{|\beta|m} \frac{1}{\mathfrak{l}_m \sqrt{\mathfrak{l}_\ell}} \\ & \leq \text{const } |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, r} \sqrt{\mathfrak{l}_\ell} \end{aligned}$$

ii) By Corollary IV.3, followed by Lemma III.14 and Lemma C.3

$$\begin{aligned} & \frac{1}{M^{|\beta|j}} \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{D}_{\mu; \mu'}^\beta \mathcal{C}_{\text{mid}}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} \\ & \leq \text{const } |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, r} \mathfrak{l}_\ell \sum_{m=i}^j \sup_{s_1, s_2 \in \Sigma_m} \frac{1}{M^{|\beta|j}} \|\mathcal{D}_{\mu; \mu'}^\beta \mathcal{C}_{\text{mid}, j, s_1, s_2}^{(m)}\|_{\text{bubble}} \\ & \quad \sup_{\kappa' \in \mathfrak{K}_\ell} \# \left\{ (u_1, u_2, s_1, s_2) \in \Sigma_\ell^2 \times \Sigma_m^2 \left| \begin{array}{l} \pi(u_1 - u_2) \cap \pi(\kappa' - s_1) \neq \emptyset \\ \pi(s_1 - s_2) \cap \pi(\kappa_1 - \kappa_2) \neq \emptyset \end{array} \right. \right\} \\ & \leq \text{const } |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, r} \mathfrak{l}_\ell \sum_{m=i}^j \frac{1}{\mathfrak{l}_m \sqrt{\mathfrak{l}_\ell}} \begin{cases} \mathfrak{l}_m \frac{M^m}{M^j} (j - m + 1) & \beta = 0 \\ \mathfrak{l}_m \frac{M^m}{M^j} & |\beta| = 1 \\ \mathfrak{l}_m \frac{M^m}{M^j} & |\beta| \geq 2 \end{cases} \end{aligned}$$

For  $\beta = 0$ ,  $\sum_{m=i}^j \frac{1}{\mathfrak{l}_m} \mathfrak{l}_m = |j - i + 1|$  as desired. For  $\beta \neq 0$ ,

$$\sum_{m=i}^j \frac{1}{\mathfrak{l}_m} \mathfrak{l}_m \frac{M^m}{M^j} (j - m + 1) = \sum_{m=i}^j M^{-(j-m)} (j - m + 1) \leq \text{const}$$

again, as desired. ■

**Proof of Theorem II.20b,c:** Replacing  $h$  by  $\frac{1}{M^{j|\delta_r|}} D_{3,4}^{\delta_r} h$  reduces consideration to  $\delta_r = 0$ . Suppose that  $\mathcal{D} = \mathcal{D}_{\nu,\text{up}}^{(\ell)}$ . Observe that, by Leibniz (Lemma II.21)

$$\begin{aligned} D_{1,2}^{\delta_1} (g_1 \bullet \mathcal{D} \bullet g_2)^f &= (D_{1,3}^{\delta_1} g_1 \bullet \mathcal{D}_{\nu,\text{up}}^{(\ell)} \bullet g_2)^f \\ &= \sum_{\substack{\beta_1, \beta_2, \beta_3 \in \mathbb{N}_0^3 \\ \beta_1 + \beta_2 + \beta_3 = \delta_1}} \binom{\delta_1}{\beta_1, \beta_2, \beta_3} (D_{1,3}^{\beta_1} g_1 \bullet D_{1,3}^{\beta_2} \mathcal{D}_{\nu,\text{up}}^{(\ell)} \bullet D_{1,3}^{\beta_3} g_2)^f \\ &= \sum_{\substack{\beta_1, \beta_2, \beta_3 \in \mathbb{N}_0^3 \\ \beta_1 + \beta_2 + \beta_3 = \delta_1}} \binom{\delta_1}{\beta_1, \beta_2, \beta_3} M^{|\beta_2 \ell|} (D_{1,3}^{\beta_1} g_1 \bullet \mathcal{D}_{\nu+\beta_2,\text{up}}^{(\ell)} \bullet D_{1,3}^{\beta_3} g_2)^f \end{aligned}$$

Replacing  $\frac{1}{M^{|\beta_1| \ell}} D_{1,3}^{\beta_1} g_1$  by  $g_1$ ,  $\nu + \beta_2$  by  $\nu$  and  $\frac{1}{M^{|\beta_1| \ell}} D_{1,3}^{\beta_3} g_2$  by  $g_2$ , Theorem IV.4, with  $r = j$ , gives bounds on

$$\frac{1}{M^{|\beta|j}} \left\| \frac{1}{M^{|\delta_1| \ell}} M^{|\beta_2 \ell|} (D_{1,3}^{\beta_1} g_1 \bullet \mathcal{D}_{\nu+\beta_2,\text{up}}^{(\ell)} \bullet D_{1,3}^{\beta_3} g_2)^f \bullet D_{\mu;\mu'}^{\beta} \mathcal{C}_{\text{loc}}^{[i,j]} \bullet h \right\|_{\kappa_1, \kappa_2}$$

for each of  $\text{loc} = \text{top, mid, bot}$ . Theorem II.20b,c now follows by Remark III.8. ■

**Theorem IV.5** *Let  $1 \leq \ell \leq i \leq j \leq r$  and  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$ . Set  $d = \kappa_1 - \kappa_2$  and let  $\mathbf{d}$ , the projection of  $d$  onto  $\{0\} \times \mathbb{R}^2$  identified with  $\mathbb{R}^2$ , be contained in a disc of radius  $2\mathfrak{l}_r$  and centre  $\boldsymbol{\tau}$ . Furthermore, set  $\tau_0 = \inf \{ |t_0| \mid (t_0, \mathbf{t}) \in d \text{ for some } \mathbf{t} \in \mathbb{R}^2 \}$ . Assume that*

$$\tau_0 \leq \frac{1}{M^{j-1}} \quad |\boldsymbol{\tau}| \leq \max \left\{ \frac{1}{M^j}, r^3 \mathfrak{l}_r \right\} \quad M^i \leq \mathfrak{l}_j M^j$$

*Also assume that  $p^{(i)}$  vanishes for all  $i > j + 1$ . Let  $\nu \in \mathbb{N}_0 \times \mathbb{N}_0^2$ , with  $\nu + \alpha \in \Delta$  for all  $|\alpha| \leq 3$  and let  $\mathcal{D}$  be either  $\mathcal{D}_{\nu,\text{up}}^{(\ell)}$  or  $\mathcal{D}_{\nu,\text{dn}}^{(\ell)}$ . For any sectorized, translation invariant functions  $g_1, g_2$  and  $h$  on  $\mathfrak{Y}_{\ell,\ell}, \mathfrak{Y}_{\ell,\ell}$  and  $\mathfrak{Y}_{i,r}$  respectively,*

$$\left\| (g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{C}^{[i,j]} \bullet h \right\|_{\kappa_1, \kappa_2} \leq \text{const} \sqrt{\mathfrak{l}_\ell} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} |g_1|_\ell^{[\alpha_{\text{up}}]} |g_2|_\ell^{[\alpha_{\text{dn}}]} |h|_{i,r}^{[\alpha_1, 0, 0]}$$

Theorem IV.5 is proven at the end of this section.

**Proof of Theorem II.20a** (assuming Theorem IV.5):

As in the proof of Theorem II.20b,c, we may assume without loss of generality that  $\delta_1 = \delta_r = 0$ . Fix  $1 \leq \ell \leq i \leq j$ ,  $\nu \in \mathbb{N}_0 \times \mathbb{N}_0^2$ ,  $\mathcal{D}$  and sectorized, translation invariant functions  $g_1, g_2$  and  $h$  on  $\mathfrak{Y}_{\ell,\ell}, \mathfrak{Y}_{\ell,\ell}$  and  $\mathfrak{Y}_{i,j}$  as in Theorem II.20. By Remark III.8, it suffices to prove that

$$\left\| (g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{C}^{[i,j]} \bullet h \right\|_{\kappa_1, \kappa_2} \leq \text{const} \ i \sqrt{\mathfrak{l}_\ell} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} |g_1|_\ell^{[\alpha_{\text{up}}]} |g_2|_\ell^{[\alpha_{\text{dn}}]} |h|_{i,j}^{[\alpha_1, 0, 0]} \quad (\text{IV.4})$$

for all  $\kappa_1, \kappa_2 \in \mathfrak{K}_j$ . Fix  $\kappa_1, \kappa_2 \in \mathfrak{K}_j$ . Set  $d = \kappa_1 - \kappa_2$  and denote by  $\mathbf{d}$  the projection of  $d$  onto  $\{0\} \times \mathbb{R}^2$  identified with  $\mathbb{R}^2$ . By Remark III.12, the set  $\mathbf{d}$  is contained in a disc of radius  $2\mathfrak{l}_j$ . We fix such a disk and denote by  $\boldsymbol{\tau}$  its centre. Furthermore, as in the proof of Theorem II.19a, we define  $\tau_0 = \inf \{ |t_0| \mid (t_0, \mathbf{t}) \in d \text{ for some } \mathbf{t} \in \mathbb{R}^2 \}$  and

$$\begin{aligned} j_0 &= \begin{cases} \max \{ n \in \mathbb{N}_0 \mid \tau_0 \leq \frac{1}{M^{n-1}} \} & \text{if } 0 < \tau_0 \leq M \\ 0 & \text{if } \tau_0 \geq M \\ \infty & \text{if } \tau_0 = 0 \end{cases} \\ j_1 &= \begin{cases} \max \{ n \in \mathbb{N}_0 \mid |\boldsymbol{\tau}| \leq \frac{1}{M^n} \} & \text{if } j^3 \mathfrak{l}_j < |\boldsymbol{\tau}| \leq 1 \\ 0 & \text{if } |\boldsymbol{\tau}| \geq 1 \\ \infty & \text{if } |\boldsymbol{\tau}| \leq j^3 \mathfrak{l}_j \end{cases} \\ \bar{j} &= \max \{ i - 1, \min \{ j, j_0, j_1 \} \} \end{aligned}$$

The analog of Proposition III.16 in the current double bubble setting is

**Proposition IV.6 (Large Transfer Momentum)**

$$\| (g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{C}^{[\bar{j}+1, j]} \bullet h \|_{\kappa_1, \kappa_2} \leq \text{const } \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, j}$$

**Proof:** If  $\min \{ j, j_0, j_1 \} = j$ , then  $\bar{j} = j$  and  $\mathcal{C}^{[\bar{j}+1, j]} = 0$  so that there is nothing to prove. So we may assume that  $\min \{ j_0, j_1 \} < j$ .

*Case 1:*  $j_0 \leq j_1$ . In this case,  $\| (g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{C}^{[\bar{j}+1, j]} \bullet h \|_{\kappa_1, \kappa_2} = 0$ , because  $\mathcal{C}^{[\bar{j}+1, j]}(p, k)$  vanishes unless  $|p_0|, |k_0| \leq \frac{\sqrt{2M}}{M^{\bar{j}+1}}$  and hence unless  $|p_0 - k_0| \leq \frac{2\sqrt{2M}}{M^{\bar{j}+1}} < \frac{1}{M^{\bar{j}}} < \tau_0$ , while  $|t_0| \geq \tau_0$  for all  $t \in d$ .

*Case 2:*  $j_1 < j_0$ . In this case  $|\boldsymbol{\tau}| \geq j^3 \mathfrak{l}_j$ . By Corollary IV.3, Lemma III.14 and Lemma C.4

$$\begin{aligned} & \| (g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{C}^{[\bar{j}+1, j]} \bullet h \|_{\kappa_1, \kappa_2} \\ & \leq \text{const } |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, j} \mathfrak{l}_\ell \sum_{m=\bar{j}+1}^j \sup_{s_1, s_2 \in \Sigma_m} \| \mathcal{C}_{s_1, s_2}^{(m)} \|_{\text{bubble}} \\ & \quad \sup_{\kappa' \in \mathfrak{K}_\ell} \# \left\{ (u_1, u_2, s_1, s_2) \in \Sigma_\ell^2 \times \Sigma_m^2 \mid \begin{array}{l} \pi(u_1 - u_2) \cap \pi(\kappa' - s_1) \neq \emptyset \\ \pi(s_1 - s_2) \cap \pi(\kappa_1 - \kappa_2) \neq \emptyset \end{array} \right\} \\ & \leq \text{const } |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, j} \sum_{m=\bar{j}+1}^j \left[ \min \left( \ell \mathfrak{l}_\ell, \frac{1+M^m \mathfrak{l}_j}{M^m |\boldsymbol{\tau}|} \right) + \frac{\sqrt{\mathfrak{l}_\ell}}{|\boldsymbol{\tau}|} \left( \frac{1}{M^m} + \mathfrak{l}_j \right) + \sqrt{\mathfrak{l}_m} \right] \end{aligned}$$

The last sum

$$\sum_{m=\bar{j}+1}^j \sqrt{\mathfrak{l}_m} \leq \sqrt{\mathfrak{l}_{\bar{j}}} \leq \sqrt{\mathfrak{l}_{i-1}} \leq \text{const } \sqrt{\mathfrak{l}_\ell}$$

As, by the definition of  $j_1$ ,

$$\frac{1}{|\mathcal{T}|} \sum_{m=j_1+1}^j \left( \frac{1}{M^m} + \mathfrak{l}_j \right) \leq \frac{1}{|\mathcal{T}|} \left( \frac{1}{M^{j_1}} + j \mathfrak{l}_j \right) \leq \text{const}$$

the middle sum

$$\sum_{m=\bar{j}+1}^j \frac{\sqrt{\mathfrak{l}_\ell}}{|\mathcal{T}|} \left( \frac{1}{M^m} + \mathfrak{l}_j \right) \leq \text{const} \sqrt{\mathfrak{l}_\ell}$$

As  $\min \left( \ell \mathfrak{l}_\ell, \frac{1+M^m \mathfrak{l}_j}{M^m |\mathcal{T}|} \right) \leq \min \left( \ell \mathfrak{l}_\ell, \frac{1}{M^m |\mathcal{T}|} \right) + \min \left( \ell \mathfrak{l}_\ell, \frac{\mathfrak{l}_j}{|\mathcal{T}|} \right)$ , the first sum

$$\begin{aligned} \sum_{m=\bar{j}+1}^j \min \left( \ell \mathfrak{l}_\ell, \frac{1+M^m \mathfrak{l}_j}{M^m |\mathcal{T}|} \right) &\leq j \min \left( \ell \mathfrak{l}_\ell, \frac{\mathfrak{l}_j}{|\mathcal{T}|} \right) + \sum_{m=\bar{j}+1}^j \min \left( \ell \mathfrak{l}_\ell, \frac{1}{M^m |\mathcal{T}|} \right) \\ &\leq j \min \left( \ell \mathfrak{l}_\ell, \frac{\mathfrak{l}_j}{|\mathcal{T}|} \right) + \sum_{m=\bar{j}+1}^j (\ell \mathfrak{l}_\ell)^{2/3} \frac{1}{(M^m |\mathcal{T}|)^{1/3}} \\ &\leq j \min \left( \ell \mathfrak{l}_\ell, \frac{\mathfrak{l}_j}{|\mathcal{T}|} \right) + (\ell \mathfrak{l}_\ell)^{2/3} \frac{1}{(M^{\bar{j}} |\mathcal{T}|)^{1/3}} \\ &\leq j \min \left( \ell \mathfrak{l}_\ell, \frac{\mathfrak{l}_j}{|\mathcal{T}|} \right) + \text{const} (\ell \mathfrak{l}_\ell)^{2/3} \end{aligned}$$

If  $j \leq \frac{1}{\mathfrak{l}_\ell^{1/3}}$ , then

$$j \ell \mathfrak{l}_\ell \leq \ell \mathfrak{l}_\ell^{2/3} \leq \text{const} \sqrt{\mathfrak{l}_\ell}$$

while, if  $j \geq \frac{1}{\mathfrak{l}_\ell^{1/3}}$ , then

$$\frac{j \mathfrak{l}_j}{|\mathcal{T}|} \leq \frac{1}{j^2} \leq \mathfrak{l}_\ell^{2/3}$$

■

**Continuation of the proof of Theorem II.20a** (assuming Theorem IV.5):

When  $M^i \geq \mathfrak{l}_{\bar{j}} M^{\bar{j}} = M^{(1-\aleph)\bar{j}}$ , we have  $|\bar{j} - i + 1| \leq \text{const } i$ . In this case Theorem IV.4, with  $r = j$  and  $j = \bar{j}$ , gives

$$\left\| (g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{C}^{[i, \bar{j}]} \bullet h \right\|_{\kappa_1, \kappa_2} \leq \text{const } i \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, j}$$

This together with Proposition IV.6 yields (IV.4). Therefore, we may assume that

$$M^i \leq \mathfrak{l}_{\bar{j}} M^{\bar{j}} \tag{IV.5}$$

Furthermore, if  $\bar{j} = i - 1$ ,  $\mathcal{C}^{[i, \bar{j}]} = 0$  and there is nothing more to prove. So we may also assume that  $j_0, j_1 \geq i$  and  $\bar{j} \leq j, j_0, j_1$ .

Set  $v' = \sum_{i=2}^{\bar{j}+1} p^{(i)}$  and  $\mathcal{C}'^{[i,\bar{j}]} = \mathcal{C}'_{\text{top}}^{[i,\bar{j}]} + \mathcal{C}'_{\text{mid}}^{[i,\bar{j}]} + \mathcal{C}'_{\text{bot}}^{[i,\bar{j}]}$  with

$$\mathcal{C}'_{\text{top}}^{[i,\bar{j}]} = \sum_{\substack{i \leq i_t \leq \bar{j} \\ i_b > \bar{j}}} C_{v'}^{(i_t)} \otimes C_{v'}^{(i_b) t}, \quad \mathcal{C}'_{\text{mid}}^{[i,\bar{j}]} = \sum_{\substack{i \leq i_t \leq \bar{j} \\ i \leq i_b \leq \bar{j}}} C_{v'}^{(i_t)} \otimes C_{v'}^{(i_b) t}, \quad \mathcal{C}'_{\text{bot}}^{[i,\bar{j}]} = \sum_{\substack{i_t > \bar{j} \\ i \leq i_b \leq \bar{j}}} C_{v'}^{(i_t)} \otimes C_{v'}^{(i_b) t}$$

as in §III. Again,  $v-v'$  is supported on the  $(\bar{j}+2)^{\text{nd}}$  extended neighbourhood and  $\mathcal{C}_{\text{mid}}^{[i,\bar{j}]} = \mathcal{C}'_{\text{mid}}^{[i,\bar{j}]}$ . Hence, by Theorem IV.4.i, with  $\beta = 0$ ,  $r = j$  and  $j = \bar{j}$ ,

$$\|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet [\mathcal{C}^{[i,\bar{j}]} - \mathcal{C}'^{[i,\bar{j}]}] \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const} \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, j} \quad (\text{IV.6})$$

By (IV.5) and the Definitions of  $\bar{j}$  and  $\mathcal{C}'^{[i,\bar{j}]}$ , the hypotheses of Theorem IV.5, with  $r = j$  and  $j = \bar{j}$ , apply to  $(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{C}'^{[i,\bar{j}]} \bullet h$ . Hence

$$\|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{C}'^{[i,\bar{j}]} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const} \sqrt{\mathfrak{l}_\ell} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} |g_1|_{\ell}^{[\alpha_{\text{up}}]} |g_2|_{\ell}^{[\alpha_{\text{dn}}]} |h|_{i, j}^{[\alpha_1, 0, 0]}$$

This together with (IV.6) and Proposition IV.6 yields (IV.4). This completes the proof that Theorem IV.5 implies Theorem II.20.a. ■

The rest of this section is devoted to the proof of Theorem IV.5. So we fix  $\nu \in \Delta$ ,  $1 \leq \ell \leq i \leq j \leq r$ ,  $\mathcal{D} = \mathcal{D}_{\nu, \text{up}}^{(\ell)}$  or  $\mathcal{D}_{\nu, \text{dn}}^{(\ell)}$  and sectorized, translation invariant functions,  $g_1$ ,  $g_2$  and  $h$ , on  $\mathfrak{Y}_{\ell, \ell}$ ,  $\mathfrak{Y}_{\ell, \ell}$  and  $\mathfrak{Y}_{i, r}$  respectively. We also fix  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$  and assume that

$$\tau_0 \leq \frac{1}{M^{\bar{j}-1}} \quad |\tau| \leq \max \left\{ \frac{1}{M^{\bar{j}}}, r^3 \mathfrak{l}_r \right\} \quad M^i \leq \mathfrak{l}_j M^{\bar{j}} \quad (\text{IV.7})$$

and that  $p^{(i)}$  vanishes for all  $i > j + 1$ . As in §III, we reduce the particle-hole bubble propagator  $\mathcal{C}^{[i, j]}$  to the model bubble propagator of (III.22). This is done in the following two lemmata.

**Lemma IV.7** *Let  $\mathcal{Z}$  be the operator defined in (III.18). Then*

$$\|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet (\mathcal{Z} \bullet \mathcal{C}^{[i, j]} \bullet \mathcal{Z}^t - \mathcal{M}) \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const} \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, r}$$

**Proof:** Expand

$$\mathcal{Z} \bullet \mathcal{C}^{[i, j]} \bullet \mathcal{Z}^t - \mathcal{M} = (\tilde{\mathcal{C}}^{[i, j]} - \mathcal{M}) + (\mathcal{Z} \bullet \mathcal{C}^{[i, j]} \bullet \mathcal{Z}^t - \tilde{\mathcal{C}}^{[i, j]})$$

where  $\tilde{\mathcal{C}}^{[i,j]}$  was defined in (III.20). Then, for  $|p_0| \leq \frac{M+4\sqrt{2M}}{M^j}$ ,

$$(\tilde{\mathcal{C}}^{[i,j]} - \mathcal{M}) + (\mathcal{Z} \bullet \mathcal{C}^{[i,j]} \bullet \mathcal{Z}^t - \tilde{\mathcal{C}}^{[i,j]}) = \sum_{m=i}^j \sum_{s_1, s_2 \in \Sigma_m} \mathcal{D}_{s_1, s_2}^{(m)} + \sum_{m=i}^j \sum_{s_1, s_2 \in \Sigma_m} (\mathcal{Z} \bullet \mathcal{C}_{s_1, s_2}^{(m)} \bullet \mathcal{Z}^t - \tilde{\mathcal{C}}_{s_1, s_2}^{(m)})$$

where  $\mathcal{D}_{s_1, s_2}^{(m)}$  was defined in (III.23). So, by Corollary IV.3,

$$\begin{aligned} & \| (g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet (\tilde{\mathcal{C}}^{[i,j]} - \mathcal{M} + \mathcal{Z} \bullet \mathcal{C}^{[i,j]} \bullet \mathcal{Z}^t - \tilde{\mathcal{C}}^{[i,j]}) \bullet h \|_{\kappa_1, \kappa_2} \\ & \leq \text{const} |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, r} \mathfrak{l}_\ell \sum_{m=i}^j \\ & \quad \sup_{\kappa' \in \mathfrak{K}_\ell} \# \{ (s_1, s_2, u_1, u_2) \mid \pi(s_1 - s_2) \cap \pi(\kappa_1 - \kappa_2) \neq \emptyset, \pi(u_1 - u_2) \cap \pi(\kappa' - s_1) \neq \emptyset \} \\ & \quad \max_{s_1, s_2 \in \Sigma_m} \left[ \|\mathcal{D}_{s_1, s_2}^{(m)}\|_{\text{bubble}} + \|\mathcal{Z} \bullet \mathcal{C}_{s_1, s_2, \phi}^{(m)} \bullet \mathcal{Z}^t - (\tilde{\mathcal{C}}_{s_1, s_2}^{(m)})_\phi\|_{\text{bubble}} \right] \end{aligned}$$

where  $\phi$  was defined at the beginning of the proof of Proposition III.22. Then by Lemma C.3, Lemma III.25 and Lemma III.23

$$\begin{aligned} & \| (g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet (\tilde{\mathcal{C}}^{[i,j]} - \mathcal{M} + \mathcal{Z} \bullet \mathcal{C}^{[i,j]} \bullet \mathcal{Z}^t - \tilde{\mathcal{C}}^{[i,j]}) \bullet h \|_{\kappa_1, \kappa_2} \\ & \leq \text{const} |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, r} \mathfrak{l}_\ell \left[ \sum_{m=i}^{i+1} \frac{1}{\mathfrak{l}_m \sqrt{\mathfrak{l}_\ell}} \mathfrak{l}_m + \sum_{m=i+2}^j \frac{j-m+1}{\mathfrak{l}_m \sqrt{\mathfrak{l}_\ell}} \frac{M^m}{M^j} \mathfrak{l}_m + \sum_{m=i}^j \frac{1}{\mathfrak{l}_m \sqrt{\mathfrak{l}_\ell}} \mathfrak{l}_m \frac{M^m}{M^j} \right] \\ & \leq \text{const} \sqrt{\mathfrak{l}_\ell} |g_1|_{\ell, \ell} |g_2|_{\ell, \ell} |h|_{i, r} \end{aligned}$$

■

## Lemma IV.8

$$\| (g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet (\mathcal{C}^{[i,j]} - \mathcal{Z} \bullet \mathcal{C}^{[i,j]} \bullet \mathcal{Z}^t) \bullet h \|_{\kappa_1, \kappa_2} \leq \text{const} \sqrt{\mathfrak{l}_\ell} \max_{|\alpha_{\text{up}}| + |\alpha_{\text{dn}}| \leq 1} |g_1|_\ell^{\llbracket \alpha_{\text{up}} \rrbracket} |g_2|_\ell^{\llbracket \alpha_{\text{dn}} \rrbracket} |h|_{i, r}^{[1, 0, 0]}$$

**Proof:** By Lemma III.20

$$\begin{aligned} & (g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet (\mathcal{C}^{[i,j]} - \mathcal{Z} \bullet \mathcal{C}^{[i,j]} \bullet \mathcal{Z}^t) \bullet h \\ & = (g_1 \bullet \mathcal{D} \bullet g_2)_r^f \bullet D_1 \mathcal{C}^{[i,j]} \bullet h + (g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{Z} \bullet D_r \mathcal{C}^{[i,j]} \bullet h_l \end{aligned}$$

Both terms are bounded as in the previous lemma, using Lemma C.3 to bound the number of allowed 4-tuples  $(s_1, s_2, u_1, u_2)$  by  $\frac{1}{\mathfrak{l}_m \sqrt{\mathfrak{l}_\ell}}$ . Lemma III.21, (and, for the second term, Lemma III.18.iii) are used to bound  $\|D_1 \mathcal{C}_{s_1, s_2}^{(m)}\|_{\text{bubble}} \leq \text{const} \frac{\mathfrak{l}_m}{M^m}$  and  $\|\mathcal{Z} \bullet D_r \mathcal{C}_{s_1, s_2}^{(m)}\|_{\text{bubble}} \leq$

$\|D_r \mathcal{C}_{s_1, s_2}^{(m)}\|_{\text{bubble}} \leq \text{const } \frac{l_m}{M^m}$ . The right derivative in  $(g_1 \bullet \mathcal{D} \bullet g_2)_r^f$  acts as a central derivative on  $g_1 \bullet \mathcal{D} \bullet g_2$  and may be written, using Leibniz's rule, as a sum of three terms with the first containing a central derivative acting on  $g_1$ , the second a central derivative acting on  $g_2$  and the third having one component of  $\mathcal{D}$ 's index  $\nu$  increased by one. Lemma IV.2 is used to bound the bubble norms of the sectorized contributions to  $\mathcal{D}$ . All together,

$$\begin{aligned}
& \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet (\mathcal{C}^{[i,j]} - \mathcal{Z} \bullet \mathcal{C}^{[i,j]} \bullet \mathcal{Z}^t) \bullet h\|_{\kappa_1, \kappa_2} \\
& \leq \|(g_1 \bullet \mathcal{D} \bullet g_2)_r^f \bullet D_1 \mathcal{C}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} + \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{Z} \bullet D_r \mathcal{C}^{[i,j]} \bullet h\|_{\kappa_1, \kappa_2} \\
& \leq \text{const} \left[ M^\ell |g_1|_\ell^{[(1,0,0)]} \mathfrak{l}_\ell |g_2|_{\ell, \ell} + |g_1|_{\ell, \ell} M^\ell \mathfrak{l}_\ell |g_2|_{\ell, \ell} + |g_1|_{\ell, \ell} \mathfrak{l}_\ell M^\ell |g_2|_\ell^{[(1,0,0)]} \right] \\
& \quad |h|_{i,r} \sum_{m=i}^j \frac{1}{l_m \sqrt{l_\ell}} \frac{l_m}{M^m} \\
& \quad + \text{const} |g_1|_{\ell, \ell} \mathfrak{l}_\ell |g_2|_{\ell, \ell} |h|_{i,r} \sum_{m=i}^j \frac{1}{l_m \sqrt{l_\ell}} \frac{l_m}{M^m} \\
& \leq \text{const} \sqrt{l_\ell} \max_{|\alpha_{\text{up}}| + |\alpha_{\text{dn}}| \leq 1} |g_1|_\ell^{[\alpha_{\text{up}}]} |g_2|_\ell^{[\alpha_{\text{dn}}]} |h|_{i,r}^{[1,0,0]} \sum_{m=i}^j \frac{M^i}{M^m} \\
& \leq \text{const} \sqrt{l_\ell} \max_{|\alpha_{\text{up}}| + |\alpha_{\text{dn}}| \leq 1} |g_1|_\ell^{[\alpha_{\text{up}}]} |g_2|_\ell^{[\alpha_{\text{dn}}]} |h|_{i,r}^{[1,0,0]}
\end{aligned}$$

■

### Proposition IV.9

$$\|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{M} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const} \sqrt{l_\ell} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} |g_1|_\ell^{[\alpha_{\text{up}}]} |g_2|_\ell^{[\alpha_{\text{dn}}]} |h|_{i,r}^{[\alpha_1, 0, 0]}$$

**Proof:** Write

$$\mathcal{M}(p, k) = \sum_{s_1, s_2 \in \Sigma_i} \mathcal{M}_{s_1, s_2}(p, k)$$

where

$$\mathcal{M}_{s_1, s_2}(p, k) = \mathcal{M}(p, k) \rho(\mathbf{p} - \mathbf{k}) \chi_{s_1}(\mathbf{p}) \chi_{s_2}(\mathbf{k})$$


and  $\rho$  was defined just before (III.24). Then

$$\begin{aligned}
& (g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{M} \bullet h \\
& = \sum_{\substack{u_1, u_2 \in \Sigma_\ell \\ w_{1,1}, w_{1,2} \in \Sigma_\ell \\ w_{2,1}, w_{2,2} \in \Sigma_\ell}} \sum_{\substack{w_{1,3}, w_{2,3} \in \Sigma_\ell \\ s_1, s_2 \in \Sigma_i \\ v_1, v_2 \in \Sigma_i}} \\
& \quad \left( g_1(y_1, (\cdot, w_{1,3}), (\cdot, w_{1,1}), (\cdot, w_{1,2})) \circ \mathcal{D}_{u_1, u_2} \circ g_2((\cdot, w_{2,1}), (\cdot, w_{2,2}), y_2, (\cdot, w_{2,3})) \right)^f \\
& \quad \circ \mathcal{M}_{s_1, s_2} \circ h((\cdot, v_1), (\cdot, v_2), y_3, y_4)
\end{aligned}$$



where, for  $\nu = 1, 2$ ,

$$y_\nu = \begin{cases} \kappa'_\nu & \text{if } \kappa'_\nu \in \mathbb{M} \\ (x_\nu, \kappa'_\nu) & \text{if } \kappa'_\nu \in \Sigma_\ell \end{cases} \quad y_{\nu+2} = \begin{cases} \kappa_\nu & \text{if } \kappa_\nu \in \mathbb{M} \\ (x_{\nu+2}, \kappa_\nu) & \text{if } \kappa_\nu \in \Sigma_r \end{cases}$$



The multiple convolution vanishes unless

$$\pi(s_1 - s_2) \cap \pi(\kappa_1 - \kappa_2) \neq \emptyset \quad \pi(u_1 - u_2) \cap \pi(\kappa'_1 - s_1) \neq \emptyset$$

and

$$\begin{aligned} w_{1,1} \cap u_1 \neq \emptyset \quad w_{2,1} \cap u_1 \neq \emptyset \quad w_{1,2} \cap u_2 \neq \emptyset \quad w_{2,2} \cap u_2 \neq \emptyset \\ \pi(w_{1,3}) \cap \pi(s_1) \neq \emptyset \quad \pi(v_1) \cap \pi(s_1) \neq \emptyset \quad \pi(w_{2,3}) \cap \pi(s_2) \neq \emptyset \quad \pi(v_2) \cap \pi(s_2) \neq \emptyset \end{aligned} \quad (\text{IV.8})$$

Fix the external sectors/momenta  $(\kappa_1, \kappa_2, \kappa'_1, \kappa'_2)$ . Then, for each fixed  $(s_1, s_2, u_1, u_2)$ , there are at most  $3^8$  8-tuples  $(w_{1,1}, w_{2,1}, w_{1,2}, w_{2,2}, w_{1,3}, v_1, w_{2,3}, v_2)$  satisfying (IV.8). By Lemma C.3, the number of allowed 4-tuples  $(s_1, s_2, u_1, u_2)$  is bounded by  $\frac{1}{l_i \sqrt{l_\ell}}$ . Set, for each  $\varsigma = (\kappa'_1, \kappa'_2, w_{1,1}, w_{1,2}, w_{1,3}, w_{2,1}, w_{2,2}, w_{2,3}, u_1, u_2) \in \mathfrak{K}_\ell^2 \times \Sigma_\ell^8$

$$g'_\varsigma = g_1(y_1, (\cdot, w_{1,3}), (\cdot, w_{1,1}), (\cdot, w_{1,2})) \circ \mathcal{D}_{u_1, u_2} \circ g_2((\cdot, w_{2,1}), (\cdot, w_{2,2}), y_2, (\cdot, w_{2,3}))$$

for each  $\tau = (w_{1,1}, w_{1,2}, w_{2,1}, w_{2,2}, u_1, u_2) \in \Sigma_\ell^6$ ,

$$g_\tau = g_1(\cdot, \cdot, (\cdot, w_{1,1}), (\cdot, w_{1,2})) \circ \mathcal{D}_{u_1, u_2} \circ g_2((\cdot, w_{2,1}), (\cdot, w_{2,2}), \cdot, \cdot)$$

and, for each  $v_1, v_2 \in \Sigma_i$ ,

$$h_{v_1, v_2}(x_1, x_2, y_3, y_4) = h((x_1, v_1), (x_2, v_2), y_3, y_4)$$

Then

$$\|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{M} \bullet h\|_{\kappa_1, \kappa_2} \leq \text{const} \frac{1}{l_i \sqrt{l_\ell}} \sup_{\varsigma \in \mathfrak{K}_\ell^2 \times \Sigma_\ell^8} \max_{\substack{s_1, s_2 \in \Sigma_i \\ v_1, v_2 \in \Sigma_i}} \|g'_\varsigma \circ \mathcal{M}_{s_1, s_2} \circ h_{v_1, v_2}\|_{1, \infty}$$

By (III.28)

$$\|g'_\varsigma{}^f \circ \mathcal{M}_{s_1, s_2} \circ h_{v_1, v_2}\|_{1, \infty} \leq \text{const } \mathfrak{l}_i \max_{\tau \in \Sigma_\ell^6} \max_{\substack{\alpha_r, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_r| + |\alpha_1| \leq 3}} |g_\tau^f|_{\ell, i}^{[0, 0, \alpha_r]} |h|_{i, r}^{[\alpha_1, 0, 0]}$$

Bounding

$$\begin{aligned} |g_\tau^f|_{\ell, i}^{[0, 0, \alpha_r]} &\leq |g_\tau|_{\ell, \ell}^{[0, \alpha_r, 0]} \\ &\leq \text{const } \mathfrak{l}_\ell \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}} \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \alpha_{\text{up}} + \alpha_{\text{dn}} \leq \alpha_r}} |g_1|_\ell^{\llbracket \alpha_{\text{up}} \rrbracket} |g_2|_\ell^{\llbracket \alpha_{\text{dn}} \rrbracket} \end{aligned}$$

by Leibniz and Lemma IV.2, yields

$$\begin{aligned} \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{M} \bullet h\|_{\kappa_1, \kappa_2} &\leq \text{const } \frac{1}{\mathfrak{l}_i \sqrt{\mathfrak{l}_\ell}} \mathfrak{l}_i \mathfrak{l}_\ell \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} |g_1|_\ell^{\llbracket \alpha_{\text{up}} \rrbracket} |g_2|_\ell^{\llbracket \alpha_{\text{dn}} \rrbracket} |h|_{i, r}^{[\alpha_1, 0, 0]} \\ &\leq \text{const } \sqrt{\mathfrak{l}_\ell} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} |g_1|_\ell^{\llbracket \alpha_{\text{up}} \rrbracket} |g_2|_\ell^{\llbracket \alpha_{\text{dn}} \rrbracket} |h|_{i, r}^{[\alpha_1, 0, 0]} \end{aligned}$$

■

**Proof of Theorem IV.5:** By Lemmas IV.7, IV.8 and Proposition IV.9,

$$\begin{aligned} \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{C}^{[i, j]} \bullet h\|_{\kappa_1, \kappa_2} &\leq \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet (\mathcal{Z} \bullet \mathcal{C}^{[i, j]} \bullet \mathcal{Z}^t - \mathcal{M}) \bullet h\|_{\kappa_1, \kappa_2} \\ &\quad + \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet (\mathcal{C}^{[i, j]} - \mathcal{Z} \bullet \mathcal{C}^{[i, j]} \bullet \mathcal{Z}^t) \bullet h\|_{\kappa_1, \kappa_2} \\ &\quad + \|(g_1 \bullet \mathcal{D} \bullet g_2)^f \bullet \mathcal{M} \bullet h\|_{\kappa_1, \kappa_2} \\ &\leq \text{const } \sqrt{\mathfrak{l}_\ell} \max_{\substack{\alpha_{\text{up}}, \alpha_{\text{dn}}, \alpha_1 \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ |\alpha_{\text{up}}| + |\alpha_{\text{dn}}| + |\alpha_1| \leq 3}} |g_1|_\ell^{\llbracket \alpha_{\text{up}} \rrbracket} |g_2|_\ell^{\llbracket \alpha_{\text{dn}} \rrbracket} |h|_{i, r}^{[\alpha_1, 0, 0]} \end{aligned}$$

as desired. ■

## Appendix A: Bounds on Propagators

Fix, as in Theorem I.20, a sequence,  $p^{(2)}, p^{(3)}, \dots$ , of sectorized, translation invariant functions  $p^{(i)}$  on  $\left((\mathbb{R} \times \mathbb{R}^2) \times \Sigma_i\right)^2$  obeying

$$|p^{(i)}|_{1, \Sigma_i} \leq \frac{\rho \mathfrak{l}_i}{M^i} \mathfrak{c}_i \quad \check{p}^{(i)}(0, \mathbf{k}) = 0$$

and set, for all  $j \geq 1$ ,

$$\begin{aligned} v(k) &= \sum_{i=2}^{\infty} \check{p}^{(i)}(k) & e'(k) &= e(\mathbf{k}) - v(k) \\ v_j(k) &= \sum_{i=2}^j \check{p}^{(i)}(k) & e'_j(k) &= e(\mathbf{k}) - v_j(k) \end{aligned}$$

**Lemma A.1** *There is a  $\tilde{\rho}_0 > 0$  such that for all  $\rho \leq \tilde{\rho}_0$  and  $(k_0, \mathbf{k})$  in the first neighbourhood*

*i)*

$$\begin{aligned} \nabla_{\mathbf{k}} e'(k_0, \mathbf{k}) &\neq 0 & \nabla_{\mathbf{k}} e'_j(k_0, \mathbf{k}) &\neq 0 \\ |\partial_{k_0} e'(k_0, \mathbf{k})| &\leq \rho < \frac{1}{2} & |\partial_{k_0} e'_j(k_0, \mathbf{k})| &\leq \rho < \frac{1}{2} \\ |\imath k_0 - e'(k)| &\geq \frac{1}{2} |\imath k_0 - e(\mathbf{k})| & |\imath k_0 - e'_j(k)| &\geq \frac{1}{2} |\imath k_0 - e(\mathbf{k})| \end{aligned}$$

*ii)* 
$$|\partial_k^\alpha e'_j(k)| \leq_{const} \begin{cases} 1 & \text{if } |\alpha| \leq 1 \\ \mathfrak{l}_j M^{(|\alpha|-1)j} & \text{if } \alpha \in \Delta, |\alpha| \geq 2 \end{cases}$$

*iii)* If  $|\gamma| = 1$ , then

$$\begin{aligned} |\partial_k^\gamma e'_j(p_0, \mathbf{k}) - \partial_k^\gamma e'_j(k_0, \mathbf{k})|, |\partial_k^\gamma e'(p_0, \mathbf{k}) - \partial_k^\gamma e'(k_0, \mathbf{k})| &\leq_{const} \rho |p_0 - k_0|^\aleph \\ |\partial_{k_0} e'_j(k) - \partial_{k_0} e'_j(k')|, |\partial_{k_0} e'(k) - \partial_{k_0} e'(k')| &\leq_{const} \rho |k - k'|^\aleph \\ |\partial_k^\gamma e'_j(k) - \partial_k^\gamma e'_j(k')|, |\partial_k^\gamma e'(k) - \partial_k^\gamma e'(k')| &\leq_{const} |k - k'|^\aleph \end{aligned}$$

**Proof:** i) By setting  $p^{(i)} = 0$  for all  $i > j$ , it suffices to prove the statements regarding  $e'$ . All statements follow from

$$\sup_k |\partial_k v(k)| \leq \sum_{i=2}^{\infty} \sup_k |\partial_k \check{p}^{(i)}(k)| \leq \sum_{i=2}^{\infty} 2\rho \mathfrak{l}_i \leq \rho$$

For the second inequality, we used Lemma XII.12 of [FKTo3].

ii) Again by Lemma XII.12 of [FKTo3],

$$\begin{aligned} |\partial_k^\alpha e'_j(k)| &\leq |\partial_k^\alpha e(\mathbf{k})| + \sum_{i=2}^j |\partial_k^\alpha \check{p}^{(i)}(k)| \leq |\partial_k^\alpha e(\mathbf{k})| + \sum_{i=2}^j 2\alpha! \rho \mathfrak{l}_i M^{(|\alpha|-1)i} \\ &\leq \text{const} \begin{cases} 1 & \text{if } |\alpha| \leq 1 \\ \mathfrak{l}_j M^{(|\alpha|-1)j} & \text{if } \alpha \in \Delta, |\alpha| \geq 2 \end{cases} \end{aligned}$$

iii) Apply Lemma C.1 of [FKTf3] with  $C_0 = C_1 = \text{const } \rho$ ,  $\alpha = \aleph$  and  $\beta = 1 - \aleph$ . When dealing with  $e'$ , use  $f_i(t) = \partial_k^\gamma \check{p}_i(t, \mathbf{k})$  for the first bound,  $f_i(t) = \partial_{k_0} \check{p}_i(k + t \frac{k' - k}{|k' - k|})$  for the second bound and  $f_i(t) = \partial_k^\gamma \check{p}_i(k + t \frac{k' - k}{|k' - k|})$  for the third bound. When dealing with  $e'_j$  use the above  $f_i$ 's for  $2 \leq i \leq j$  and zero otherwise. The contribution from  $e(\mathbf{k})$  vanishes in the first two bounds and is bounded by  $\text{const } |k - k'|$  in the third. ■

Recall that, for  $j \geq 1$ ,  $C_v^{(j)}(k) = \frac{\nu^{(j)}(k)}{ik_0 - e'(k)}$ . Set, for  $m \geq 1$  and  $s \in \Sigma_m$ ,

$$c_s^{(j)}(k) = C_v^{(j)}(k) \chi_s(k)$$

and denote by  $c_s^{(j)}(x)$  its Fourier transform.

**Lemma A.2** *There are  $M$ -dependent constants  $\text{const}$  and  $\tilde{\rho}_0$  such that the following holds for all  $\rho \leq \tilde{\rho}_0$ ,  $\beta \in \Delta$  and  $j \geq 1$ .*

i) For  $s \in \Sigma_j$ ,

$$\|x^\beta c_s^{(j)}(x)\|_{L^1} \leq \text{const } M^{(1+|\beta|)j}$$

ii) For  $1 \leq m \leq j$  and  $s \in \Sigma_m$ ,

$$\|x^\beta c_s^{(j)}(x)\|_{L^\infty} \leq \text{const } \mathfrak{l}_m M^{(|\beta|-1)j}$$

iii) For  $m \geq 1$  and  $s \in \Sigma_m$ ,

$$\|c_s^{(j)}(x)\|_{L^\infty} \leq \text{const } \frac{\mathfrak{l}_m}{M^j} \quad \left\| \frac{\partial}{\partial x_0} c_s^{(j)}(x) \right\|_{L^\infty} \leq \text{const } \frac{\mathfrak{l}_m}{M^{2j}}$$

iv) For  $s \in \Sigma_j$ ,

$$\left\| \frac{\partial}{\partial x_0} c_s^{(j)}(x) \right\|_{L^1} \leq \text{const}$$

**Proof:** For any sector  $s'$  of any scale,  $c_{s'}^{(j)}(k)$  is supported on the  $j^{\text{th}}$  shell and  $\check{p}^{(i)}(k)$  is supported on the  $i^{\text{th}}$  extended neighbourhood. If  $i > j + 1$ , the  $j$  shell and  $i^{\text{th}}$  extended neighbourhood do not intersect, so we may assume that  $p^{(i)} = 0$  for all  $i > j + 1$ . Therefore, by Corollary XIX.13 and Proposition XIX.4.iii of [FKTo4],

$$|v|_{1, \Sigma_j} \leq \sum_{i=2}^{j+1} |p^{(i)}|_{1, \Sigma_j} \leq \text{const} \sum_{i=2}^{j+1} \frac{\rho \iota_i}{M^j} \mathbf{c}_j \leq \text{const} \frac{\rho}{M^j} \mathbf{c}_j$$

Hence the hypotheses of Proposition XIII.5 of [FKTo3] are fulfilled. To apply this Proposition, we let, for  $s', s'' \in \Sigma_j$ ,  $\tilde{c}((\xi, s'), (\xi', s''))$  be the Fourier transform, as in Definition IX.3 of [FKTo2], of  $\chi_{s'}(k) C_v^{(j)}(k) \chi_{s''}(k)$ . Comparing this Fourier transform with that specified before Definition I.17, we see that

$$c_{s'}^{(j)}(x) = \sum_{s'' \in \Sigma_j} \tilde{c}((0, \uparrow, 0, s'), (x, \uparrow, a, s''))$$

By conservation of momentum, only three  $s''$ 's give nonzero contributions to the right hand side for each  $s'$ . Hence, by parts (ii) and (iii) of Proposition XIII.5 of [FKTo3] and Corollary A.5.i of [FKTo1],

$$\sum_{\delta \in \mathbb{N}_0 \times \mathbb{N}_0^2} \frac{1}{\delta!} \|x^\delta c_{s'}^{(j)}(x)\|_{L^1} t^\delta \leq \text{const} |\tilde{c}|_{1, \Sigma_j} \leq \text{const} \frac{M^j \mathbf{c}_j}{1 - \text{const} \rho \mathbf{c}_j} \leq \text{const} M^j \mathbf{c}_j \quad (\text{A.1})$$

$$\sum_{\delta \in \mathbb{N}_0 \times \mathbb{N}_0^2} \frac{1}{\delta!} \|x^\delta c_{s'}^{(j)}(x)\|_{L^\infty} t^\delta \leq \text{const} \frac{\iota_j}{M^j} \frac{\mathbf{c}_j}{1 - \text{const} \rho \mathbf{c}_j} \leq \text{const} \frac{\iota_j}{M^j} \mathbf{c}_j \quad (\text{A.2})$$

i) follows from (A.1) by choosing  $s' = s$ .

ii) By (A.2),

$$\|x^\delta c_{s'}^{(j)}(x)\|_{L^\infty} \leq \text{const} \frac{\iota_j}{M^j} M^{|\delta|j}$$

for all  $s' \in \Sigma_j$  and  $\delta \in \Delta$ . Write  $c_s^{(j)}(k) = \sum_{\substack{s' \in \Sigma_j \\ s \cap s' \neq \emptyset}} c_{s'}^{(j)}(k) \chi_s(k)$ . By Lemma XII.3.iii of [FKTo3]

$$\begin{aligned} \|x^\beta c_s^{(j)}(x)\|_{L^\infty} &\leq \text{const} \sum_{\substack{s' \in \Sigma_j \\ s \cap s' \neq \emptyset}} \sum_{\substack{\delta, \delta' \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \delta + \delta' = \beta}} \|x^\delta c_{s'}^{(j)}(x)\|_{L^\infty} \|x^{\delta'} \hat{\chi}_s(x)\|_{L^1} \\ &\leq \text{const} \sum_{\substack{s' \in \Sigma_j \\ s \cap s' \neq \emptyset}} \sum_{\substack{\delta, \delta' \in \mathbb{N}_0 \times \mathbb{N}_0^2 \\ \delta + \delta' = \beta}} \frac{\iota_j}{M^j} M^{|\delta|j} M^{|\delta'|j} \leq \text{const} \frac{\iota_m}{\iota_j} \frac{\iota_j}{M^j} M^{|\beta|j} \end{aligned}$$

iii) follows from the observations that

$$\sup_k |c_s^{(j)}(k)| \leq \text{const} M^j \quad \sup_k |k_0 c_s^{(j)}(k)| \leq \text{const}$$

and  $c_s^{(j)}(k)$  is supported in a region of volume  $\text{const } \frac{l_m}{M^{2j}}$ .

iv) follows from part (iv) of Proposition XIII.5 of [FKTo3]. ■

Recall from Lemma III.25 that

$$\Delta_{s_1, s_2}^{m_1, m_2}(\mathbf{p}, \mathbf{k}) = \int_{-b(m_1, m_2)}^{b(m_1, m_2)} d\omega \frac{\mu^{(m_1, m_2)}(\omega, \mathbf{k}, \mathbf{p}) \chi_{s_1}(\mathbf{p}) \chi_{s_2}(\mathbf{k})}{[i\omega - e'(\omega, \mathbf{p})][i\omega - e'(\omega, \mathbf{k})]}$$

where

$$\mu^{(m_1, m_2)}(\omega, \mathbf{k}, \mathbf{p}) = \left[1 - \nu_0(\omega) \nu_1(\mathbf{p}, \mathbf{k})\right] \nu^{(m_1)}((\omega, \mathbf{p})) \nu^{(m_2)}((\omega, \mathbf{k}))$$

and

$$b(m_1, m_2) = \begin{cases} \frac{\text{const}}{M^{\max\{m_1, m_2\}}} & \text{if } m = i, i+1 \\ \min \left\{ \frac{1}{M^{j-3/4}}, \frac{\text{const}}{M^{\max\{m_1, m_2\}}} \right\} & \text{if } m \geq i+2 \end{cases}$$

The functions  $\nu_0$  and  $\nu_1$  were defined just before (III.22).

**Lemma A.3** *Let  $i \leq m \leq j$ ,  $\min\{m_1, m_2\} = m$  and  $s_1, s_2 \in \Sigma_m$ . Then*

$$\begin{aligned} \int d\mathbf{z}_1 \sup_{\mathbf{z}_2} |\hat{\Delta}_{s_1, s_2}^{m_1, m_2}(\mathbf{z}_1, \mathbf{z}_2)| &\leq \text{const } b(m_1, m_2) \, l_m^2 \frac{M^{m_1}}{l_{m_1}} \\ \int d\mathbf{z}_2 \sup_{\mathbf{z}_1} |\hat{\Delta}_{s_1, s_2}^{m_1, m_2}(\mathbf{z}_1, \mathbf{z}_2)| &\leq \text{const } b(m_1, m_2) \, l_m^2 \frac{M^{m_2}}{l_{m_2}} \end{aligned}$$

**Proof:** We may write  $\Delta_{s_1, s_2}^{m_1, m_2}(\mathbf{p}, \mathbf{k})$  as a sum of two pieces, each of the form

$$\sum_{\substack{u_1, v_1 \in \Sigma_{m_1} \\ u_1 \cap s_1 \neq \emptyset \\ u_1 \cap v_1 \neq \emptyset}} \sum_{\substack{u_2, v_2 \in \Sigma_{m_2} \\ u_2 \cap s_2 \neq \emptyset \\ u_2 \cap v_2 \neq \emptyset}} \Delta_{u_1, v_1, u_2, v_2}(\mathbf{p}, \mathbf{k})$$

with

$$\Delta_{u_1, v_1, u_2, v_2}(\mathbf{p}, \mathbf{k}) = \pm \int d\omega \, \zeta_{m_1, m_2}(\omega) c_{u_1}^{(m_1)}(\omega, \mathbf{p}) c_{u_2}^{(m_2)}(\omega, \mathbf{k}) \chi_{1, v_1}(\omega, \mathbf{p}) \chi_{2, v_2}(\omega, \mathbf{k})$$

where

$$\begin{aligned} \chi_{1, v_1}(\omega, \mathbf{p}) &= \left\{ \sum_{\ell=i+1}^{\infty} \nu(M^{2\ell} e(\mathbf{p})^2) \quad \text{or} \right\} \chi_{v_1}(\omega, \mathbf{p}) \chi_{s_1}(0, \mathbf{p}) \\ \chi_{2, v_2}(\omega, \mathbf{k}) &= \left\{ \sum_{\ell=i+1}^{\infty} \nu(M^{2\ell} e(\mathbf{k})^2) \quad \text{or} \right\} \chi_{v_2}(\omega, \mathbf{k}) \chi_{s_2}(0, \mathbf{k}) \\ \zeta_{m_1, m_2}(\omega) &= \left\{ \begin{matrix} 1 \\ \nu_0(\omega) \end{matrix} \quad \text{or} \right\} \chi_{m_1, m_2}(\omega) \end{aligned}$$

and  $\chi_{m_1, m_2}(\omega)$  is the characteristic function of the interval  $[-b(m_1, m_2), b(m_1, m_2)]$ . The Fourier transform of  $\Delta_{u_1, v_1, u_2, v_2}(\mathbf{p}, \mathbf{k})$  is then

$$\begin{aligned} \hat{\Delta}_{u_1, v_1, u_2, v_2}(\mathbf{z}_1, \mathbf{z}_2) &= \int d\mathbf{x}_1 d\mathbf{x}_2 \prod_{\ell=1}^4 dt_\ell c_{u_\ell}^{(m_\ell)}(t_\ell, \mathbf{x}_\ell) \hat{\chi}_{1, v_1}(t_2 - t_1, \mathbf{z}_1 - \mathbf{x}_1) \\ &\quad c_{u_2}^{(m_2)}(t_3 - t_2, \mathbf{x}_2) \hat{\chi}_{2, v_2}(t_4 - t_3, \mathbf{z}_2 - \mathbf{x}_2) \hat{\zeta}_{m_1, m_2}(-t_4) \end{aligned}$$

This is bounded using

$$\begin{aligned} \sup_{t_4} |\hat{\zeta}_{m_1, m_2}(-t_4)| &\leq 2b(m_1, m_2) \\ \sup_{t_3, \mathbf{x}_2, \mathbf{z}_2} \int dt_4 |\hat{\chi}_{2, v_2}(t_4 - t_3, \mathbf{z}_2 - \mathbf{x}_2)| &\leq \text{const } \frac{\mathfrak{l}_{m_2}}{M^{m_2}} \\ \sup_{t_2} \int dt_3 d\mathbf{x}_2 |c_{u_2}^{(m_2)}(t_3 - t_2, \mathbf{x}_2)| &\leq \text{const } M^{m_2} \\ \sup_{t_1, \mathbf{x}_1} \int dt_2 d\mathbf{z}_1 |\hat{\chi}_{1, v_1}(t_2 - t_1, \mathbf{z}_1 - \mathbf{x}_1)| &\leq \text{const} \\ \int dt_1 d\mathbf{x}_1 |c_{u_1}^{(m_1)}(t_1, \mathbf{x}_1)| &\leq \text{const } M^{m_1} \end{aligned} \tag{A.3}$$

The supremum of  $|\hat{\zeta}_{m_1, m_2}(-t_4)|$  was bounded by the  $L^1$  norm of  $\zeta_{m_1, m_2}(\omega)$ . The bounds on  $c_{u_\ell}^{(m_\ell)}$ ,  $\ell = 1, 2$  are immediate consequences of Lemma A.2.i with  $\beta = 0$ . The bounds on  $\hat{\chi}_{\ell, v_\ell}$ ,  $\ell = 1, 2$  are proven much as Lemma XIII.3 of [FKTo3]. Indeed (XIII.3) of [FKTo3] applies with  $j = m_\ell$  and  $\mathfrak{l} = \mathfrak{l}_{m_\ell}$ . Denoting by  $\mathbf{x}_\perp$  and  $\mathbf{x}_\parallel$  the components of  $\mathbf{x}$  perpendicular and parallel, respectively, to the Fermi curve at the centre of  $v_\ell$ , Proposition XIII.1.i gives

$$|\hat{\chi}_{\ell, v_\ell}(t, \mathbf{x})| \leq \text{const } \frac{\mathfrak{l}_{m_\ell}}{M^{2m_\ell}} \frac{1}{[1 + M^{-m_\ell} |t|]^2 [1 + M^{-m_\ell} |\mathbf{x}_\perp| + \mathfrak{l}_{m_\ell} |\mathbf{x}_\parallel|]^3}$$

from which the desired bounds follow. Applying, in order, the bounds of (A.3) yields

$$\int d\mathbf{z}_1 \sup_{\mathbf{z}_2} |\hat{\Delta}_{u_1, v_1, u_2, v_2}(\mathbf{z}_1, \mathbf{z}_2)| \leq \text{const } b(m_1, m_2) \mathfrak{l}_{m_2} M^{m_1}$$

Similarly,

$$\int d\mathbf{z}_2 \sup_{\mathbf{z}_1} |\hat{\Delta}_{u_1, v_1, u_2, v_2}(\mathbf{z}_1, \mathbf{z}_2)| \leq \text{const } b(m_1, m_2) \mathfrak{l}_{m_1} M^{m_2}$$

For each fixed  $s_1 \in \Sigma_m$ , there are at most  $\text{const } \frac{\mathfrak{l}_m}{\mathfrak{l}_{m_1}}$  pairs  $(u_1, v_1) \in \Sigma_{m_1}^2$  obeying  $u_1 \cap s_1 \neq \emptyset$ ,  $u_1 \cap v_1 \neq \emptyset$  and for each fixed  $s_2 \in \Sigma_m$ , there are at most  $\text{const } \frac{\mathfrak{l}_m}{\mathfrak{l}_{m_2}}$  pairs  $(u_2, v_2) \in \Sigma_{m_2}^2$  obeying  $u_2 \cap s_2 \neq \emptyset$ ,  $u_2 \cap v_2 \neq \emptyset$ . Hence

$$\begin{aligned} \int d\mathbf{z}_1 \sup_{\mathbf{z}_2} |\hat{\Delta}_{s_1, s_2}^{m_1, m_2}(\mathbf{z}_1, \mathbf{z}_2)| &\leq \text{const } b(m_1, m_2) \frac{\mathfrak{l}_m}{\mathfrak{l}_{m_1}} \frac{\mathfrak{l}_m}{\mathfrak{l}_{m_2}} \mathfrak{l}_{m_2} M^{m_1} \leq \text{const } b(m_1, m_2) \mathfrak{l}_m^2 \frac{M^{m_1}}{\mathfrak{l}_{m_1}} \\ \int d\mathbf{z}_2 \sup_{\mathbf{z}_1} |\hat{\Delta}_{s_1, s_2}^{m_1, m_2}(\mathbf{z}_1, \mathbf{z}_2)| &\leq \text{const } b(m_1, m_2) \frac{\mathfrak{l}_m}{\mathfrak{l}_{m_1}} \frac{\mathfrak{l}_m}{\mathfrak{l}_{m_2}} \mathfrak{l}_{m_1} M^{m_2} \leq \text{const } b(m_1, m_2) \mathfrak{l}_m^2 \frac{M^{m_2}}{\mathfrak{l}_{m_2}} \end{aligned}$$

■

## Appendix B: Bound on the Generalized Model Bubble

Fix, as in Theorem I.20, a sequence,  $p^{(2)}, p^{(3)}, \dots$ , of sectorized, translation invariant functions  $p^{(i)}$  on  $\left((\mathbb{R} \times \mathbb{R}^2) \times \Sigma_i\right)^2$  obeying

$$|p^{(i)}|_{1, \Sigma_i} \leq \frac{\rho \mathfrak{l}_i}{M^i} \mathbf{c}_i \quad \check{p}^{(i)}(0, \mathbf{k}) = 0$$

Let  $I$  be an interval of length  $\mathfrak{l}$  on the Fermi surface  $F$  and  $u(\mathbf{k}, \mathbf{t})$  a function that vanishes unless  $\pi_F(\mathbf{k}) \in I$ , where  $\pi_F$  is projection on the Fermi curve  $F$ . Set, for  $1 \leq i \leq j$ ,

$$B_{i,j}(\mathbf{t}) = \int dk \frac{\nu_0^{(i,j)}(k_0) u(\mathbf{k}, \mathbf{t})}{[ik_0 - e'(k_0, \mathbf{k})][ik_0 - e'(k_0, \mathbf{k} + \mathbf{t})]}$$

where

$$\nu_0^{(i,j)}(k_0) = \sum_{\ell=i+1}^{j-1} \nu(M^{2\ell} k_0^2) \quad e'(k) = e(\mathbf{k}) - \sum_{\ell=2}^{j+1} \check{p}^{(\ell)}(k)$$

**Lemma B.1** *Let  $1 \leq i \leq j$  obey  $M^i \leq \mathfrak{l}_j M^j$ . Then*

$$|\partial_{\mathbf{t}}^\alpha B_{i,j}(\mathbf{t})| \leq \text{const } \mathfrak{l} \max\{1, j \mathfrak{l}_j M^{|\alpha|j}\} \max_{|\beta+\gamma| \leq |\alpha|} \sup_{\mathbf{k}, \mathbf{t}} \frac{1}{M^{i|\beta+\gamma|}} |\partial_{\mathbf{t}}^\beta \partial_{\mathbf{k}}^\gamma u(\mathbf{k}, \mathbf{t})|$$

for all  $|\alpha| \leq 4$  and all  $\mathbf{t}$  in a neighbourhood of the origin.

**Proof:** Let

$$\begin{aligned} E'(k_0, \mathbf{k}, \mathbf{t}, s) &= s e'(k_0, \mathbf{k}) + (1-s) e'(k_0, \mathbf{k} + \mathbf{t}) \\ E(\mathbf{k}, \mathbf{t}, s) &= E'(0, \mathbf{k}, \mathbf{t}, s) = s e(\mathbf{k}) + (1-s) e(\mathbf{k} + \mathbf{t}) \\ w(\mathbf{k}, \mathbf{t}, s) &= 1 - \frac{1}{i} \frac{\partial E'}{\partial k_0}(0, \mathbf{k}, \mathbf{t}, s) \\ \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s) &= E'(k_0, \mathbf{k}, \mathbf{t}, s) - E'(0, \mathbf{k}, \mathbf{t}, s) - k_0 \frac{\partial E'}{\partial k_0}(0, \mathbf{k}, \mathbf{t}, s) \end{aligned}$$

Then

$$\begin{aligned} B_{i,j}(\mathbf{t}) &= \int dk \frac{\nu_0^{(i,j)}(k_0) u(\mathbf{k}, \mathbf{t})}{[ik_0 - e'(k_0, \mathbf{k})][ik_0 - e'(k_0, \mathbf{k} + \mathbf{t})]} \\ &= \int dk \frac{\nu_0^{(i,j)}(k_0) u(\mathbf{k}, \mathbf{t})}{e'(k_0, \mathbf{k}) - e'(k_0, \mathbf{k} + \mathbf{t})} \left[ \frac{1}{ik_0 - e'(k_0, \mathbf{k})} - \frac{1}{ik_0 - e'(k_0, \mathbf{k} + \mathbf{t})} \right] \\ &= \int dk \int_0^1 ds \frac{\nu_0^{(i,j)}(k_0) u(\mathbf{k}, \mathbf{t})}{[ik_0 - E'(k_0, \mathbf{k}, \mathbf{t}, s)]^2} \\ &= \int dk \int_0^1 ds \frac{\nu_0^{(i,j)}(k_0) u(\mathbf{k}, \mathbf{t})}{[w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)]^2} \end{aligned} \tag{B.1}$$



Case i:  $|\alpha| \geq 1$  Make, for each fixed  $s$ , the change of variables from  $\mathbf{k}$  to  $E$  and an “angular” variable  $\theta$ . Denote by  $J(E, \mathbf{t}, \theta, s)$  the Jacobian of this change of variables. Then

$$B_{i,j}(\mathbf{t}) = \int_0^1 ds \int dk_0 \int d\theta dE \frac{\nu_0^{(i,j)}(k_0) u(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}) J(E, \mathbf{t}, \theta, s)}{[w(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E - \tilde{E}(k_0, \mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}, s)]^2} \quad (\text{B.2})$$

Since  $E(\mathbf{k}, \mathbf{t}, s) = se(\mathbf{k}) + (1-s)e(\mathbf{k} + \mathbf{t})$  and  $\mathbf{t}$  is restricted to a small neighbourhood of the origin,

$$|\nabla_{\mathbf{k}} E(\mathbf{k}, \mathbf{t}, s)| \geq \text{const} > 0 \quad |\partial_{\mathbf{k}}^\alpha \partial_{\mathbf{t}}^\beta E(\mathbf{k}, \mathbf{t}, s)| \leq \text{const}' \quad (\text{B.3})$$

for all  $\alpha + \beta$  having spatial component at most  $r$ . Using

$$\partial_{\mathbf{t}_i} \mathbf{k}_\ell(E, \mathbf{t}, \theta, s) = - \frac{\partial_{\mathbf{k}_2} \theta(\mathbf{k}) \delta_{\ell,1} - \partial_{\mathbf{k}_1} \theta(\mathbf{k}) \delta_{\ell,2}}{\partial_{\mathbf{k}_1} E(\mathbf{k}, \mathbf{t}, s) \partial_{\mathbf{k}_2} \theta(\mathbf{k}) - \partial_{\mathbf{k}_2} E(\mathbf{k}, \mathbf{t}, s) \partial_{\mathbf{k}_1} \theta(\mathbf{k})} \partial_{\mathbf{t}_i} E(\mathbf{k}, \mathbf{t}, s) \Big|_{\mathbf{k}=\mathbf{k}(E, \mathbf{t}, \theta, s)}$$

one proves, by induction on  $|\beta|$ , that, for  $|\beta| \leq r$ ,

$$|\partial_{\mathbf{t}}^\beta \mathbf{k}(E, \mathbf{t}, \theta, s)| \leq \text{const}'' \quad (\text{B.4})$$

Using this bound and

$$J(E, \mathbf{t}, \theta, s) = \frac{1}{|\partial_{\mathbf{k}_1} E(\mathbf{k}, \mathbf{t}, s) \partial_{\mathbf{k}_2} \theta(\mathbf{k}) - \partial_{\mathbf{k}_2} E(\mathbf{k}, \mathbf{t}, s) \partial_{\mathbf{k}_1} \theta(\mathbf{k})|} \Big|_{\mathbf{k}=\mathbf{k}(E, \mathbf{t}, \theta, s)}$$

one proves that, for  $|\beta| \leq r-1$ ,

$$|\partial_{\mathbf{t}}^\beta J(E, \mathbf{t}, \theta, s)| \leq \text{const}'' \quad (\text{B.5})$$

By Lemma A.1.ii, for  $\alpha + \beta + (1, 0, 0) \in \Delta$ ,

$$|\partial_{\mathbf{k}}^\alpha \partial_{\mathbf{t}}^\beta w(\mathbf{k}, \mathbf{t}, s)| \leq \text{const} \begin{cases} 1 & \text{if } |\alpha| + |\beta| = 0 \\ \mathfrak{l}_j M^{(|\alpha|+|\beta|)j} & \text{if } |\alpha| + |\beta| \geq 1 \end{cases} \quad (\text{B.6})$$

Since

$$\begin{aligned} \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s) &= E'(k_0, \mathbf{k}, \mathbf{t}, s) - E'(0, \mathbf{k}, \mathbf{t}, s) - k_0 \frac{\partial E'}{\partial k_0}(0, \mathbf{k}, \mathbf{t}, s) \\ &= \int_0^{k_0} d\kappa \left[ \frac{\partial E'}{\partial k_0}(\kappa, \mathbf{k}, \mathbf{t}, s) - \frac{\partial E'}{\partial k_0}(0, \mathbf{k}, \mathbf{t}, s) \right] \end{aligned}$$

parts (ii) and (iii) of Lemma A.1 imply that, again for  $\alpha + \beta + (1, 0, 0) \in \Delta$ ,

$$|\partial_{\mathbf{k}}^\alpha \partial_{\mathbf{t}}^\beta \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)| \leq \text{const} \rho \begin{cases} |k_0|^{1+\aleph} & \text{if } |\alpha| + |\beta| = 0 \\ |k_0|^\aleph + |k_0| \mathfrak{l}_j M^j & \text{if } |\alpha| + |\beta| = 1 \\ \mathfrak{l}_j M^{(|\alpha|+|\beta|-1)j} + |k_0| \mathfrak{l}_j M^{(|\alpha|+|\beta|)j} & \text{if } |\alpha| + |\beta| > 1 \end{cases}$$

For  $k_0$  in the support of  $\nu_0^{(i,j)}(k_0)$ ,  $|k_0| \geq \text{const} \frac{1}{M^j}$  and

$$|\partial_k^\alpha \partial_t^\beta \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)| \leq \text{const} \rho \begin{cases} |k_0|^{1+\aleph} & \text{if } |\alpha| + |\beta| = 0 \\ |k_0| \mathfrak{l}_j M^{(|\alpha|+|\beta|)j} & \text{if } |\alpha| + |\beta| \geq 1 \end{cases} \quad (\text{B.7})$$

Applying  $\partial_t^\alpha$  to (B.2) yields an integral whose integrand is a sum of terms (whose number is bounded by a universal constant) of the form a combinatorial factor (which is bounded by a universal constant) times  $\nu_0^{(i,j)}(k_0)$  times

$$\frac{1}{[\imath w k_0 - E - \tilde{E}]^{m+2}} \prod_{p=1}^m \left[ -\partial_{\mathbf{k}}^{\gamma^{(p)}} \partial_{\mathbf{t}}^{\beta^{(p)}} (\imath k_0 w - \tilde{E}) \prod_{\ell=1}^{|\gamma^{(p)}|} \partial_{\mathbf{t}}^{\alpha^{(p,\ell)}} \mathbf{k}_{i_\ell} \right] \left[ \partial_{\mathbf{k}}^{\gamma'} \partial_{\mathbf{t}}^{\beta'} u \prod_{\ell'=1}^{|\gamma'|} \partial_{\mathbf{t}}^{\alpha'^{(\ell')}} \mathbf{k}_{i_{\ell'}} \right] \partial_{\mathbf{t}}^{\beta''} J$$

with the various degrees obeying

$$\begin{aligned} \beta'' + \beta' + \sum_{\ell'=1}^{|\gamma'|} \alpha'^{(\ell')} + \sum_{p=1}^m \left[ \beta^{(p)} + \sum_{\ell=1}^{|\gamma^{(p)}|} \alpha^{(p,\ell)} \right] &= \alpha \\ |\gamma^{(p)}| + |\beta^{(p)}| &\geq 1 \quad \text{for all } 1 \leq p \leq m \\ |\alpha^{(p,\ell)}| &\geq 1 \quad \text{for all } 1 \leq p \leq m, 1 \leq \ell \leq |\gamma^{(p)}| \\ |\alpha'^{(\ell')}| &\geq 1 \quad \text{for all } 1 \leq \ell' \leq |\gamma'| \end{aligned}$$

Using the fact, from Lemma A.1.i, that  $|w(\mathbf{k}, \mathbf{t}, s) - 1| \leq \rho \leq \frac{1}{2}$  and the bounds on the derivatives of  $\mathbf{k}$ ,  $J$ ,  $w$  and  $\tilde{E}$  of (B.4–B.7), we may bound this term by

$$\begin{aligned} &\text{const}' \frac{1}{|\imath k_0 - E|^{m+2}} \prod_{p=1}^m \left[ |k_0| \mathfrak{l}_j M^{(|\gamma^{(p)}|+|\beta^{(p)}|)j} \right] \sup_{\mathbf{k}, \mathbf{t}} |\partial_{\mathbf{k}}^{\gamma'} \partial_{\mathbf{t}}^{\beta'} u| \\ &\leq \text{const}' \frac{|k_0|^m}{|\imath k_0 - E|^{m+2}} \prod_{p=1}^m \left[ \mathfrak{l}_j M^{(|\gamma^{(p)}|+|\beta^{(p)}|)j} \right] M^{i(|\beta'|+|\gamma'|)} \max_{|\beta+\gamma| \leq |\alpha|} \sup_{\mathbf{k}, \mathbf{t}} \frac{1}{M^{i|\beta+\gamma|}} |\partial_{\mathbf{t}}^\beta \partial_{\mathbf{k}}^\gamma u(\mathbf{k}, \mathbf{t})| \\ &\leq \text{const}' \frac{|k_0|^m}{|\imath k_0 - E|^{m+2}} M^{|\alpha-\beta''|j} M^{|\beta'+\gamma'|j} \mathfrak{l}_j^m \max_{|\beta+\gamma| \leq |\alpha|} \sup_{\mathbf{k}, \mathbf{t}} \frac{1}{M^{i|\beta+\gamma|}} |\partial_{\mathbf{t}}^\beta \partial_{\mathbf{k}}^\gamma u(\mathbf{k}, \mathbf{t})| \\ &\leq \text{const}' \frac{1}{|\imath k_0 - E|^2} M^{|\alpha|j} \mathfrak{l}_j \max_{|\beta+\gamma| \leq |\alpha|} \sup_{\mathbf{k}, \mathbf{t}} \frac{1}{M^{i|\beta+\gamma|}} |\partial_{\mathbf{t}}^\beta \partial_{\mathbf{k}}^\gamma u(\mathbf{k}, \mathbf{t})| \end{aligned}$$

For the second inequality, we used that  $|\beta'| + |\gamma'| + \sum_{p=1}^m [|\gamma^{(p)}| + |\beta^{(p)}|] \leq |\alpha - \beta''|$ . For the final inequality we used that one of  $m$ ,  $|\gamma'|$ ,  $|\beta'|$ ,  $|\beta''|$  must be nonzero for  $|\alpha|$  to be nonzero and we also used the hypothesis that  $M^{i-j} \leq \mathfrak{l}_j$ . The bound is completed by applying

$$\int_{\frac{\text{const}}{M^j}}^{\text{const}} dk_0 \int_I d\theta \int_{-\text{const}}^{\text{const}} dE \frac{1}{|\imath k_0 - E|^2} \leq \text{const}' \mathfrak{l} \int_{\frac{\text{const}}{M^j}}^{\text{const}} dR \frac{1}{R} \leq \text{const} j \mathfrak{l}$$

Case ii:  $\alpha = 0$ . Recall from (B.1) that

$$B_{i,j}(\mathbf{t}) = \int dk \int_0^1 ds \frac{\nu_0^{(i,j)}(k_0)u(\mathbf{k}, \mathbf{t})}{[iw(\mathbf{k}, \mathbf{t}, s)k_0 - E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)]^2}$$

and set

$$B'_{i,j}(\mathbf{t}) = \int dk \int_0^1 ds \frac{\nu_0^{(i,j)}(k_0)u(\mathbf{k}, \mathbf{t})}{[iw(\mathbf{k}, \mathbf{t}, s)k_0 - E(\mathbf{k}, \mathbf{t}, s)]^2}$$

By Lemma A.1.i, (B.7) and the reality of  $k_0$  and  $E(\mathbf{k}, \mathbf{t}, s)$ ,

$$\frac{1}{4}|ik_0 - E(\mathbf{k}, \mathbf{t}, s)| \leq |iw(\mathbf{k}, \mathbf{t}, s)k_0 - E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)| \leq 2|ik_0 - E(\mathbf{k}, \mathbf{t}, s)|$$

Hence, by (B.7),

$$\begin{aligned} & |B_{i,j}(\mathbf{t}) - B'_{i,j}(\mathbf{t})| \\ & \leq \int dk \int_0^1 ds |\nu_0^{(i,j)}(k_0)u(\mathbf{k}, \mathbf{t})| \left| \frac{[iw(\mathbf{k}, \mathbf{t}, s)k_0 - E(\mathbf{k}, \mathbf{t}, s)]^2 - [iw(\mathbf{k}, \mathbf{t}, s)k_0 - E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)]^2}{[iw(\mathbf{k}, \mathbf{t}, s)k_0 - E(\mathbf{k}, \mathbf{t}, s)]^2 [iw(\mathbf{k}, \mathbf{t}, s)k_0 - E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)]^2} \right| \\ & = \int dk \int_0^1 ds |\nu_0^{(i,j)}(k_0)u(\mathbf{k}, \mathbf{t})| \left| \frac{\tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)[2iw(\mathbf{k}, \mathbf{t}, s)k_0 - 2E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)]}{[iw(\mathbf{k}, \mathbf{t}, s)k_0 - E(\mathbf{k}, \mathbf{t}, s)]^2 [iw(\mathbf{k}, \mathbf{t}, s)k_0 - E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)]^2} \right| \\ & \leq \int dk \int_0^1 ds \text{const } |k_0|^{\aleph} |\nu_0^{(i,j)}(k_0)u(\mathbf{k}, \mathbf{t})| \frac{|\tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)|}{|ik_0 - E(\mathbf{k}, \mathbf{t}, s)|^3} \\ & \leq \int dk \int_0^1 ds \text{const } |k_0|^{\aleph} |\nu_0^{(i,j)}(k_0)u(\mathbf{k}, \mathbf{t})| \frac{1}{|ik_0 - E(\mathbf{k}, \mathbf{t}, s)|^2} \\ & \leq \text{const} \sup_{\mathbf{k}, \mathbf{t}} |u(\mathbf{k}, \mathbf{t})| \int_I d\theta \int_{-const}^{const} dk_0 \int_{-const}^{const} dE \frac{|k_0|^{\aleph}}{|ik_0 - E|^2} \\ & \leq \text{const} \sup_{\mathbf{k}, \mathbf{t}} |u(\mathbf{k}, \mathbf{t})| \int_I d\theta \int_{-const}^{const} dk_0 |k_0|^{\aleph-1} \int dE' \frac{1}{|i - E'|^2} \\ & \leq \text{const} \sup_{\mathbf{k}, \mathbf{t}} |u(\mathbf{k}, \mathbf{t})| \end{aligned}$$

and it suffices to consider  $B'_{i,j}(\mathbf{t})$ .

Make, for each fixed  $s$ , the change of variables from  $\mathbf{k}$  to  $E = E(\mathbf{k}, \mathbf{t}, s)$  and an “angular” variable  $\theta$ . Then

$$B'_{i,j}(\mathbf{t}) = \int_0^1 ds \int d\theta \int dk_0 dE \nu_0^{(i,j)}(k_0) \frac{u(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t})J(E, \mathbf{t}, \theta, s)}{[iw(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}, s)k_0 - E]^2}$$

Integrating by parts with respect to  $k_0$ ,

$$\begin{aligned} B'_{i,j}(\mathbf{t}) &= \int_0^1 ds \int d\theta dE \int_{-\infty}^{\infty} dk_0 \nu_0^{(i,j)}(k_0) \frac{uJ}{w} i \frac{d}{dk_0} \frac{1}{iwk_0 - E} \\ &= \int_0^1 ds \int d\theta dE \int_{-\infty}^{\infty} dk_0 \frac{\frac{uJ}{w} (-i) \frac{d}{dk_0} \nu_0^{(i,j)}(k_0)}{iwk_0 - E} \end{aligned}$$

Since  $\frac{d}{dk_0}\nu_0^{(i,j)}(k_0)$  is odd under  $k_0 \rightarrow -k_0$ ,

$$\begin{aligned} B'_{i,j}(\mathbf{t}) &= \int_0^1 ds \int d\theta dE \int_0^\infty dk_0 \frac{uJ}{w} (-i) \left[ \frac{d}{dk_0}\nu_0^{(i,j)}(k_0) \right] \left[ \frac{1}{iwk_0-E} - \frac{1}{-iwk_0-E} \right] \\ &= \int_0^1 ds \int d\theta dE \int_0^\infty dk_0 \frac{uJ}{w} (-i) \left[ \frac{d}{dk_0}\nu_0^{(i,j)}(k_0) \right] \frac{-2iwk_0}{w^2k_0^2+E^2} \\ &= -2 \int_0^1 ds \int d\theta \int_{-\infty}^\infty dE \int_0^\infty dk_0 \left[ \frac{d}{dk_0}\nu_0^{(i,j)}(k_0) \right] \frac{u(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}) J(E, \mathbf{t}, \theta, s) k_0}{w(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}, s)^2 k_0^2 + E^2} \end{aligned}$$

Hence, since  $|w(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}, s) - 1| \leq \rho \leq \frac{1}{5}$ ,  $|w(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}, s)^2 - 1| \leq \frac{1}{2}$  and

$$|B'_{i,j}(\mathbf{t})| \leq 4 \sup |uJ| \int_0^1 ds \int_{I'} d\theta \int_0^\infty dk_0 \int_{-\infty}^\infty dE \left| \frac{d}{dk_0}\nu_0^{(i,j)}(k_0) \right| \frac{k_0}{k_0^2 + E^2}$$

where  $I'$  is some interval of length  $\text{const } \mathfrak{l}$ . Finally,

$$\begin{aligned} |B'_{i,j}(\mathbf{t})| &\leq \text{const } \mathfrak{l} \sup |u| \int_0^\infty dk_0 \left| \frac{d}{dk_0}\nu_0^{(i,j)}(k_0) \right| \\ &\leq \text{const } \mathfrak{l} \sup_{\mathbf{k}, \mathbf{t}} |u(\mathbf{k}, \mathbf{t})| \end{aligned}$$

since  $\int_0^\infty dk_0 \left| \frac{d}{dk_0}\nu_0^{(i,j)}(k_0) \right| = 2$ . ■

**Theorem B.2** *Let  $1 \leq i \leq j$  obey  $M^i \leq \mathfrak{l}_j M^j$ . Let  $\mathbf{t}$  and  $\mathbf{n}$  be mutually perpendicular unit vectors in  $\mathbb{R}^2$  and  $\rho(\mathbf{k})$  be a function that is supported in a rectangle in  $\mathbf{k}$  having one side of length  $\frac{\text{const}}{M^j}$  parallel to  $\mathbf{n}$  and one side of length  $\text{const } \mathfrak{l}_j$  parallel to  $\mathbf{t}$ . Furthermore assume that, for all  $\alpha_1, \alpha_2 \leq 2$ ,*

$$\left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}})^{\alpha_2} \rho(\mathbf{k}) \right| \leq \text{const } M^{\alpha_1 j} \frac{1}{\mathfrak{l}_j^{\alpha_2}}$$

Let  $a = \frac{1/2}{1-\mathfrak{R}}$  and  $\hat{B}_{i,j}(\mathbf{x})$  be the Fourier transform of  $\rho(\mathbf{k}) B_{i,j}(\mathbf{k})$ . Then

$$|\hat{B}_{i,j}(\mathbf{x})| \leq \text{const } \frac{\mathfrak{l}_j}{M^j} \frac{1}{(1+|\mathbf{n} \cdot \mathbf{x}/M^j|^{3/2})(1+|\mathfrak{l}_j \mathbf{t} \cdot \mathbf{x}|^{(1+a)/2})} \max_{|\beta+\gamma| \leq 3} \sup_{\mathbf{k}, \mathbf{t}} \frac{1}{M^{i|\beta+\gamma|}} |\partial_{\mathbf{t}}^\beta \partial_{\mathbf{k}}^\gamma u(\mathbf{k}, \mathbf{t})|$$

**Proof:** Denote

$$U = \max_{|\beta+\gamma| \leq 3} \sup_{\mathbf{k}, \mathbf{t}} \frac{1}{M^{i|\beta+\gamma|}} |\partial_{\mathbf{t}}^\beta \partial_{\mathbf{k}}^\gamma u(\mathbf{k}, \mathbf{t})|$$

Note that  $1 < a < \frac{3}{2}$ . The first step is to prove that for all  $\alpha_1 \in \{0, 1\}$  and  $0 \leq \alpha_2 \leq a$

$$\left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}})^{[\alpha_2]} [B_{i,j}(\mathbf{k} + q\mathbf{t}) - B_{i,j}(\mathbf{k})] \right| \leq C U M^{\alpha_1 j} \frac{1}{\mathfrak{l}_j^{\alpha_2}} |q|^{\alpha_2 - [\alpha_2]} \quad (\text{B.8})$$

where  $[\alpha_2]$  is the integer part of  $\alpha_2$ . Here  $C$  is a constant that is independent of  $i, j, \mathbf{k}$  and  $q$ . To prove (B.8) when  $[\alpha_2] = 0$ , apply Lemma B.1 twice to obtain

$$\begin{aligned} \left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} [B_{i,j}(\mathbf{k} + q\mathbf{t}) - B_{i,j}(\mathbf{k})] \right| &\leq \left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} B_{i,j}(\mathbf{k} + q\mathbf{t}) \right| + \left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} B_{i,j}(\mathbf{k}) \right| \\ &\leq 2 \text{const } \mathcal{U} M^{\alpha_1 j} \\ \left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} [B_{i,j}(\mathbf{k} + q\mathbf{t}) - B_{i,j}(\mathbf{k})] \right| &\leq |q| \sup_{\mathbf{p}} \left| (\mathbf{n} \cdot \partial_{\mathbf{p}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{p}}) B_{i,j}(\mathbf{p}) \right| \\ &\leq \text{const } \mathcal{U} |q| j \mathfrak{l}_j M^{(\alpha_1+1)j} \end{aligned}$$

Multiplying the  $(1 - \alpha_2)^{\text{th}}$  power of the first bound by the  $\alpha_2^{\text{th}}$  power of the second gives

$$\begin{aligned} \left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} [B_{i,j}(\mathbf{k} + q\mathbf{t}) - B_{i,j}(\mathbf{k})] \right| &\leq 2^{1-\alpha_2} \text{const } \mathcal{U} |q|^{\alpha_2} M^{\alpha_1 j} (j \mathfrak{l}_j M^j)^{\alpha_2} \\ &= 2^{1-\alpha_2} \text{const } \mathcal{U} M^{\alpha_1 j} \frac{1}{\mathfrak{l}_j^{\alpha_2}} |q|^{\alpha_2} (j M^{(1-2\aleph)j})^{\alpha_2} \\ &\leq C \mathcal{U} M^{\alpha_1 j} \frac{1}{\mathfrak{l}_j^{\alpha_2}} |q|^{\alpha_2} \end{aligned}$$

To prove (B.8) when  $[\alpha_2] = 1$ , again apply Lemma B.1 twice to obtain

$$\begin{aligned} \left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}}) [B_{i,j}(\mathbf{k} + q\mathbf{t}) - B_{i,j}(\mathbf{k})] \right| &\leq 2 \text{const } \mathcal{U} j \mathfrak{l}_j M^{(\alpha_1+1)j} \\ \left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}}) [B_{i,j}(\mathbf{k} + q\mathbf{t}) - B_{i,j}(\mathbf{k})] \right| &\leq |q| \sup_{\mathbf{p}} \left| (\mathbf{n} \cdot \partial_{\mathbf{p}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{p}})^2 B_{i,j}(\mathbf{p}) \right| \\ &\leq \text{const } \mathcal{U} |q| j \mathfrak{l}_j M^{(\alpha_1+2)j} \end{aligned}$$

Multiplying the  $(2 - \alpha_2)^{\text{th}}$  power of the first bound by the  $(\alpha_2 - 1)^{\text{th}}$  power of the second gives

$$\begin{aligned} \left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}}) [B_{i,j}(\mathbf{k} + q\mathbf{t}) - B_{i,j}(\mathbf{k})] \right| &\leq 2^{2-\alpha_2} \text{const } \mathcal{U} |q|^{\alpha_2-1} M^{(\alpha_1+1)j} j \mathfrak{l}_j M^{(\alpha_2-1)j} \\ &= 2^{2-\alpha_2} \text{const } \mathcal{U} M^{\alpha_1 j} \frac{1}{\mathfrak{l}_j^{\alpha_2}} |q|^{\alpha_2-1} (j M^{-(\aleph+\alpha_2\aleph-\alpha_2)j}) \\ &\leq C \mathcal{U} M^{\alpha_1 j} \frac{1}{\mathfrak{l}_j^{\alpha_2}} |q|^{\alpha_2-1} \end{aligned}$$

since  $\aleph + \alpha_2\aleph - \alpha_2 = \aleph - (1 - \aleph)\alpha_2 \geq \aleph - (1 - \aleph)a = \aleph - \frac{1}{2} > 0$ . We also have, for all  $\alpha_1 \in \{0, 1, 2\}$  and  $\alpha_2 \in \{0, 1\}$ ,

$$\left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}})^{\alpha_2} B_{i,j}(\mathbf{k}) \right| \leq \text{const } \mathcal{U} M^{\alpha_1 j} \frac{1}{\mathfrak{l}_j^{\alpha_2}} \quad (\text{B.9})$$

since  $j \mathfrak{l}_j M^{\alpha_2 j} = \frac{1}{\mathfrak{l}_j^{\alpha_2}} j M^{-(\aleph+\alpha_2\aleph-\alpha_2)j} \leq \text{const } \frac{1}{\mathfrak{l}_j^{\alpha_2}}$  for  $\alpha_2 = 0, 1$ .

The next step is to prove that for all  $\alpha_1 \in \{0, 1\}$  and  $0 \leq \alpha_2 \leq a$

$$\left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}})^{[\alpha_2]} [\rho(\mathbf{k} + q\mathbf{t}) B_{i,j}(\mathbf{k} + q\mathbf{t}) - \rho(\mathbf{k}) B_{i,j}(\mathbf{k})] \right| \leq C' \mathcal{U} M^{\alpha_1 j} \frac{1}{\mathfrak{l}_j^{\alpha_2}} |q|^{\alpha_2 - [\alpha_2]} \quad (\text{B.10})$$

Applying the hypothesis on  $\rho$  twice,

$$\begin{aligned} \left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}})^{[\alpha_2]} [\rho(\mathbf{k} + q\mathbf{t}) - \rho(\mathbf{k})] \right| &\leq 2 \text{const } M^{\alpha_1 j} \frac{1}{l_j^{[\alpha_2]}} \\ \left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}})^{[\alpha_2]} [\rho(\mathbf{k} + q\mathbf{t}) - \rho(\mathbf{k})] \right| &\leq \text{const } M^{\alpha_1 j} \frac{1}{l_j^{[\alpha_2]+1}} |q| \end{aligned}$$

Multiplying the  $(1 + [\alpha_2] - \alpha_2)^{\text{th}}$  power of the first bound by the  $(\alpha_2 - [\alpha_2])^{\text{th}}$  power of the second gives

$$\left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}})^{[\alpha_2]} [\rho(\mathbf{k} + q\mathbf{t}) - \rho(\mathbf{k})] \right| \leq 2^{1+[\alpha_2]-\alpha_2} \text{const } M^{\alpha_1 j} \frac{1}{l_j^{\alpha_2}} |q|^{\alpha_2 - [\alpha_2]} \quad (\text{B.11})$$

The bound (B.10) follows from the product rule and

$$\begin{aligned} &\left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}})^{[\alpha_2]} [\rho(\mathbf{k} + q\mathbf{t}) B_{i,j}(\mathbf{k} + q\mathbf{t}) - \rho(\mathbf{k}) B_{i,j}(\mathbf{k})] \right| \\ &\leq \left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}})^{[\alpha_2]} [\rho(\mathbf{k} + q\mathbf{t}) B_{i,j}(\mathbf{k} + q\mathbf{t}) - \rho(\mathbf{k}) B_{i,j}(\mathbf{k} + q\mathbf{t})] \right| \\ &\quad + \left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}})^{[\alpha_2]} [\rho(\mathbf{k}) B_{i,j}(\mathbf{k} + q\mathbf{t}) - \rho(\mathbf{k}) B_{i,j}(\mathbf{k})] \right| \end{aligned}$$

by using (B.11) and (B.9) to bound the first line and (B.8) and the hypothesis on the derivatives of  $\rho$ , to bound the second.

The Lemma will follow from

$$\sup_{\mathbf{x}} \left| \mathbf{n} \cdot \mathbf{x} / M^j \right|^{\alpha_1} |l_j \mathbf{t} \cdot \mathbf{x}|^{\alpha_2} |\hat{B}_{i,j}(\mathbf{x})| \leq \text{const } lU \frac{l_j}{M^j}$$

for all  $\alpha_1 \in \{0, \frac{3}{2}\}$  and  $\alpha_2 \in \{0, \frac{1+a}{2}\}$ . This in turn will follow from

$$\sup_{\mathbf{x}} \left| \mathbf{n} \cdot \mathbf{x} / M^j \right|^{\alpha_1} |l_j \mathbf{t} \cdot \mathbf{x}|^{\alpha_2} |\hat{B}_{i,j}(\mathbf{x})| \leq \text{const } lU \frac{l_j}{M^j} \quad (\text{B.12})$$

for all  $\alpha_1 \in \{0, 1, 2\}$ ,  $\alpha_2 \in \{0, 1, a\}$ ,  $(\alpha_1, \alpha_2) \neq (2, a)$ , by taking various geometric means. In particular, to handle the case  $(\alpha_1, \alpha_2) = (\frac{3}{2}, \frac{1+a}{2})$ , take the geometric mean of the bounds with  $(\alpha_1, \alpha_2) = (1, a)$  and  $(\alpha_1, \alpha_2) = (2, 1)$ . For  $\alpha_2 = 0, 1$ , (B.12) follows, by integration by parts, from

$$\left| (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (\mathbf{t} \cdot \partial_{\mathbf{k}})^{\alpha_2} [\rho(\mathbf{k}) B_{i,j}(\mathbf{k})] \right| \leq \text{const } lU M^{\alpha_1 j} \frac{1}{l_j^{\alpha_2}}$$

and the fact that  $\rho(\mathbf{k}) B_{i,j}(\mathbf{k})$  is supported in a region of volume  $\text{const } \frac{l_j}{M^j}$ . Furthermore, if  $|l_j \mathbf{t} \cdot \mathbf{x}| \leq 1$ ,

$$\left| \mathbf{n} \cdot \mathbf{x} / M^j \right|^{\alpha_1} |l_j \mathbf{t} \cdot \mathbf{x}|^a |\hat{B}_{i,j}(\mathbf{x})| \leq \left| \mathbf{n} \cdot \mathbf{x} / M^j \right|^{\alpha_1} |\hat{B}_{i,j}(\mathbf{x})| \leq \text{const } lU \frac{l_j}{M^j}$$

so it suffices to consider  $\alpha_1 = 0, 1$ ,  $\alpha_2 = a$  and  $|\mathbf{t} \cdot \mathbf{x}| \geq \frac{1}{l_j}$ . Let  $\tilde{D}_j(\mathbf{x})$  denote the Fourier transform of

$$D_{i,j}(\mathbf{k}) = \frac{1}{M^{\alpha_1 j}} (\mathbf{n} \cdot \partial_{\mathbf{k}})^{\alpha_1} (l_j \mathbf{t} \cdot \partial_{\mathbf{k}}) [\rho(\mathbf{k}) B_{i,j}(\mathbf{k})]$$

Then

$$\begin{aligned} |e^{-i q \mathbf{t} \cdot \mathbf{x}} - 1| |\mathbf{n} \cdot \mathbf{x} / M^j|^{\alpha_1} |l_j \mathbf{t} \cdot \mathbf{x}| |\hat{B}_{i,j}(\mathbf{x})| &= |e^{-i q \mathbf{t} \cdot \mathbf{x}} - 1| \left| \int \frac{d^2 \mathbf{k}}{(2\pi)^2} e^{i \mathbf{k} \cdot \mathbf{x}} D_{i,j}(\mathbf{k}) \right| \\ &= \left| \int \frac{d^2 \mathbf{k}}{(2\pi)^2} [e^{i(\mathbf{k} - q \mathbf{t}) \cdot \mathbf{x}} - e^{i \mathbf{k} \cdot \mathbf{x}}] D_{i,j}(\mathbf{k}) \right| \\ &= \left| \int \frac{d^2 \mathbf{k}}{(2\pi)^2} e^{i \mathbf{k} \cdot \mathbf{x}} [D_{i,j}(\mathbf{k} + q \mathbf{t}) - D_{i,j}(\mathbf{k})] \right| \end{aligned}$$

By (B.10)

$$|D_{i,j}(\mathbf{k} + q \mathbf{t}) - D_{i,j}(\mathbf{k})| \leq C' l_j \frac{|q|^{a-1}}{l_j^{a-1}}$$

Furthermore, if  $|q| < l_j$ ,  $D_{i,j}(\mathbf{k} + q \mathbf{t}) - D_{i,j}(\mathbf{k})$  is supported in a region of volume  $\text{const } \frac{l_j}{M^j}$ , so that

$$|e^{-i q \mathbf{t} \cdot \mathbf{x}} - 1| |\mathbf{n} \cdot \mathbf{x} / M^j|^{\alpha_1} |l_j \mathbf{t} \cdot \mathbf{x}| |\hat{B}_{i,j}(\mathbf{x})| \leq \text{const } l_j \frac{|q|^{a-1}}{M^j l_j^{a-1}}$$

To finish the proof of (B.12), and the Lemma, it now suffices to choose  $q = \frac{1}{10 \mathbf{t} \cdot \mathbf{x}}$  and observe that then  $|e^{-i q \mathbf{t} \cdot \mathbf{x}} - 1| = |e^{-i/10} - 1| > 0$ . ■

**Lemma B.3** *Let  $I$  be an interval of length  $l$  on the Fermi surface  $F$ ,  $K$  be a compact subset of  $\mathbb{R}^2$ , and  $u_j(k, t_0, z) = u_j((k_0, \mathbf{k}), t_0, z)$  and  $v_j(k, \mathbf{t}, z) = v_j((k_0, \mathbf{k}), \mathbf{t}, z)$ ,  $j > 1$ , be functions that vanish unless  $\mathbf{k} \in K$  and  $\pi_F(\mathbf{k}) \in I$ , where  $\pi_F$  is projection on the Fermi surface. The variable  $z$  runs over  $\mathbb{R}^n$  for some  $n \in \mathbb{N}$ . Let  $n_j(\omega)$ ,  $j > 1$ , be functions that take values in  $[0, 1]$ , vanish in a ( $j$ -dependent) neighbourhood of  $\omega = 0$ , are supported in a compact set that is independent of  $j$  and converge pointwise as  $j \rightarrow \infty$ . Set, for  $j > 1$ ,*

$$\begin{aligned} A_j(t_0, z) &= \int dk \frac{n_j(e(\mathbf{k})) u_j(k, t_0, z)}{[i k_0 - e'(k_0, \mathbf{k})][i(k_0 + t_0) - e'(k_0 + t_0, \mathbf{k})]} \\ B_j(\mathbf{t}, z) &= \int dk \frac{n_j(k_0) v_j(k, \mathbf{t}, z)}{[i k_0 - e'(k_0, \mathbf{k})][i k_0 - e'(k_0, \mathbf{k} + \mathbf{t})]} \end{aligned}$$

and let

$$\begin{aligned} U_{\tilde{\mathbf{x}}} &= \sup_{j,k,t_0,z} |u_j(k, t_0, z)| + \sup_{j,k,t_0,\kappa,z} \frac{|u_j(k - (\kappa, \mathbf{0}), t_0, z) - u_j(k, t_0, z)|}{|\kappa|^{\tilde{\mathbf{x}}}} \\ &\quad + \sup_{j,k,t_0,z} \frac{|u_j(k, t_0, z) - u_j(k, 0, z)|}{|t_0|^{\tilde{\mathbf{x}}}} + \sup_{j,k,t_0,z,z'} \frac{|u_j(k, t_0, z) - u_j(k, t_0, z')|}{|z - z'|^{\tilde{\mathbf{x}}}} \\ V_{\tilde{\mathbf{x}}} &= \sup_{j,k,\mathbf{t},z} |v_j(k, \mathbf{t}, z)| + \sup_{j,k_0,\mathbf{k},\mathbf{k}',\mathbf{t},z} \frac{|v_j((k_0, \mathbf{k}'), \mathbf{t}, z) - v_j((k_0, \mathbf{k}), \mathbf{t}, z)|}{|\mathbf{k} - \mathbf{k}'|^{\tilde{\mathbf{x}}}} \\ &\quad + \sup_{j,k,\mathbf{t},z} \frac{|v_j(k, \mathbf{t}, z) - v_j(k, \mathbf{0}, z)|}{|\mathbf{t}|^{\tilde{\mathbf{x}}}} + \sup_{j,k,\mathbf{t},z,z'} \frac{|v_j(k, \mathbf{t}, z) - v_j(k, \mathbf{t}, z')|}{|z - z'|^{\tilde{\mathbf{x}}}} \end{aligned}$$

be finite.

a) Let  $0 \leq \aleph' < \tilde{\aleph} < \aleph$ . Then

$$\begin{aligned} |A_j(t_0, z) - A_j(0, z)| &\leq \text{const } \mathfrak{l} U_{\tilde{\aleph}} |t_0|^{\aleph'} \\ |A_j(t_0, z) - A_j(t_0, z')| &\leq \text{const } \mathfrak{l} U_{\tilde{\aleph}} |z - z'|^{\aleph'} \end{aligned}$$

for all  $t_0$  in a ( $j$ -independent) neighbourhood of the origin.

b) If, in addition,  $u_j(k, t_0, z)$  converges pointwise as  $j \rightarrow \infty$ , the limit  $A(t_0, z) = \lim_{j \rightarrow \infty} A_j(t_0, z)$  exists for  $t_0$  in a neighbourhood of the origin and obeys

$$\begin{aligned} |A(t_0, z) - A(0, z)| &\leq \text{const } \mathfrak{l} U_{\tilde{\aleph}} |t_0|^{\aleph'} \\ |A(t_0, z) - A(t_0, z')| &\leq \text{const } \mathfrak{l} U_{\tilde{\aleph}} |z - z'|^{\aleph'} \end{aligned}$$

for all  $t_0$  in a neighbourhood of the origin.

c) Let  $0 \leq \aleph' < \tilde{\aleph} < \aleph$ . Then

$$\begin{aligned} |B_j(\mathbf{t}, z) - B_j(\mathbf{0}, z)| &\leq \text{const } \mathfrak{l} V_{\tilde{\aleph}} |\mathbf{t}|^{\aleph'} \\ |B_j(\mathbf{t}, z) - B_j(\mathbf{t}, z')| &\leq \text{const } \mathfrak{l} V_{\tilde{\aleph}} |z - z'|^{\aleph'} \end{aligned}$$

for all  $\mathbf{t}$  in a ( $j$ -independent) neighbourhood of the origin.

d) If, in addition,  $v_j(k, \mathbf{t}, z)$  converges pointwise as  $j \rightarrow \infty$ , the limit  $B(\mathbf{t}, z) = \lim_{j \rightarrow \infty} B_j(\mathbf{t}, z)$  exists for  $\mathbf{t}$  in a neighbourhood of the origin and obeys

$$\begin{aligned} |B(\mathbf{t}, z) - B(\mathbf{0}, z)| &\leq \text{const } \mathfrak{l} V_{\tilde{\aleph}} |\mathbf{t}|^{\aleph'} \\ |B(\mathbf{t}, z) - B(\mathbf{t}, z')| &\leq \text{const } \mathfrak{l} V_{\tilde{\aleph}} |z - z'|^{\aleph'} \end{aligned}$$

**Proof:** The proofs of parts (a) and (b) are very similar to the proofs of parts (c) and (d) respectively. So we only give the latter.

c) As in Lemma B.1,

$$B_j(\mathbf{t}, z) = \int dk \int_0^1 ds \frac{n_j(k_0) v_j(k, \mathbf{t}, z)}{[w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)]^2}$$

where

$$\begin{aligned} E'(k_0, \mathbf{k}, \mathbf{t}, s) &= s e'(k_0, \mathbf{k}) + (1 - s) e'(k_0, \mathbf{k} + \mathbf{t}) \\ E(\mathbf{k}, \mathbf{t}, s) &= E'(0, \mathbf{k}, \mathbf{t}, s) = s e(\mathbf{k}) + (1 - s) e(\mathbf{k} + \mathbf{t}) \\ w(\mathbf{k}, \mathbf{t}, s) &= 1 - \frac{1}{i} \frac{\partial E'}{\partial k_0}(0, \mathbf{k}, \mathbf{t}, s) \\ \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s) &= E'(k_0, \mathbf{k}, \mathbf{t}, s) - E'(0, \mathbf{k}, \mathbf{t}, s) - k_0 \frac{\partial E'}{\partial k_0}(0, \mathbf{k}, \mathbf{t}, s) \end{aligned}$$



Set

$$B'_j(\mathbf{t}, z) = \int dk \int_0^1 ds \frac{n_j(k_0) v_j(k, \mathbf{t}, z)}{[\imath w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s)]^2}$$

and

$$\begin{aligned} I(k_0, \mathbf{k}, \mathbf{t}, s) &= \frac{1}{[\imath w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)]^2} - \frac{1}{[\imath w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s)]^2} \\ &= \frac{[\imath w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s)]^2 - [\imath w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)]^2}{[\imath w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s)]^2 [\imath w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)]^2} \\ &= \frac{\tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s) [2\imath w(\mathbf{k}, \mathbf{t}, s) k_0 - 2E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)]}{[\imath w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s)]^2 [\imath w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)]^2} \end{aligned}$$

so that

$$B_j(\mathbf{t}, z) - B'_j(\mathbf{t}, z) = \int dk \int_0^1 ds n_j(k_0) v_j(k, \mathbf{t}, z) I(k_0, \mathbf{k}, \mathbf{t}, s) \quad (\text{B.13})$$

By (B.7) and Lemma A.1, using  $\tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s) = \int_0^{k_0} d\kappa \left[ \frac{\partial E'}{\partial k_0}(\kappa, \mathbf{k}, \mathbf{t}, s) - \frac{\partial E'}{\partial k_0}(0, \mathbf{k}, \mathbf{t}, s) \right]$ ,

$$\begin{aligned} |\tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s)| &\leq \text{const } \rho |k_0|^{1+\aleph} & |\tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{0}, s)| &\leq \text{const } |k_0| |\mathbf{t}|^\aleph \\ |w(\mathbf{k}, \mathbf{t}, s) - 1| &\leq \rho & |w(\mathbf{k}, \mathbf{t}, s) - w(\mathbf{k}, \mathbf{0}, s)| &\leq \text{const } \rho |\mathbf{t}|^\aleph \\ & & |E(\mathbf{k}, \mathbf{t}, s) - E(\mathbf{k}, \mathbf{0}, s)| &\leq \text{const } |\mathbf{t}| \end{aligned}$$

Consequently

$$|I(k_0, \mathbf{k}, \mathbf{t}, s)| \leq \text{const } \frac{|k_0|^{1+\aleph} |\imath k_0 - E(\mathbf{k}, \mathbf{t}, s)|}{|\imath k_0 - E(\mathbf{k}, \mathbf{t}, s)|^4} \leq \text{const } \frac{|k_0|^\aleph}{|\imath k_0 - E(\mathbf{k}, \mathbf{t}, s)|^2} \quad (\text{B.14})$$

and, we claim, for  $0 \leq \aleph' \leq \aleph$ ,

$$|I(k_0, \mathbf{k}, \mathbf{t}, s) - I(k_0, \mathbf{k}, \mathbf{0}, s)| \leq \text{const } |k_0|^{\aleph-\aleph'} |\mathbf{t}|^{\aleph'} \left[ \frac{1}{|\imath k_0 - E(\mathbf{k}, \mathbf{0}, s)|^2} + \frac{1}{|\imath k_0 - E(\mathbf{k}, \mathbf{t}, s)|^2} \right]$$

We prove the last bound in the case  $|E(\mathbf{k}, \mathbf{t}, s)| \leq |E(\mathbf{k}, \mathbf{0}, s)|$ . The other case is similar. Set

$$\begin{aligned} c &= \imath w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s) & C &= \imath w(\mathbf{k}, \mathbf{0}, s) k_0 - E(\mathbf{k}, \mathbf{0}, s) \\ d &= \imath w(\mathbf{k}, \mathbf{t}, s) k_0 - E(\mathbf{k}, \mathbf{t}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s) & D &= \imath w(\mathbf{k}, \mathbf{0}, s) k_0 - E(\mathbf{k}, \mathbf{0}, s) - \tilde{E}(k_0, \mathbf{k}, \mathbf{0}, s) \\ a &= \tilde{E}(k_0, \mathbf{k}, \mathbf{t}, s) & A &= \tilde{E}(k_0, \mathbf{k}, \mathbf{0}, s) \\ b &= c + d & B &= C + D \end{aligned}$$

Then, for  $|k_0|, |\mathbf{t}| \leq \text{const}$ ,

$$\begin{aligned} |a|, |A| &\leq \text{const } |k_0|^{1+\aleph} \\ |b|, |c|, |d| &\leq \text{const } |\imath k_0 - E(\mathbf{k}, \mathbf{t}, s)| \leq \text{const } |\imath k_0 - E(\mathbf{k}, \mathbf{0}, s)| \\ |c|, |d| &\geq \text{const } |\imath k_0 - E(\mathbf{k}, \mathbf{t}, s)| \\ |B|, |C|, |D| &\leq \text{const } |\imath k_0 - E(\mathbf{k}, \mathbf{0}, s)| \\ |C|, |D| &\geq \text{const } |\imath k_0 - E(\mathbf{k}, \mathbf{0}, s)| \\ |a - A| &\leq \text{const } \min \{ |k_0|^{1+\aleph}, |k_0| |\mathbf{t}|^\aleph \} \leq \text{const } |k_0|^{1+\aleph-\aleph'} |\mathbf{t}|^{\aleph'} \\ |c - C|, |d - D|, |b - B| &\leq \text{const } |k_0| |\mathbf{t}|^\aleph + \text{const } \min \{ |E(\mathbf{k}, \mathbf{0}, s)| + |E(\mathbf{k}, \mathbf{t}, s)|, |\mathbf{t}| \} \\ &\leq \text{const } |\mathbf{t}|^{\aleph'} |\imath k_0 - E(\mathbf{k}, \mathbf{0}, s)|^{1-\aleph'} \end{aligned}$$

Applying these to

$$\begin{aligned}\frac{AB}{C^2 D^2} - \frac{ab}{c^2 d^2} &= \frac{(A-a)B}{C^2 D^2} + \frac{a(B-b)}{C^2 D^2} + \frac{ab}{c^2 d^2} \frac{c^2 d^2 - C^2 D^2}{C^2 D^2} \\ &= \frac{(A-a)B}{C^2 D^2} + \frac{a(B-b)}{C^2 D^2} + \frac{ab}{c^2 d^2} \frac{c^2 (d-D)(d+D)}{C^2 D^2} + \frac{ab}{c^2 d^2} \frac{(c-C)(c+C)}{C^2}\end{aligned}$$

gives

$$\begin{aligned}|I(k_0, \mathbf{k}, \mathbf{t}, s) - I(k_0, \mathbf{k}, \mathbf{0}, s)| &\leq \text{const} \frac{|k_0|^{\aleph-\aleph'} |\mathbf{t}|^{\aleph'}}{|ik_0 - E(\mathbf{k}, \mathbf{0}, s)|^2} + \text{const} \frac{|k_0|^{\aleph}}{|ik_0 - E(\mathbf{k}, \mathbf{t}, s)|^2} \frac{|\mathbf{t}|^{\aleph'}}{|ik_0 - E(\mathbf{k}, \mathbf{0}, s)|^{\aleph'}} \\ &\leq \text{const} |k_0|^{\aleph-\aleph'} |\mathbf{t}|^{\aleph'} \left[ \frac{1}{|ik_0 - E(\mathbf{k}, \mathbf{0}, s)|^2} + \frac{1}{|ik_0 - E(\mathbf{k}, \mathbf{t}, s)|^2} \right]\end{aligned}$$

which is the desired bound.

Using these bounds gives, for  $0 \leq \aleph' < \tilde{\aleph} < \aleph$ ,

$$\begin{aligned}&|B_j(\mathbf{t}, z) - B'_j(\mathbf{t}, z) - B_j(\mathbf{0}, z) + B'_j(\mathbf{0}, z)| \\ &\leq \int dk \int_0^1 ds n_j(k_0) \left| v_j(k, \mathbf{t}, z) I(k_0, \mathbf{k}, \mathbf{t}, s) - v_j(k, \mathbf{0}, z) I(k_0, \mathbf{k}, \mathbf{0}, s) \right| \\ &\leq \text{const} \int dk \int_0^1 ds n_j(k_0) \left[ \frac{|v_j(k, \mathbf{t}, z) - v_j(k, \mathbf{0}, z)| |k_0|^{\aleph}}{|ik_0 - E(\mathbf{k}, \mathbf{t}, s)|^2} + \frac{|v_j(k, \mathbf{0}, z)| |k_0|^{\aleph-\aleph'} |\mathbf{t}|^{\aleph'}}{|ik_0 - E(\mathbf{k}, \mathbf{0}, s)|^2} + \frac{|v_j(k, \mathbf{0}, z)| |k_0|^{\aleph-\aleph'} |\mathbf{t}|^{\aleph'}}{|ik_0 - E(\mathbf{k}, \mathbf{t}, s)|^2} \right] \\ &\leq \text{const} \int_I d\theta \int_{-\text{const}}^{\text{const}} dk_0 \int_{-\text{const}}^{\text{const}} dE \left[ \frac{|k_0|^{\aleph} \sup_k |v_j(k, \mathbf{t}, z) - v_j(k, \mathbf{0}, z)|}{|ik_0 - E|^2} + \frac{|k_0|^{\aleph-\aleph'} |\mathbf{t}|^{\aleph'} \sup_k |v_j(k, \mathbf{0}, z)|}{|ik_0 - E|^2} \right] \\ &\leq \text{const} \mathfrak{l} \left[ \sup_k |v_j(k, \mathbf{t}, z) - v_j(k, \mathbf{0}, z)| + |\mathbf{t}|^{\aleph'} \sup_k |v_j(k, \mathbf{0}, z)| \right] \\ &\leq \text{const} \mathfrak{l} V_{\tilde{\aleph}} |\mathbf{t}|^{\aleph'}\end{aligned}$$

and

$$\begin{aligned}&|B_j(\mathbf{t}, z) - B'_j(\mathbf{t}, z) - B_j(\mathbf{t}, z') + B'_j(\mathbf{t}, z')| \\ &\leq \int dk \int_0^1 ds n_j(k_0) \left| v_j(k, \mathbf{t}, z) I(k_0, \mathbf{k}, \mathbf{t}, s) - v_j(k, \mathbf{t}, z') I(k_0, \mathbf{k}, \mathbf{t}, s) \right| \\ &\leq \text{const} \int dk \int_0^1 ds n_j(k_0) \frac{|v_j(k, \mathbf{t}, z) - v_j(k, \mathbf{t}, z')| |k_0|^{\aleph}}{|ik_0 - E(\mathbf{k}, \mathbf{t}, s)|^2} \\ &\leq \text{const} \int_I d\theta \int_{-\text{const}}^{\text{const}} dk_0 \int_{-\text{const}}^{\text{const}} dE \frac{|k_0|^{\aleph} \sup_k |v_j(k, \mathbf{t}, z) - v_j(k, \mathbf{t}, z')|}{|ik_0 - E|^2} \\ &\leq \text{const} \mathfrak{l} \sup_k |v_j(k, \mathbf{t}, z) - v_j(k, \mathbf{t}, z')| \\ &\leq \text{const} \mathfrak{l} V_{\tilde{\aleph}} |z - z'|^{\aleph'}\end{aligned}$$

Hence it suffices to consider  $B'_j(\mathbf{t}, z)$ .

Making, as in Lemma B.1, for each fixed  $s$ , the change of variables from  $\mathbf{k}$  to  $E = E(\mathbf{k}, \mathbf{t}, s)$  and an “angular” variable  $\theta$

$$B'_j(\mathbf{t}, z) = \int_0^1 ds \int d\theta \int dk_0 dE n_j(k_0) \frac{v_j((k_0, \mathbf{k}(E, \mathbf{t}, \theta, s)), \mathbf{t}, z) J(E, \mathbf{t}, \theta, s)}{[iw(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2}$$

Set

$$B_j''(\mathbf{t}, z) = \int_0^1 ds \int d\theta \int dk_0 dE n_j(k_0) \frac{v_j((k_0, \mathbf{k}(0, \mathbf{t}, \theta, s)), \mathbf{t}, z) J(0, \mathbf{t}, \theta, s)}{[w(\mathbf{k}(0, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2}$$

Using

$$\partial_E \mathbf{k}_\ell(E, \mathbf{t}, \theta, s) = \frac{\partial_{\mathbf{k}_2} \theta(\mathbf{k}) \delta_{\ell,1} - \partial_{\mathbf{k}_1} \theta(\mathbf{k}) \delta_{\ell,2}}{\partial_{\mathbf{k}_1} E(\mathbf{k}, \mathbf{t}, s) \partial_{\mathbf{k}_2} \theta(\mathbf{k}) - \partial_{\mathbf{k}_2} E(\mathbf{k}, \mathbf{t}, s) \partial_{\mathbf{k}_1} \theta(\mathbf{k})} \Big|_{\mathbf{k}=\mathbf{k}(E, \mathbf{t}, \theta, s)}$$

one proves, by induction on  $n$ , that, for  $n \leq r$ ,

$$|\partial_E^n \mathbf{k}(E, \mathbf{t}, \theta, s)| \leq \text{const}'' \quad (\text{B.15})$$

Using this bound and the observation that the Jacobian

$$J(E, \mathbf{t}, \theta, s) = \frac{1}{|\partial_{\mathbf{k}_1} E(\mathbf{k}, \mathbf{t}, s) \partial_{\mathbf{k}_2} \theta(\mathbf{k}) - \partial_{\mathbf{k}_2} E(\mathbf{k}, \mathbf{t}, s) \partial_{\mathbf{k}_1} \theta(\mathbf{k})|} \Big|_{\mathbf{k}=\mathbf{k}(E, \mathbf{t}, \theta, s)}$$

one proves that, for  $n \leq r - 1$ ,

$$|\partial_E^n J(E, \mathbf{t}, \theta, s)| \leq \text{const}'' \quad (\text{B.16})$$

Since

$$\begin{aligned} |w(\mathbf{k}, \mathbf{t}, s) - w(\mathbf{k}', \mathbf{t}, s)| &\leq s \left| \frac{\partial e'}{\partial k_0}(0, \mathbf{k}) - \frac{\partial e'}{\partial k_0}(0, \mathbf{k}') \right| + (1-s) \left| \frac{\partial e'}{\partial k_0}(0, \mathbf{k} + \mathbf{t}) - \frac{\partial e'}{\partial k_0}(0, \mathbf{k}' + \mathbf{t}) \right| \\ &\leq \text{const} |\mathbf{k} - \mathbf{k}'|^\aleph \end{aligned} \quad (\text{B.17})$$

(B.15), (B.16) and the fact, from Lemma A.1.i, that  $|w(\mathbf{k}, \mathbf{t}, s) - 1| \leq \rho \leq \frac{1}{2}$  imply that

$$\begin{aligned} &\left| \frac{v_j((k_0, \mathbf{k}(E, \mathbf{t}, \theta, s)), \mathbf{t}, z) J(E, \mathbf{t}, \theta, s)}{[w(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2} - \frac{v_j((k_0, \mathbf{k}(0, \mathbf{t}, \theta, s)), \mathbf{t}, z) J(0, \mathbf{t}, \theta, s)}{[w(\mathbf{k}(0, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2} \right| \\ &\leq \text{const} \left| v_j((k_0, \mathbf{k}(E, \mathbf{t}, \theta, s)), \mathbf{t}, z) - v_j((k_0, \mathbf{k}(0, \mathbf{t}, \theta, s)), \mathbf{t}, z) \right| \frac{1}{k_0^2 + E^2} \\ &\quad + \text{const} \sup_{k, \mathbf{t}} |v_j(k, \mathbf{t}, z)| \frac{|E|}{k_0^2 + E^2} + \text{const} \sup_{k, \mathbf{t}} |v_j(k, \mathbf{t}, z)| \frac{[|k_0| + |E|][k_0][|E|]^\aleph}{[k_0^2 + E^2]^2} \\ &\leq \text{const} \left[ \sup_{\mathbf{k}, \mathbf{k}'} \frac{|v_j((k_0, \mathbf{k}'), \mathbf{t}, z) - v_j((k_0, \mathbf{k}), \mathbf{t}, z)|}{|\mathbf{k} - \mathbf{k}'|^\aleph} + \sup_{k, \mathbf{t}} |v_j(k, \mathbf{t}, z)| \right] \frac{|E|^\aleph}{k_0^2 + E^2} \leq \text{const} V_\aleph \frac{|E|^\aleph}{k_0^2 + E^2} \end{aligned}$$

for all small  $E$ . For  $E$  bounded away from zero

$$\left| \frac{v_j((k_0, \mathbf{k}(E, \mathbf{t}, \theta, s)), \mathbf{t}, z) J(E, \mathbf{t}, \theta, s)}{[w(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2} - \frac{v_j((k_0, \mathbf{k}(0, \mathbf{t}, \theta, s)), \mathbf{t}, z) J(0, \mathbf{t}, \theta, s)}{[w(\mathbf{k}(0, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2} \right| \leq \text{const} \sup_{k, \mathbf{t}} |v_j(k, \mathbf{t}, z)| \frac{1}{k_0^2 + E^2}$$

so

$$\left| \frac{v_j((k_0, \mathbf{k}(E, \mathbf{t}, \theta, s)), \mathbf{t}, z) J(E, \mathbf{t}, \theta, s)}{[w(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2} - \frac{v_j((k_0, \mathbf{k}(0, \mathbf{t}, \theta, s)), \mathbf{t}, z) J(0, \mathbf{t}, \theta, s)}{[w(\mathbf{k}(0, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2} \right| \leq \text{const} V_\aleph \frac{\min\{|E|^\aleph, 1\}}{k_0^2 + E^2} \quad (\text{B.18})$$

Since

$$|w(\mathbf{k}, \mathbf{t}, s) - w(\mathbf{k}, \mathbf{0}, s)| = |1 - s| \left| \frac{\partial e'}{\partial k_0}(0, \mathbf{k} + \mathbf{t}) - \frac{\partial e'}{\partial k_0}(0, \mathbf{k}) \right| \leq \text{const } |\mathbf{t}|^{\aleph}$$

(B.17), (B.4) and (B.5) imply that

$$\begin{aligned} & \left| \frac{v_j((k_0, \mathbf{k}(E, \mathbf{t}, \theta, s)), \mathbf{t}, z) J(E, \mathbf{t}, \theta, s)}{[i w(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2} - \frac{v_j((k_0, \mathbf{k}(E, \mathbf{0}, \theta, s)), \mathbf{0}, z) J(E, \mathbf{0}, \theta, s)}{[i w(\mathbf{k}(E, \mathbf{0}, \theta, s), \mathbf{0}, s) k_0 - E]^2} \right| \\ & \leq \text{const } \left| v_j((k_0, \mathbf{k}(E, \mathbf{t}, \theta, s)), \mathbf{t}, z) - v_j((k_0, \mathbf{k}(E, \mathbf{0}, \theta, s)), \mathbf{0}, z) \right| \frac{1}{k_0^2 + E^2} \\ & \quad + \text{const } |\mathbf{t}| \sup_k |v_j(k, \mathbf{0}, z)| \frac{1}{k_0^2 + E^2} + \text{const } |\mathbf{t}|^{\aleph} \sup_k |v_j(k, \mathbf{0}, z)| \frac{|k_0| [|k_0| + |E|]}{[k_0^2 + E^2]^2} \\ & \leq \text{const } \left[ \sup_{\substack{\mathbf{k}, \mathbf{k}' \\ |\mathbf{k} - \mathbf{k}'| \leq \text{const } |\mathbf{t}|}} |v_j((k_0, \mathbf{k}'), \mathbf{t}, z) - v_j((k_0, \mathbf{k}), \mathbf{0}, z)| + |\mathbf{t}|^{\aleph} \sup_{\mathbf{k}} |v_j(k, \mathbf{0}, z)| \right] \frac{1}{k_0^2 + E^2} \\ & \leq \text{const } V_{\tilde{\aleph}} |\mathbf{t}|^{\tilde{\aleph}} \frac{1}{k_0^2 + E^2} \end{aligned} \tag{B.19}$$

and that

$$\begin{aligned} & \left| \frac{v_j((k_0, \mathbf{k}(0, \mathbf{t}, \theta, s)), \mathbf{t}, z) J(0, \mathbf{t}, \theta, s)}{[i w(\mathbf{k}(0, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2} - \frac{v_j((k_0, \mathbf{k}(0, \mathbf{0}, \theta, s)), \mathbf{0}, z) J(0, \mathbf{0}, \theta, s)}{[i w(\mathbf{k}(0, \mathbf{0}, \theta, s), \mathbf{0}, s) k_0 - E]^2} \right| \\ & \leq \text{const } \left| v_j((k_0, \mathbf{k}(0, \mathbf{t}, \theta, s)), \mathbf{t}, z) - v_j((k_0, \mathbf{k}(0, \mathbf{0}, \theta, s)), \mathbf{0}, z) \right| \frac{1}{k_0^2 + E^2} \\ & \quad + \text{const } |\mathbf{t}| \sup_k |v_j(k, \mathbf{0}, z)| \frac{1}{k_0^2 + E^2} + \text{const } |\mathbf{t}|^{\aleph} \sup_k |v_j(k, \mathbf{0}, z)| \frac{|k_0| [|k_0| + |E|]}{[k_0^2 + E^2]^2} \\ & \leq \text{const } V_{\tilde{\aleph}} |\mathbf{t}|^{\tilde{\aleph}} \frac{1}{k_0^2 + E^2} \end{aligned} \tag{B.20}$$

Combining, (B.18), a second copy of (B.18) with  $\mathbf{t} = \mathbf{0}$ , (B.19) and (B.20), gives, for all  $0 \leq \aleph' < \tilde{\aleph} < \aleph$ ,

$$\begin{aligned} & \left| \frac{v_j((k_0, \mathbf{k}(E, \mathbf{t}', \theta, s)), \mathbf{t}', z) J(E, \mathbf{t}', \theta, s)}{[i w(\mathbf{k}(E, \mathbf{t}', \theta, s), \mathbf{t}', s) k_0 - E]^2} \right|_{\substack{\mathbf{t}' = \mathbf{t} \\ \mathbf{t}' = \mathbf{0}}} - \frac{v_j((k_0, \mathbf{k}(0, \mathbf{t}', \theta, s)), \mathbf{t}', z) J(0, \mathbf{t}', \theta, s)}{[i w(\mathbf{k}(0, \mathbf{t}', \theta, s), \mathbf{t}', s) k_0 - E]^2} \right|_{\substack{\mathbf{t}' = \mathbf{t} \\ \mathbf{t}' = \mathbf{0}}} \\ & \leq \text{const } V_{\tilde{\aleph}} |\mathbf{t}|^{\aleph'} \frac{\min\{|E|^{\tilde{\aleph} - \aleph'}, 1\}}{k_0^2 + E^2} \end{aligned}$$

and

$$\begin{aligned} & |B'_j(\mathbf{t}, z) - B''_j(\mathbf{t}, z) - B'_j(\mathbf{0}, z) + B''_j(\mathbf{0}, z)| \\ & \leq \text{const } V_{\tilde{\aleph}} |\mathbf{t}|^{\aleph'} \int_I d\theta \int_{-\text{const}}^{\text{const}} dk_0 \int dE \frac{\min\{|E|^{\tilde{\aleph} - \aleph'}, 1\}}{k_0^2 + E^2} \\ & \leq \text{const } V_{\tilde{\aleph}} |\mathbf{t}|^{\aleph'} \end{aligned}$$

Similarly, combining, (B.18), a second copy of (B.18) with  $z \rightarrow z'$  and

$$\left| \frac{v_j((k_0, \mathbf{k}(E', \mathbf{t}, \theta, s)), \mathbf{t}, z) J(E', \mathbf{t}, \theta, s)}{[i w(\mathbf{k}(E', \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2} - \frac{v_j((k_0, \mathbf{k}(E', \mathbf{t}, \theta, s)), \mathbf{t}, z') J(E', \mathbf{t}, \theta, s)}{[i w(\mathbf{k}(E', \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2} \right| \leq \text{const } V_{\tilde{\aleph}} |z - z'|^{\tilde{\aleph}} \frac{1}{k_0^2 + E^2}$$

gives, for all  $0 \leq \aleph' < \tilde{\aleph} < \aleph$ ,

$$\begin{aligned} & \left| \frac{v_j((k_0, \mathbf{k}(E, \mathbf{t}, \theta, s)), \mathbf{t}, \tilde{z}) J(E, \mathbf{t}, \theta, s)}{[i w(\mathbf{k}(E, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2} \right|_{\substack{\tilde{z} = z \\ \tilde{z} = z'}} - \frac{v_j((k_0, \mathbf{k}(0, \mathbf{t}, \theta, s)), \mathbf{t}, \tilde{z}) J(0, \mathbf{t}, \theta, s)}{[i w(\mathbf{k}(0, \mathbf{t}, \theta, s), \mathbf{t}, s) k_0 - E]^2} \right|_{\substack{\tilde{z} = z \\ \tilde{z} = z'}} \\ & \leq \text{const } V_{\tilde{\aleph}} |z - z'|^{\aleph'} \frac{\min\{|E|^{\tilde{\aleph} - \aleph'}, 1\}}{k_0^2 + E^2} \end{aligned}$$

and

$$\begin{aligned}
& |B'_j(\mathbf{t}, z) - B''_j(\mathbf{t}, z) - B'_j(\mathbf{t}, z') + B''_j(\mathbf{t}, z')| \\
& \leq \text{const } V_{\tilde{\mathbf{R}}} |z - z'|^{\aleph'} \int_I d\theta \int_{-\text{const}}^{\text{const}} dk_0 \int dE \frac{\min\{|E|^{\tilde{\mathbf{R}} - \aleph'}, 1\}}{k_0^2 + E^2} \\
& \leq \text{const } \mathfrak{I} V_{\tilde{\mathbf{R}}} |z - z'|^{\aleph'}
\end{aligned}$$

Fix  $s$ ,  $\mathbf{t}$  and  $\theta$  and write  $w$  for  $w(\mathbf{k}(0, \mathbf{t}, \theta, s), \mathbf{t}, s)$ . Then, for all  $k_0 \neq 0$ ,

$$\int_{-\infty}^{\infty} dE \frac{1}{[iwk_0 - E]^2} = \int_{-\infty}^{\infty} dE \frac{d}{dE} \frac{1}{iwk_0 - E} = 0$$

Thus  $B''_j(\mathbf{t}, z) \equiv 0$ .

d) By part (a), it suffices to prove that  $B_j(\mathbf{t}, z)$  converges as  $j \rightarrow \infty$ . By (B.14),  $\sup_j |v_j(k, \mathbf{t}, z)I(k_0, \mathbf{k}, \mathbf{t}, s)|$  is locally  $L^1$  in  $k$  and  $s$ . Hence, by the Lebesgue dominated convergence theorem applied to the integral in (B.13),  $B_j(\mathbf{t}, z) - B'_j(\mathbf{t}, z)$  converges as  $j \rightarrow \infty$ . So it suffices to prove that  $B'_j(\mathbf{t}, z)$  converges. By (B.18), the Lebesgue dominated convergence theorem also implies that  $B'_j(\mathbf{t}, z) - B''_j(\mathbf{t}, z)$  converges as  $j \rightarrow \infty$ . We have already observed that  $B''_j(\mathbf{t}, z) \equiv 0$ . ■

The function  $A(t_0, z)$  of Lemma B.3.b was constructed in such a way that the cutoff in the  $k_0$  direction was removed first (it does not even appear in the definition of  $A_j(t_0, z)$ ) and the cutoff in the  $e(\mathbf{k})$  direction was removed second (in the limit  $j \rightarrow \infty$ ). On the other hand, for the function  $B(\mathbf{t}, z)$  of Lemma B.3.d, the cutoff in the  $e(\mathbf{k})$  direction was removed before the cutoff in the  $k_0$  direction. The following Lemma illustrates, in a simplified setting, that the order of removal of the two cutoffs matters when  $t = 0$ .

**Lemma B.4** *Let, for  $0 < a, b < 1$  and  $w > 0$ ,*

$$B_{a,b}^{(1)} = \int_{a < |k_0| \leq 1} dk_0 \int_{b < |E| \leq 1} dE \frac{1}{[iwk_0 - E]^2}$$

*Then  $B_{a,b}^{(1)}$  is bounded uniformly on  $w \geq \epsilon > 0$ ,  $0 < a, b < 1$  and*

$$\lim_{a \rightarrow 0} \lim_{b \rightarrow 0} B_{a,b}^{(1)} = -\frac{4}{w} \tan^{-1} w \qquad \lim_{b \rightarrow 0} \lim_{a \rightarrow 0} B_{a,b}^{(1)} = -\frac{4}{w} \left[ \tan^{-1} w - \frac{\pi}{2} \right]$$

**Proof:**

$$\begin{aligned}
B_{a,b}^{(1)} &= \int_{a < |k_0| \leq 1} dk_0 \left[ \frac{1}{iwk_0 - E} \Big|_b^1 + \frac{1}{iwk_0 - E} \Big|_{-1}^{-b} \right] = -2 \int_{a < |k_0| \leq 1} dk_0 \left[ \frac{1}{1+w^2k_0^2} - \frac{b}{b^2+w^2k_0^2} \right] \\
&= -4 \int_a^1 dk_0 \left[ \frac{1}{1+w^2k_0^2} - \frac{b}{b^2+w^2k_0^2} \right] = -\frac{4}{w} \int_{wa}^w dx \frac{1}{1+x^2} + \frac{4}{w} \int_{wa/b}^{w/b} dx \frac{b}{b^2+x^2} \\
&= -\frac{4}{w} \int_{wa}^w dx \frac{1}{1+x^2} + \frac{4}{w} \int_{wa/b}^{w/b} dy \frac{1}{1+y^2} \\
&= -\frac{4}{w} \left[ \tan^{-1} w - \tan^{-1} wa - \tan^{-1} \frac{w}{b} + \tan^{-1} \frac{wa}{b} \right]
\end{aligned}$$


■

## Appendix C: Sector Counting with Specified Transfer Momentum

As pointed out in the introduction to §IV, we are interested in translates  $F + \mathbf{t} = \{ \mathbf{k} + \mathbf{t} \mid \mathbf{k} \in F \}$  of the Fermi surface  $F$ , and in particular in the distances from points of  $F + \mathbf{t}$  to  $F$ .

**Lemma C.1** *There are constants  $\delta, \text{const} > 0$  that depend only on the Fermi curve  $F$  such that the following holds: Let  $\mathbf{p} \in F$ ,  $|\mathbf{t}| \leq \delta$  such that  $\mathbf{p} - \mathbf{t} \in F$ . Denote by  $U$  the disc of radius  $\delta$  around  $\mathbf{p}$ . Then for any  $\mathbf{k} \in F \cap U$ .*


$$|\mathbf{k} - \mathbf{p}| \leq \frac{\text{const}}{|\mathbf{t}|} \text{dist}(\mathbf{k}, F + \mathbf{t})$$



**Proof:** If  $\text{dist}(\mathbf{k}, F + \mathbf{t}) \geq \frac{1}{2}|\mathbf{t}|$  or if  $\mathbf{k} = \mathbf{p}$  there is nothing to prove. So assume that  $\text{dist}(\mathbf{k}, F + \mathbf{t}) \leq \frac{1}{2}|\mathbf{t}|$ . If  $\delta$  was chosen small enough, the angle between the chords  $-\mathbf{t} = (\mathbf{p} - \mathbf{t}) - \mathbf{p}$  and  $\mathbf{k} - \mathbf{p}$  of  $F$  is sufficiently small. In particular,  $(\mathbf{k} - \mathbf{p}) \cdot \mathbf{t} \neq 0$ .

*Case 1:  $(\mathbf{k} - \mathbf{p}) \cdot \mathbf{t} > 0$*

We may assume without loss of generality that  $\mathbf{p} = (0, 0)$ , that  $\mathbf{t} = (0, t_2)$  with  $t_2 > 0$  and that the tangent direction of  $F$  at  $\mathbf{p}$  is  $(\alpha, 1)$  with some  $\alpha > 0$ . As  $F$  is strictly convex, and both  $\mathbf{p} = (0, 0)$  and  $\mathbf{t} = (0, t_2)$  are on  $F$ ,  $F' = \{ \mathbf{k}' \in F \cap U \mid (\mathbf{k}' - \mathbf{p}) \cdot \mathbf{t} \geq 0 \}$  is contained in the first quadrant, if  $\delta$  was chosen small enough. By the implicit function theorem,  $F'$  can be parametrized in the form  $F' = \{ (x, y(x)) \mid 0 \leq x < \text{const} \}$  with an  $r_e + 3$  times differentiable function satisfying  $y'(0) = \frac{1}{\alpha}$ ,  $y'' < 0$ .




Since the curvature of  $F$  is bounded above and below, there are constants  $\text{const}_1, \text{const}_2 > 0$  such that

$$\text{const}_1 |\mathbf{t}| \leq \alpha \leq \text{const}_2 |\mathbf{t}|$$

If  $\delta$  was chosen small enough,  $y' > 1$ . Let  $c_1$  resp.  $c_2$  be the maximal resp. minimal curvature of  $F$ , and let  $C_1$  resp.  $C_2$  be the circles of curvature  $c_1$  resp.  $c_2$  that are tangent to  $F$  at  $\mathbf{p}$

and curved in the same direction as  $F$  at  $\mathbf{p}$ . Then  $F'$  lies between  $C_1$  and  $C_2$ , and the slope of  $F'$  at a point  $(x, y(x))$  lies between the slopes of  $C_1$  resp.  $C_2$  at the points with the same  $x$ -coordinate in the first quadrant.




If  $C$  is a circle of a radius  $r > 0$  that is tangent to  $F$  at  $\mathbf{p}$  and curved in the same direction as  $F$  at  $\mathbf{p}$  then the slope of  $C$  at any of its points  $(x, y)$  in the first quadrant is equal to

$$\frac{\frac{r}{\sqrt{1+\alpha^2}} - x}{y + \frac{\alpha r}{\sqrt{1+\alpha^2}}}$$

Therefore, for any point  $(x, y(x))$  of  $F'$

$$y'(x) \leq \frac{\frac{1}{c_2\sqrt{1+\alpha^2}} - x}{y(x) + \frac{\alpha}{c_2\sqrt{1+\alpha^2}}} \leq \frac{const_3}{y(x) + const_4|\mathbf{t}|}$$

Let  $F''$  be the union of  $\mathbf{t} + F'$  and the segment joining  $\mathbf{p} = (0, 0)$  to  $\mathbf{t}$ .



Then


$$\text{dist}(\mathbf{k}, F + \mathbf{t}) \geq \text{dist}(\mathbf{k}, F'') \geq \min \{k_1, \text{dist}(\mathbf{k}, F' + \mathbf{t})\}$$

As  $k_1 \geq \alpha k_2 \geq const|\mathbf{t}||\mathbf{k}|$ , we get that

$$\frac{const}{|\mathbf{t}|} \text{dist}(\mathbf{k}, F + \mathbf{t}) \geq \min \{|\mathbf{k} - \mathbf{p}|, \frac{1}{|\mathbf{t}|} \text{dist}(\mathbf{k}, F' + \mathbf{t})\} \quad (\text{C.1})$$

If  $k_2 \leq |\mathbf{t}|$  then the distance from  $\mathbf{k}$  to  $F' + \mathbf{t}$  is larger than the distance from  $\mathbf{k}$  to the ray through  $\mathbf{t}$  in the direction  $(1, 1)$ , since  $y' > 1$ .






Consequently

$$\text{dist}(\mathbf{k}, F' + \mathbf{t}) \geq \frac{1}{\sqrt{2}}k_1 \geq \text{const}|\mathbf{t}| |\mathbf{k} - \mathbf{p}|$$

Together with (C.1) this gives the claim of the Lemma in the situation that  $k_2 \leq |\mathbf{t}|$ .

Now assume that  $k_2 \geq |\mathbf{t}|$ . Let  $\mathbf{k}' = (k'_1, k_2 - |\mathbf{t}|)$  be the point of  $F'$  with  $y$ -coordinate  $k'_2 = k_2 - |\mathbf{t}|$  that lies to the left of  $\mathbf{k}$ , i.e.  $k'_1 < k_1$ .



By the convexity of  $F$ , the distance of  $\mathbf{k}$  to  $\mathbf{t} + F$  is bounded below by the distance of  $\mathbf{k}$  to the line segment joining  $\mathbf{k}' + \mathbf{t}$  and  $\mathbf{k} + \mathbf{t}$ . Thus

$$\text{dist}(\mathbf{k}, F' + \mathbf{t}) \geq \frac{1}{\sqrt{2}} \min\{|\mathbf{t}|, k_1 - k'_1\}$$

Since  $F'$  is strictly convex

$$k_1 - k'_1 \geq \frac{|\mathbf{t}|}{y'(k'_1)} \geq \text{const}_5 |\mathbf{t}| (y(k'_1) + \text{const}_4 |\mathbf{t}|) = \text{const}_5 |\mathbf{t}| (k'_2 + \text{const}_4 |\mathbf{t}|)$$

If  $k_2 \leq 2|\mathbf{t}|$  then  $|\mathbf{t}| \geq \text{const} |\mathbf{k}| = \text{const} |\mathbf{k} - \mathbf{p}|$  and

$$k_1 - k'_1 \geq \text{const} |\mathbf{t}|^2 \geq \text{const} |\mathbf{t}| |\mathbf{k} - \mathbf{p}|$$

and if  $k_2 \geq 2|\mathbf{t}|$

$$k_1 - k'_1 \geq \text{const} |\mathbf{t}| k'_2 \geq \text{const} |\mathbf{t}| k_2 \geq \text{const} |\mathbf{t}| |\mathbf{k}| = \text{const} |\mathbf{t}| |\mathbf{k} - \mathbf{p}|$$

Therefore

$$\frac{1}{|\mathbf{t}|} \text{dist}(\mathbf{k}, F' + \mathbf{t}) \geq \text{const} \min\{1, |\mathbf{k} - \mathbf{p}|\} \geq \text{const} |\mathbf{k} - \mathbf{p}|$$

Again, (C.1) implies the claim of the Lemma in the situation that  $k_2 \geq |\mathbf{t}|$ .

Case 2:  $(\mathbf{k} - \mathbf{p}) \cdot \mathbf{t} < 0$


Let  $\mathbf{k}' \in F + \mathbf{t}$  such that  $\text{dist}(\mathbf{k}, F + \mathbf{t}) = |\mathbf{k} - \mathbf{k}'|$ . Then

$$\text{dist}(\mathbf{k}', (F + \mathbf{t}) - \mathbf{t}) \leq |\mathbf{k}' - \mathbf{k}| = \text{dist}(\mathbf{k}, F + \mathbf{t})$$

and

$$|\mathbf{k} - \mathbf{p}| \leq |\mathbf{k} - \mathbf{k}'| + |\mathbf{k}' - \mathbf{p}| \leq \text{dist}(\mathbf{k}, F + \mathbf{t}) + |\mathbf{k}' - \mathbf{p}| \quad (\text{C.2})$$

Observe that the point  $\mathbf{p}$  of  $F + \mathbf{t}$  has the property that  $\mathbf{p} - (-\mathbf{t})$  lies also in  $F + \mathbf{t}$ . Also, if  $\delta$  was chosen small enough, the angle between the tangents to  $F$  at  $\mathbf{p}$  and to  $F + \mathbf{t}$  at  $\mathbf{p}$  (which is parallel to the tangent to  $F$  at  $\mathbf{p} - \mathbf{t}$ ) is very small and  $(\mathbf{k}' - \mathbf{p}) \cdot (-\mathbf{t}) > 0$ .



Thus we can apply the results of Case 1, with  $F$  replaced by  $F + \mathbf{t}$ ,  $\mathbf{t}$  replaced by  $-\mathbf{t}$  and  $\mathbf{k}$  replaced by  $\mathbf{k}'$  and get

$$|\mathbf{k}' - \mathbf{p}| \leq \frac{\text{const}}{|\mathbf{t}|} \text{dist}(\mathbf{k}', (F + \mathbf{t}) - \mathbf{t}) \leq \frac{\text{const}}{|\mathbf{t}|} \text{dist}(\mathbf{k}, F + \mathbf{t})$$

This, together with (C.2), proves the Lemma in Case 2. ■

**Lemma C.2** *There are constants  $\delta_F, \text{const}$  that depend only on  $F$  and  $M$  such that the following holds:*

*Let  $\boldsymbol{\tau} \in \mathbb{R}^2$ ,  $\epsilon > 0$  and  $D$  the disc centered at  $\boldsymbol{\tau}$  with radius  $\epsilon$ . Let  $m \geq 1$  be a scale with  $\mathfrak{l}_m \geq \frac{1}{2}\epsilon$ . Define*

$$N = \#\{ (s_1, s_2) \in \Sigma_m \times \Sigma_m \mid (s_1 - s_2) \cap D \neq \emptyset \}$$

*where  $D \subset \mathbb{R}^2$  is viewed as  $\{ (0, \mathbf{t}) \mid \mathbf{t} \in D \} \subset \mathbb{R} \times \mathbb{R}^2$ .*

*a) If  $|\boldsymbol{\tau}| \geq \delta_F$ , then  $N \leq \frac{\text{const}}{\sqrt{\mathfrak{l}_m}}$ .*

*b) If  $|\boldsymbol{\tau}| \leq \delta_F$ , then*

$$N \leq \frac{\text{const}}{\mathfrak{l}_m |\boldsymbol{\tau}|} \left( \frac{1}{M^m} + \epsilon \right) + \text{const}$$

**Proof:** We first observe that, given any fixed  $s_1 \in \Sigma_m$ ,  $(s_1 - s_2) \cap D \neq \emptyset$  only if  $s_2 \cap (s_1 - D) \neq \emptyset$ . As  $s_1 - D$  is contained in a ball of radius at most  $3\mathfrak{l}_m$ , there are at most

five sectors  $s_2 \in \Sigma_m$  that intersect it. Hence

$$\begin{aligned} N &\leq \text{const} \# \{ s_1 \in \Sigma_m \mid \exists s_2 \in \Sigma_m \text{ such that } s_2 \cap (s_1 - D) \neq \emptyset \} \\ &\leq \text{const} \# \{ s_1 \in \Sigma_m \mid \exists \mathbf{k} \in s_1 \cap F \text{ such that } \text{dist}(\mathbf{k} - \boldsymbol{\tau}, F) \leq \text{const}_1 \left( \frac{1}{M^m} + \epsilon \right) \} \end{aligned}$$

Define

$$I = \{ \mathbf{k} \in F \mid \text{dist}(\mathbf{k} - \boldsymbol{\tau}, F) \leq \text{const}_1 \left( \frac{1}{M^m} + \epsilon \right) \}$$

Then

$$N \leq \text{const} \# \{ s \in \Sigma_m \mid s \cap I \neq \emptyset \} \quad (\text{C.3})$$

Clearly  $I \subset I'$ , where

$$I' = \{ \mathbf{k} \in F \mid \text{dist}(\mathbf{k}, F + \boldsymbol{\tau}) \leq \text{const} \, \mathfrak{l}_m \}$$


and hence

$$N \leq \text{const} \# \{ s \in \Sigma_m \mid s \cap I' \neq \emptyset \} \leq \frac{\text{const}}{\mathfrak{l}_m} \text{length}(I') + \text{const} \quad (\text{C.4})$$

We choose  $\delta_F$  to be smaller than the constant  $\delta$  of Lemma C.1.

a) If  $|\boldsymbol{\tau}| \geq \delta_F$ , we use (C.4) and that

$$\text{length}(I') \leq \text{const} \sqrt{\mathfrak{l}_m}$$



b) Assume that  $|\boldsymbol{\tau}| \leq \delta_F$ . Since  $F$  is strictly convex,  $F \cap (F + \boldsymbol{\tau})$  consists of two points, say  $\mathbf{p}_1, \mathbf{p}_2$ . Let  $U_1$  and  $U_2$  be the discs of radius  $\delta_F$  around  $\mathbf{p}_1$  and  $\mathbf{p}_2$ , respectively. If  $\delta_F$  is small enough,  $U_1$  and  $U_2$  are disjoint. By Lemma C.1, for  $i = 1, 2$ ,  $U_i \cap I$  is contained in an interval of length  $\frac{\text{const}}{|\boldsymbol{\tau}|} \left( \frac{1}{M^m} + \epsilon \right)$ . Also by Lemma C.1, for  $i = 1, 2$ , the distance between  $F + \boldsymbol{\tau}$  and any endpoint of  $F \cap U_i$  is bigger than  $\text{const} \, \delta_F |\boldsymbol{\tau}|$ . If  $\frac{1}{|\boldsymbol{\tau}|} \left( \frac{1}{M^m} + \epsilon \right) \geq \frac{\text{const}}{2\text{const}_1} \delta_F$ , the desired bound follows immediately from (C.3) and the fact that  $\#\Sigma_m \leq \frac{\text{const}}{\mathfrak{l}_m}$ . So we may assume that, for  $i = 1, 2$  the distance between  $F + \boldsymbol{\tau}$  and any endpoint of  $F \cap U_i$  is bigger than  $2\text{const}_1 \left( \frac{1}{M^m} + \epsilon \right)$ .

We now show that  $I \subset U_1 \cup U_2$ . For this purpose, let  $\mathbf{k} \in I$ . Then there is  $\mathbf{v} \in \mathbb{R}^2$  with  $|\mathbf{v}| \leq \text{const}_1 \left( \frac{1}{M^m} + \epsilon \right)$  such that  $\mathbf{k} \in F + \boldsymbol{\tau} + \mathbf{v}$ . Now for  $i = 1, 2$  there is point  $\mathbf{k}_i \in F \cap U_i$

such that  $\mathbf{k}_i - \mathbf{v} \in F + \boldsymbol{\tau}$ . (At the two endpoints  $\mathbf{k}'$  of  $F \cap U_i$ , the points  $\mathbf{k}' - \mathbf{v}$  lie on opposite sides of  $F + \boldsymbol{\tau}$ .) Since  $F \cap (F + \boldsymbol{\tau} + \mathbf{v})$  consists of only two points,  $\mathbf{k} = \mathbf{k}_1$  or  $\mathbf{k} = \mathbf{k}_2$ ; in particular  $\mathbf{k} \in U_1 \cup U_2$ .

Therefore  $I$  is contained in two intervals of length  $\frac{\text{const}}{|\boldsymbol{\tau}|} \left( \frac{1}{M^m} + \epsilon \right)$  and, by (C.3),

$$N \leq \frac{\text{const}}{\mathfrak{l}_m} \frac{\text{const}}{|\boldsymbol{\tau}|} \left( \frac{1}{M^m} + \epsilon \right) + \text{const}$$

■

Recall that  $\boldsymbol{\pi} : k = (k_0, \mathbf{k}) \mapsto \mathbf{k}$  is the projection of  $\mathbb{M} = \mathbb{R} \times \mathbb{R}^2$  onto its second factor.

**Lemma C.3** *Let  $1 \leq \ell \leq m \leq r$  and  $\kappa' \in \mathfrak{K}_\ell$ ,  $\kappa_1, \kappa_2 \in \mathfrak{K}_r$ . Then the number of 4-tuples  $(u_1, u_2, s_1, s_2) \in \Sigma_\ell \times \Sigma_\ell \times \Sigma_m \times \Sigma_m$  fulfilling*

$$\begin{aligned} \boldsymbol{\pi}(s_1 - s_2) \cap \boldsymbol{\pi}(\kappa_1 - \kappa_2) &\neq \emptyset \\ \boldsymbol{\pi}(u_1 - u_2) \cap \boldsymbol{\pi}(\kappa' - s_1) &\neq \emptyset \end{aligned} \tag{C.5}$$

*is bounded by  $\frac{\text{const}}{\mathfrak{l}_m \sqrt{\mathfrak{l}_\ell}}$  with the constant  $\text{const}$  independent of  $\kappa_1, \kappa_2, \kappa', \ell, m$  and  $r$ .*

**Proof:** Observe that for each fixed  $s_1 \in \Sigma_m$  there are at most  $\text{const}$  sectors  $s_2 \in \Sigma_m$  fulfilling  $\boldsymbol{\pi}(s_1 - s_2) \cap \boldsymbol{\pi}(\kappa_1 - \kappa_2) \neq \emptyset$ . Recall that for any sector  $s$ ,  $\mathbf{k}_s$  denotes the center of  $F \cap s$ . When  $\kappa' \in \mathbb{M}$ , set  $\mathbf{k}_{\kappa'} = \kappa'$ . We bound each of the three terms in

$$\begin{aligned} &\#\{(u_1, u_2, s_1, s_2) \mid \text{(C.5) holds}\} \\ &\leq \#\{(u_1, u_2, s_1, s_2) \mid \text{(C.5) holds, } |\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}| \leq \text{const } \mathfrak{l}_\ell\} \\ &\quad + \#\{(u_1, u_2, s_1, s_2) \mid \text{(C.5) holds, } \text{const } \mathfrak{l}_\ell \leq |\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}| \leq \delta_F\} \\ &\quad + \#\{(u_1, u_2, s_1, s_2) \mid \text{(C.5) holds, } \delta_F \leq |\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}|\} \end{aligned}$$

separately. For the first term observe that there are at most  $\text{const} \left[ \frac{\mathfrak{l}_\ell}{\mathfrak{l}_m} + 1 \right]$  sectors  $s_1$  with  $|\mathbf{k}_{s_1} - \mathbf{k}_{\kappa'}| \leq \text{const } \mathfrak{l}_\ell$ , and that for any given  $s_1$ , there are at most  $\frac{\text{const}}{\mathfrak{l}_\ell}$  pairs  $(u_1, u_2)$  such that  $\boldsymbol{\pi}(u_1 - u_2) \cap \boldsymbol{\pi}(\kappa' - s_1) \neq \emptyset$ . Hence

$$\#\{(u_1, u_2, s_1, s_2) \mid \text{(C.5) holds, } |\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}| \leq \text{const } \mathfrak{l}_\ell\} \leq \text{const} \left[ \frac{\mathfrak{l}_\ell}{\mathfrak{l}_m} + 1 \right] \frac{1}{\mathfrak{l}_\ell} \leq \frac{\text{const}}{\mathfrak{l}_m}$$

We next bound the third term. There are at most  $\frac{\text{const}}{\mathfrak{l}_m}$  pairs  $(s_1, s_2)$  obeying  $\boldsymbol{\pi}(s_1 - s_2) \cap \boldsymbol{\pi}(\kappa_1 - \kappa_2) \neq \emptyset$  and  $|\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}| \geq \delta_F$ . For each such a pair  $(s_1, s_2)$ ,  $\boldsymbol{\pi}(\kappa' - s_1)$  is contained in a disc of radius  $2\mathfrak{l}_\ell$ , centered a distance at least  $\delta_F$  from the origin, so, by Lemma C.2a, with  $m$  replaced by  $\ell$ , there are at most  $\frac{\text{const}}{\sqrt{\mathfrak{l}_\ell}}$  pairs  $(u_1, u_2)$  such that (C.5) holds. Hence

$$\#\{(u_1, u_2, s_1, s_2) \mid \text{(C.5) holds, } \delta_F \leq |\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}|\} \leq \frac{\text{const}}{\mathfrak{l}_m \sqrt{\mathfrak{l}_\ell}}$$

Finally, for the second term, we observe that, for each fixed  $(s_1, s_2)$  satisfying  $\text{const } \mathfrak{l}_\ell \leq |\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}| \leq \delta_F$  there are, by Lemma C.2b with  $\epsilon = 2\mathfrak{l}_\ell$  and  $m$  replaced by  $\ell$ , at most  $\frac{\text{const}}{\mathfrak{l}_\ell |\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}|} \left[ \frac{1}{M^\ell} + \mathfrak{l}_\ell \right] + \text{const} \leq \frac{\text{const}}{|\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}|}$  pairs  $(u_1, u_2)$  such that (C.5) holds. Furthermore, we may order the allowed  $s_1$ 's so that the  $\mu^{\text{th}}$  obeys  $|\mathbf{k}_{s_1} - \mathbf{k}_{\kappa'}| \geq \text{const} (\mathfrak{l}_\ell + \mu \mathfrak{l}_m)$ . Hence

$$\begin{aligned} \#\{(u_1, u_2, s_1, s_2) \mid (\text{C.5}) \text{ holds, } \text{const } \mathfrak{l}_\ell \leq |\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}| \leq \delta_F\} &\leq \sum_{\mu=1}^{\text{const}/\mathfrak{l}_m} \frac{\text{const}}{\mathfrak{l}_\ell + \mu \mathfrak{l}_m} \\ &\leq \frac{\text{const}}{\mathfrak{l}_m} \sum_{\mu=1}^{\text{const}/\mathfrak{l}_m} \frac{1}{\frac{\mathfrak{l}_\ell}{\mathfrak{l}_m} + \mu} \leq \frac{\text{const}}{\mathfrak{l}_m} \ln \frac{\frac{\mathfrak{l}_\ell}{\mathfrak{l}_m} + \frac{\text{const}}{\mathfrak{l}_m}}{\frac{\mathfrak{l}_\ell}{\mathfrak{l}_m}} \leq \frac{\text{const}}{\mathfrak{l}_m} \ln \left( 1 + \frac{\text{const}}{\mathfrak{l}_\ell} \right) \\ &\leq \frac{\text{const}}{\mathfrak{l}_m} \ell \leq \frac{\text{const}}{\mathfrak{l}_m \sqrt{\mathfrak{l}_\ell}} \end{aligned}$$

■

**Lemma C.4** *Let  $1 \leq \ell \leq m$  and  $\kappa' \in \mathfrak{K}_\ell$ . Let  $D$  be the disc of radius  $\epsilon$  centered at  $\boldsymbol{\tau}$ , with  $\boldsymbol{\tau} \in \mathbb{R}^2$  and  $0 \leq \epsilon \leq 2\mathfrak{l}_m$ . Let  $N$  be the number of 4-tuples  $(u_1, u_2, s_1, s_2) \in \Sigma_\ell \times \Sigma_\ell \times \Sigma_m \times \Sigma_m$  fulfilling*

$$\begin{aligned} \pi(s_1 - s_2) \cap D &\neq \emptyset \\ \pi(u_1 - u_2) \cap \pi(\kappa' - s_1) &\neq \emptyset \end{aligned} \tag{C.6}$$

a) If  $|\boldsymbol{\tau}| \geq \delta_F$ , then  $N \leq \frac{\text{const}}{\mathfrak{l}_\ell \sqrt{\mathfrak{l}_m}}$ .

b) If  $|\boldsymbol{\tau}| \leq \delta_F$ , then

$$N \leq \frac{\text{const}}{\mathfrak{l}_m \mathfrak{l}_\ell} \left[ \min \left( \ell \mathfrak{l}_\ell, \frac{1+M^m \epsilon}{M^m |\boldsymbol{\tau}|} \right) + \frac{\sqrt{\mathfrak{l}_\ell}}{|\boldsymbol{\tau}|} \left( \frac{1}{M^m} + \epsilon \right) + \mathfrak{l}_m \right]$$

**Proof:** a) By Lemma C.2.a,  $\#\{(s_1, s_2) \in \Sigma_m^2 \mid \pi(s_1 - s_2) \cap D \neq \emptyset\} \leq \frac{\text{const}}{\sqrt{\mathfrak{l}_m}}$ . For each fixed  $(s_1, s_2)$  there are at most  $\frac{\text{const}}{\mathfrak{l}_\ell}$  pairs  $(u_1, u_2)$  such that (C.6) holds. The desired bound follows.

b) As in the proof of Lemma C.3, we bound each of the three terms in

$$\begin{aligned} &\#\{(u_1, u_2, s_1, s_2) \mid (\text{C.6}) \text{ holds}\} \\ &\leq \#\{(u_1, u_2, s_1, s_2) \mid (\text{C.6}) \text{ holds, } |\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}| \leq \text{const } \mathfrak{l}_\ell\} \\ &\quad + \#\{(u_1, u_2, s_1, s_2) \mid (\text{C.6}) \text{ holds, } \text{const } \mathfrak{l}_\ell \leq |\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}| \leq \delta_F\} \\ &\quad + \#\{(u_1, u_2, s_1, s_2) \mid (\text{C.6}) \text{ holds, } \delta_F \leq |\mathbf{k}_{\kappa'} - \mathbf{k}_{s_1}|\} \end{aligned}$$

separately.

By Lemma C.2.b

$$\#\{(s_1, s_2) \in \Sigma_m^2 \mid \boldsymbol{\pi}(s_1 - s_2) \cap D \neq \emptyset\} \leq \text{const} \left[1 + \frac{1+M^m \epsilon}{M^m \mathfrak{l}_m |\boldsymbol{\tau}|}\right] \quad (\text{C.7})$$

As well, for each fixed  $s_1$  there are at most  $\text{const}$   $s_2 \in \Sigma_m$  such that  $\boldsymbol{\pi}(s_1 - s_2) \cap D \neq \emptyset$ . Hence

$$\#\{(s_1, s_2) \in \Sigma_m^2 \mid \boldsymbol{\pi}(s_1 - s_2) \cap D \neq \emptyset, |\mathbf{k}_{s_1} - \mathbf{k}_{s_2}| \leq \text{const} \mathfrak{l}_\ell\} \leq \text{const} \min \left\{1 + \frac{1+M^m \epsilon}{M^m \mathfrak{l}_m |\boldsymbol{\tau}|}, \frac{\mathfrak{l}_\ell}{\mathfrak{l}_m}\right\}$$

Also, for each fixed  $(s_1, s_2)$  there are at most  $\frac{\text{const}}{\mathfrak{l}_\ell}$  pairs  $(u_1, u_2)$  such that (C.6) holds. Hence

$$\begin{aligned} \#\{(u_1, u_2, s_1, s_2) \mid (\text{C.6}) \text{ holds}, |\mathbf{k}_{s_1} - \mathbf{k}_{s_2}| \leq \text{const} \mathfrak{l}_\ell\} &\leq \frac{\text{const}}{\mathfrak{l}_m \mathfrak{l}_\ell} \min \left\{\mathfrak{l}_m + \frac{1+M^m \epsilon}{M^m |\boldsymbol{\tau}|}, \mathfrak{l}_\ell\right\} \\ &\leq \frac{\text{const}}{\mathfrak{l}_m \mathfrak{l}_\ell} \left[\min \left\{\frac{1+M^m \epsilon}{M^m |\boldsymbol{\tau}|}, \mathfrak{l}_\ell\right\} + \mathfrak{l}_m\right] \end{aligned}$$

This gives the desired bound for the first term.

We next bound the third term. For each fixed  $(s_1, s_2)$  with  $|\mathbf{k}_{s_1} - \mathbf{k}_{s_2}| \geq \delta_F$ , there are, by Lemma C.2a, at most  $\frac{\text{const}}{\sqrt{\mathfrak{l}_\ell}}$  pairs  $(u_1, u_2)$  such that (C.6) holds. Hence

$$\begin{aligned} \#\{(u_1, u_2, s_1, s_2) \mid (\text{C.6}) \text{ holds}, \delta_F \leq |\mathbf{k}_{s_1} - \mathbf{k}_{s_2}|\} &\leq \frac{\text{const}}{\sqrt{\mathfrak{l}_\ell}} \left[1 + \frac{1+M^m \epsilon}{M^m \mathfrak{l}_m |\boldsymbol{\tau}|}\right] \\ &\leq \frac{\text{const}}{\mathfrak{l}_m \mathfrak{l}_\ell} \left[\mathfrak{l}_m \sqrt{\mathfrak{l}_\ell} + \frac{\sqrt{\mathfrak{l}_\ell}}{|\boldsymbol{\tau}|} \left(\frac{1}{M^m} + \epsilon\right)\right] \end{aligned}$$

which is smaller than the desired bound.

Finally, for the second term, we observe that, for each fixed  $(s_1, s_2)$  satisfying  $\text{const} \mathfrak{l}_\ell \leq |\mathbf{k}_{s_1} - \mathbf{k}_{s_2}| \leq \delta_F$  there are, by Lemma C.2b, with  $\epsilon = 2\mathfrak{l}_\ell$  and  $m$  replaced by  $\ell$ , at most  $\frac{\text{const}}{\mathfrak{l}_\ell |\mathbf{k}_{s_1} - \mathbf{k}_{s_2}|} \left[\frac{1}{M^\ell} + \mathfrak{l}_\ell\right] \leq \frac{\text{const}}{|\mathbf{k}_{s_1} - \mathbf{k}_{s_2}|} \leq \frac{\text{const}}{\mathfrak{l}_\ell}$  pairs  $(u_1, u_2)$  such that (C.6) holds. Hence, by (C.7) and the last argument of Lemma C.3,

$$\begin{aligned} &\#\{(u_1, u_2, s_1, s_2) \mid (\text{C.6}) \text{ holds}, \text{const} \mathfrak{l}_\ell \leq |\mathbf{k}_{s_1} - \mathbf{k}_{s_2}| \leq \delta_F\} \\ &\leq \min \left\{ \sum_{\mu=1}^{\text{const}/\mathfrak{l}_m} \frac{\text{const}}{\mathfrak{l}_\ell + \mu \mathfrak{l}_m}, \frac{\text{const}}{\mathfrak{l}_\ell} \left[1 + \frac{1+M^m \epsilon}{M^m \mathfrak{l}_m |\boldsymbol{\tau}|}\right] \right\} \\ &\leq \frac{\text{const}}{\mathfrak{l}_m \mathfrak{l}_\ell} \min \left\{ \mathfrak{l}_\ell \sum_{\mu=1}^{\text{const}/\mathfrak{l}_m} \frac{1}{\frac{\mathfrak{l}_\ell}{\mathfrak{l}_m} + \mu}, \mathfrak{l}_m + \frac{1+M^m \epsilon}{M^m |\boldsymbol{\tau}|} \right\} \\ &\leq \frac{\text{const}}{\mathfrak{l}_m \mathfrak{l}_\ell} \min \left\{ \mathfrak{l}_\ell \ell, \mathfrak{l}_m + \frac{1+M^m \epsilon}{M^m |\boldsymbol{\tau}|} \right\} \\ &\leq \frac{\text{const}}{\mathfrak{l}_m \mathfrak{l}_\ell} \left[ \mathfrak{l}_m + \min \left\{ \ell \mathfrak{l}_\ell, \frac{1+M^m \epsilon}{M^m |\boldsymbol{\tau}|} \right\} \right] \end{aligned}$$

■

# References

- [FHN] H. Fukuyama, Y. Hasegawa and O. Narikiyo, **Two-Dimensional Hubbard Model at Low Electron Density**, Journal of the Physical Society of Japan, **60** (1991), 2013–2030.
- [FKLT1] J. Feldman, H. Knörrer, D. Lehmann and E. Trubowitz, **Fermi Liquids in Two Space Dimensions**, in “*Constructive Physics*”, edited by V. Rivasseau, Springer LNP **446**, 267–299 (1995).
- [FKLT2] J. Feldman, H. Knörrer, D. Lehmann and E. Trubowitz, **A Class of Fermi Liquids**, in “*Particles and Fields '94*”, edited by G. Semenoff and L. Vinet, Springer 35-62 (1999).
- [FMST] J. Feldman, H. Knörrer, R. Sinclair and E. Trubowitz, **Superconductivity in a Repulsive Model**, Helvetica Physica Acta, **70** (1997) 154–191.
- [FKTf1] J. Feldman, H. Knörrer, E. Trubowitz, **A Two Dimensional Fermi Liquid, Part 1: Overview**, preprint.
- [FKTf2] J. Feldman, H. Knörrer, E. Trubowitz, **A Two Dimensional Fermi Liquid, Part 2: Convergence**, preprint.
- [FKTf3] J. Feldman, H. Knörrer, E. Trubowitz, **A Two Dimensional Fermi Liquid, Part 3: The Fermi Surface**, preprint.
- [FKTo1] J. Feldman, H. Knörrer, E. Trubowitz, **Single Scale Analysis of Many Fermion Systems, Part 1: Insulators**, preprint.
- [FKTo2] J. Feldman, H. Knörrer, E. Trubowitz, **Single Scale Analysis of Many Fermion Systems, Part 2: The First Scale**, preprint.
- [FKTo3] J. Feldman, H. Knörrer, E. Trubowitz, **Single Scale Analysis of Many Fermion Systems, Part 3: Sectorized Norms**, preprint.
- [FKTo4] J. Feldman, H. Knörrer, E. Trubowitz, **Single Scale Analysis of Many Fermion Systems, Part 4: Sector Counting**, preprint.
- [FKTr1] J. Feldman, H. Knörrer, E. Trubowitz, **Convergence of Perturbation Expansions in Fermionic Models, Part 1: Nonperturbative Bounds**, preprint.
- [FKTr2] J. Feldman, H. Knörrer, E. Trubowitz, **Convergence of Perturbation Expansions in Fermionic Models, Part 2: Overlapping Loops**, preprint.
- [FMRT] J. Feldman, J. Magnen, V. Rivasseau, E. Trubowitz, **Two Dimensional Many Fermion Systems as Vector Models**, Europhysics Letters, **24** (1993) 521-526.

- [N] P. Nozières, *Theory of Interacting Fermi Systems*, Benjamin (1964).
- [S] M. Salmhofer, private communication.



# Notation

## Configuration Spaces

| Symbol                                | Interpretation                                           | Reference               |
|---------------------------------------|----------------------------------------------------------|-------------------------|
| $\mathbb{M}$                          | momentum                                                 | after Definition I.3    |
| $\mathfrak{Y}$                        | momentum or position                                     | before Definition III.1 |
| $\mathfrak{Y}_\Sigma$                 | momentum or (position, sector)                           | after Definition I.3    |
| $\mathfrak{Y}_{0,\Sigma}$             | momentum                                                 | (I.2)                   |
| $\mathfrak{Y}_{1,\Sigma}$             | (position, sector)                                       | (I.2)                   |
| $\mathfrak{Y}_{2,\Sigma}$             | (momentum, sector)                                       | Definition I.5          |
| $\mathfrak{Y}_\Sigma^\uparrow$        | (momentum, spin) or (position, spin, sector)             | after Definition I.3    |
| $\mathfrak{X}_\Sigma$                 | (momentum, spin, creation/annihilation index)            | after Definition I.3    |
|                                       | or (position, spin, creation/annihilation index, sector) | after Definition I.3    |
| $\mathcal{B}^\uparrow$                | (position, spin)                                         | after Definition I.3    |
| $\check{\mathcal{B}}^\uparrow$        | (momentum, spin)                                         | after Definition I.3    |
| $\mathcal{B}$                         | (position, spin, creation/annihilation index)            | after Definition I.3    |
| $\check{\mathcal{B}}$                 | (momentum, spin, creation/annihilation index)            | after Definition I.3    |
| $\mathfrak{Y}_{\Sigma,\Sigma'}^{(4)}$ | $\mathfrak{Y}_\Sigma^2 \times \mathfrak{Y}_{\Sigma'}^2$  | (I.2)                   |
| $\mathfrak{Y}_{\ell,r}$               | $\mathfrak{Y}_{\Sigma_\ell,\Sigma_r}^{(4)}$              | Convention II.13        |
| $\mathfrak{K}_r$                      | momentum or sector                                       | Definition III.7        |

## Norms

| Norm                                                          | Characteristics                                                                           | Reference           |
|---------------------------------------------------------------|-------------------------------------------------------------------------------------------|---------------------|
| $\  \cdot \ _{1,\infty}$                                      | no derivatives, external positions only                                                   | Definition I.11     |
| $\  \cdot \ $                                                 | no derivatives, external positions and momenta                                            | Definition III.1    |
| $\  \cdot \ _{\text{bubble}}$                                 | operator norm for bubble propagators                                                      | Definition III.1    |
| $  \cdot  _{1,\Sigma}^\delta$                                 | two-legged kernel, $\delta$ derivatives, sectors                                          | Definition I.12     |
| $  \cdot  _{\Sigma,\Sigma'}^{(\delta_l, \delta_c, \delta_r)}$ | four-legged kernel, $(\delta_l, \delta_c, \delta_r)$ derivatives, sectors                 | Definition I.14     |
| $  \cdot  _{1,\Sigma}$                                        | two-legged kernel, all derivatives, sectors                                               | Definition I.15     |
| $  \cdot  _\Sigma$                                            | four-legged kernel, all derivatives, sectors                                              | Definition I.15     |
| $\  \cdot \ _{\ell,r}^{(\delta_l, \delta_c, \delta_r)}$       | $(\delta_l, \delta_c, \delta_r)$ scaled derivatives, sectors $\Sigma_\ell, \Sigma_r$      | Definition II.14    |
| $  \cdot  _{\ell,r}^{[\delta_l, \delta_c, \delta_r]}$         | $\leq (\delta_l, \delta_c, \delta_r)$ scaled derivatives, sectors $\Sigma_\ell, \Sigma_r$ | Def'ns II.14, II.16 |
| $  \cdot  _j^{[\delta]}$                                      | $\delta_l + \delta_c + \delta_r \leq \delta$ scaled derivatives, sectors $\Sigma_j$       | Def'ns II.14, II.16 |
| $  \cdot  _{\ell,r}$                                          | no derivatives, sectors $\Sigma_\ell, \Sigma_r$                                           | Definition III.6    |
| $\  \cdot \ _{\kappa_1, \kappa_2}$                            | no derivatives, specified right hand momenta/sectors                                      | Definition III.7    |

## Propagators and Ladders

| Symbol                                  | Interpretation                                                                                                              | Reference              |
|-----------------------------------------|-----------------------------------------------------------------------------------------------------------------------------|------------------------|
| $C_v^{(j)}$                             | $C_v^{(j)}(k) = \frac{\nu^{(j)}(k)}{ik_0 - e(\mathbf{k}) - v(k)}$ , single scale propagator                                 | before Definition I.17 |
| $C_v^{(\geq j)}$                        | $C_v^{(\geq j)}(k) = \frac{\nu^{(\geq j)}(k)}{ik_0 - e(\mathbf{k}) - v(k)}$ , multi scale propagator                        | before Definition I.17 |
| $\mathcal{C}(A, B)$                     | $A \otimes A^t + A \otimes B^t + B \otimes A^t$ , bubble propagator                                                         | Definition I.8         |
| $\mathcal{C}^{(j)}$                     | $\sum_{\substack{i_1, i_2 \geq 1 \\ \min(i_1, i_2) = j}} C_v^{(i_1)} \otimes C_v^{(i_2)t}$ , single scale bubble propagator | before Convention II.1 |
| $\mathcal{C}^{[i, j]}$                  | $\sum_{i \leq \ell \leq j} \mathcal{C}^{(\ell)}$ , multi scale bubble propagator                                            | before Convention II.1 |
| $\mathcal{C}_{\text{top}}^{[i, j]}$     | $\sum_{\substack{i \leq i_t \leq j \\ i_b > j}} C_v^{(i_t)} \otimes C_v^{(i_b)t}$                                           | Definition II.18       |
| $\mathcal{C}_{\text{mid}}^{[i, j]}$     | $\sum_{\substack{i \leq i_t \leq j \\ i \leq i_b \leq j}} C_v^{(i_t)}$                                                      | Definition II.18       |
| $\mathcal{C}_{\text{bot}}^{[i, j]}$     | $\sum_{\substack{i_t > j \\ i \leq i_b \leq j}} C_v^{(i_t)} \otimes C_v^{(i_b)t}$                                           | Definition II.18       |
| $\mathcal{D}_{\nu, \text{up}}^{(\ell)}$ | $\frac{1}{M^{ \nu ^\ell}} \sum_{m=\ell}^{\infty} D_{1,3}^\nu C_v^{(\ell)} \otimes C_v^{(m)t}$                               | Theorem II.20          |
| $\mathcal{D}_{\nu, \text{dn}}^{(\ell)}$ | $\frac{1}{M^{ \nu ^\ell}} \sum_{m=\ell+1}^{\infty} C_v^{(m)} \otimes D_{2,4}^\nu C_v^{(\ell)t}$                             | Theorem II.20          |
| $\mathcal{M}$                           | model particle–hole bubble propagator                                                                                       | (III.22)               |
| $\mathcal{L}_v^{(j)}(\vec{F})$          | compound particle–hole ladder                                                                                               | Definition I.19        |
| $L^{(j)}$                               | single scale compound particle–hole ladder                                                                                  | Definition II.2        |

## Scales and Sectors

| Symbol                                       | Interpretation                                                             | Reference              |
|----------------------------------------------|----------------------------------------------------------------------------|------------------------|
| $M$                                          | scale parameter, $M > 1$ , large enough                                    | Lemma I.1              |
| $\nu, \varphi$                               | used in constructing scale functions                                       | Definition I.2         |
| $\nu^{(j)}, j \geq 1$                        | partition of unity that implements scales                                  | Definition I.2         |
| $\nu^{(\geq j)}$                             | basically $\sum_{i \geq j} \nu^{(i)}$                                      | Definition I.2         |
| $\nu_0(\omega)\nu_1(\mathbf{p}, \mathbf{k})$ | factorized cutoff for model bubble propagator                              | before (III.22)        |
| $\aleph$                                     | $\frac{1}{2} < \aleph < \frac{2}{3}$ , parameter controlling sector length | before Definition I.17 |
| $\mathfrak{l}_j$                             | $\mathfrak{l}_j = \frac{1}{M^{\aleph j}}$ , sector length for scale $j$    | before Definition I.17 |
| $\chi_s, s \in \Sigma$                       | partition of unity that implements sectorization                           | before Definition I.17 |
| $\Sigma_j$                                   | set of sectors of scale $j$                                                | before Definition I.17 |
| $p_\Sigma, f_{\Sigma, \Sigma'}$              | resectorization                                                            | Definition I.17        |

## Miscellaneous

| Symbol                       | Interpretation                                                                                                                                  | Reference             |
|------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| $\text{const}$               | generic constant, independent of scale                                                                                                          |                       |
| $\text{const}$               | generic constant, independent of scale and $M$                                                                                                  |                       |
| $F$                          | Fermi curve = $\{ \mathbf{k} \in \mathbb{R}^2 \mid e(\mathbf{k}) = 0 \}$                                                                        | before Definition I.2 |
| $r_0, r_e$                   | $r_0, r_e \geq 6$ , number of derivatives controlled                                                                                            | before Definition I.2 |
| $\pi_F$                      | projection on the Fermi surface                                                                                                                 | Definition I.3        |
| $\boldsymbol{\pi}$           | $\boldsymbol{\pi}(k_0, \mathbf{k}) = \mathbf{k}$                                                                                                | Remark III.13         |
| $\langle k, x \rangle_-$     | $-k_0 x_0 + \mathbf{k}_1 \mathbf{x}_1 + \mathbf{k}_2 \mathbf{x}_2$                                                                              | Definition I.4        |
| $b_\nu$                      | $b_1 = b_4 = 0, b_1 = b_2 = 1$                                                                                                                  | Definition I.4        |
| $K^f$                        | flipped vertex                                                                                                                                  | (I.5)                 |
| $\mathbf{c}_j$               | $\sum_{\delta \in \Delta} M^{j \delta } t^\delta + \sum_{\delta \notin \Delta} \infty t^\delta$                                                 | (I.4)                 |
| $\Delta$                     | $\{ \delta \in \mathbb{N}_0 \times \mathbb{N}_0^2 \mid \delta_0 \leq r_0, \delta_1 + \delta_2 \leq r_e \}$                                      | (II.1)                |
| $\vec{\Delta}$               | $\{ \vec{\delta} = (\delta_1, \delta_c, \delta_r) \in (\mathbb{N}_0 \times \mathbb{N}_0^2)^3 \mid \delta_1 + \delta_c + \delta_r \in \Delta \}$ | (II.1)                |
| $\bullet$                    | convolution with sector sums                                                                                                                    | Definition I.8        |
| $\circ$                      | convolution without sector sums                                                                                                                 | before (III.2)        |
| $f_C, f_S$                   | charge and spin components                                                                                                                      | Lemma II.8            |
| $W_R$                        | $W_R(p, k) = W(p, k)R(p - k)$ , transfer momentum cutoff                                                                                        | Definition III.3      |
| $\mathcal{R}(d)$             | set of functions $R(t)$ that are identically one on $d$                                                                                         | Definition III.10     |
| $\mathcal{Z}, \mathcal{Z}^t$ | zero component localization operator and transpose                                                                                              | (III.18)              |
| $\widetilde{W}$              | $(\mathcal{Z} \circ W \circ \mathcal{Z}^t)(p, k) = \delta(k_0) \int d\omega W((\omega, \mathbf{0}) + p, (\omega, \mathbf{k}))$                  |                       |
| $\widetilde{W}$              | $\widetilde{W}(p, k) = \delta(k_0) \int d\omega W((\omega, \mathbf{p}), (\omega, \mathbf{k}))$                                                  | (III.20)              |
| $\bar{j}$                    | boundary between large and small transfer momentum                                                                                              | before Prop'n III.16  |